



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 07:20 am GMT

PDB ID : 5T3P
Title : Crystal structure of Human Peroxisomal coenzyme A diphosphatase NUDT7
Authors : Srikannathasan, V.; Nunez, C.A.; Tallant, C.; Siejka, P.; Mathea, S.; Kopec, J.; Elkins, J.M.; Burgess-Brown, N.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; von Delft, F.; Huber, K.
Deposited on : 2016-08-26
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	9172 (2.04-2.00)
Clashscore	122126	10355 (2.04-2.00)
Ramachandran outliers	120053	10237 (2.04-2.00)
Sidechain outliers	120020	10236 (2.04-2.00)
RSRZ outliers	108989	8961 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>7%</div> </div> </div>
1	B	236	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>6%</div> <div>7%</div> </div> </div>
1	C	236	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>6%</div> </div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5567 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisomal coenzyme A diphosphatase NUDT7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1724	1112	292	314	6			
1	B	220	Total	C	N	O	S	0	0	0
			1736	1121	292	317	6			
1	C	221	Total	C	N	O	S	0	0	0
			1753	1128	300	319	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP P0C024
B	1	SER	-	expression tag	UNP P0C024
C	1	SER	-	expression tag	UNP P0C024

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	102	Total O 102 102	0	0
3	B	113	Total O 113 113	0	0
3	C	99	Total O 99 99	0	0

- Molecule 1: Peroxisomal coenzyme A diphosphatase NUDT7



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	113.73Å 113.73Å 108.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	56.87 – 2.03 72.84 – 2.04	Depositor EDS
% Data completeness (in resolution range)	100.0 (56.87-2.03) 94.9 (72.84-2.04)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 2.05Å)	Xtriage
Refinement program	PHENIX 1.9_1682	Depositor
R, R_{free}	0.177 , 0.225 0.190 , 0.187	Depositor DCC
R_{free} test set	2603 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5567	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1766	0.54	0/2405
1	B	0.43	0/1779	0.53	0/2423
1	C	0.40	0/1796	0.49	0/2444
All	All	0.42	0/5341	0.52	0/7272

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1724	0	1707	18	0
1	B	1736	0	1718	14	0
1	C	1753	0	1737	6	0
2	A	16	0	24	3	0
2	B	12	0	18	5	0
2	C	12	0	18	1	1
3	A	102	0	0	1	0
3	B	113	0	0	1	0
3	C	99	0	0	0	1
All	All	5567	0	5222	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:ASN:HB2	2:A:301:EDO:H22	1.60	0.83
1:B:64:LYS:NZ	3:B:401:HOH:O	2.16	0.78
1:A:67:ARG:NH1	3:A:402:HOH:O	2.26	0.69
1:A:64:LYS:HB2	2:B:302:EDO:H12	1.80	0.63
1:B:66:ARG:HH11	2:B:303:EDO:H11	1.72	0.55
1:B:101:ARG:NH2	1:C:95:GLN:HE21	2.04	0.55
1:A:171:PHE:HD2	1:A:214:GLN:NE2	2.05	0.55
1:A:183:ASP:OD1	1:A:185:VAL:HG12	2.09	0.52
1:A:185:VAL:HA	1:C:14:ASN:HA	1.91	0.52
1:C:226:GLU:O	1:C:230:LYS:HG3	2.11	0.51
1:B:159:HIS:HE1	1:B:174:HIS:ND1	2.10	0.50
1:A:54:LEU:C	1:A:55:HIS:HD2	2.14	0.50
1:B:29:ILE:HD13	1:B:221:LEU:HD11	1.95	0.49
1:C:29:ILE:HG23	1:C:32:LYS:HD3	1.95	0.49
1:B:41:TYR:OH	2:B:303:EDO:H12	2.14	0.48
1:A:66:ARG:NH2	1:B:63:GLU:O	2.45	0.48
1:A:21:LYS:HB2	1:A:128:LEU:HD11	1.96	0.47
1:A:22:ALA:HA	1:A:25:ARG:NH1	2.29	0.47
1:B:161:GLN:OE1	1:B:211:PHE:HB2	2.14	0.47
1:B:64:LYS:HE3	1:B:140:ALA:O	2.14	0.47
1:B:157:GLN:HA	1:B:157:GLN:NE2	2.30	0.47
1:A:22:ALA:O	1:A:25:ARG:HG2	2.16	0.46
2:A:303:EDO:H11	1:B:64:LYS:HA	1.98	0.45
1:B:28:ASP:OD2	2:B:301:EDO:H12	2.17	0.45
1:C:67:ARG:HH21	2:C:302:EDO:H12	1.80	0.45
1:A:169:HIS:NE2	1:A:227:LEU:HD11	2.32	0.44
1:A:228:PHE:O	1:A:231:VAL:HG12	2.17	0.44
1:A:28:ASP:OD2	2:A:302:EDO:H21	2.16	0.44
1:A:163:TYR:CG	1:A:170:ARG:HD2	2.52	0.44
1:B:66:ARG:NH1	2:B:302:EDO:H21	2.33	0.43
1:A:168:GLY:O	1:A:169:HIS:HB3	2.18	0.43
1:C:14:ASN:HD21	1:C:17:LEU:HD12	1.83	0.42
1:A:29:ILE:HG13	1:A:29:ILE:H	1.78	0.41
1:A:54:LEU:C	1:A:55:HIS:CD2	2.93	0.41
1:B:29:ILE:O	1:B:32:LYS:HB3	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:303:EDO:O1	3:C:430:HOH:O[4_556]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/236 (92%)	208 (96%)	8 (4%)	1 (0%)	31	24
1	B	218/236 (92%)	208 (95%)	8 (4%)	2 (1%)	19	11
1	C	219/236 (93%)	212 (97%)	7 (3%)	0	100	100
All	All	654/708 (92%)	628 (96%)	23 (4%)	3 (0%)	31	24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	15	SER
1	A	169	HIS
1	B	32	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/208 (90%)	187 (100%)	0	100	100
1	B	189/208 (91%)	189 (100%)	0	100	100
1	C	191/208 (92%)	190 (100%)	1 (0%)	90	92

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	567/624 (91%)	566 (100%)	1 (0%)	94	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	157	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	HIS
1	B	95	GLN
1	B	157	GLN
1	C	95	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	301	-	3,3,3	0.41	0	2,2,2	0.40	0
2	EDO	A	302	-	3,3,3	0.46	0	2,2,2	0.34	0
2	EDO	A	303	-	3,3,3	0.48	0	2,2,2	0.43	0
2	EDO	A	304	-	3,3,3	0.44	0	2,2,2	0.52	0
2	EDO	B	301	-	3,3,3	0.50	0	2,2,2	0.32	0
2	EDO	B	302	-	3,3,3	0.48	0	2,2,2	0.33	0
2	EDO	B	303	-	3,3,3	0.46	0	2,2,2	0.31	0
2	EDO	C	301	-	3,3,3	0.54	0	2,2,2	0.22	0
2	EDO	C	302	-	3,3,3	0.44	0	2,2,2	0.47	0
2	EDO	C	303	-	3,3,3	0.37	0	2,2,2	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	301	-	-	0/1/1/1	0/0/0/0
2	EDO	A	302	-	-	0/1/1/1	0/0/0/0
2	EDO	A	303	-	-	0/1/1/1	0/0/0/0
2	EDO	A	304	-	-	0/1/1/1	0/0/0/0
2	EDO	B	301	-	-	0/1/1/1	0/0/0/0
2	EDO	B	302	-	-	0/1/1/1	0/0/0/0
2	EDO	B	303	-	-	0/1/1/1	0/0/0/0
2	EDO	C	301	-	-	0/1/1/1	0/0/0/0
2	EDO	C	302	-	-	0/1/1/1	0/0/0/0
2	EDO	C	303	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	EDO	1	0
2	A	302	EDO	1	0
2	A	303	EDO	1	0
2	B	301	EDO	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	302	EDO	2	0
2	B	303	EDO	2	0
2	C	302	EDO	1	0
2	C	303	EDO	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	219/236 (92%)	-0.16	5 (2%) 60 60	32, 46, 83, 120	0
1	B	220/236 (93%)	-0.17	4 (1%) 68 67	31, 46, 84, 110	0
1	C	221/236 (93%)	-0.16	2 (0%) 84 84	30, 47, 73, 106	0
All	All	660/708 (93%)	-0.16	11 (1%) 70 69	30, 47, 80, 120	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	167	LEU	3.5
1	A	169	HIS	3.2
1	B	167	LEU	2.9
1	A	116	ILE	2.8
1	C	167	LEU	2.8
1	C	29	ILE	2.4
1	A	118	THR	2.3
1	B	29	ILE	2.1
1	A	33	TYR	2.1
1	B	165	THR	2.0
1	B	30	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	C	301	4/4	0.73	0.16	58,69,76,79	0
2	EDO	B	303	4/4	0.76	0.17	72,76,77,81	0
2	EDO	A	304	4/4	0.79	0.15	70,74,77,81	0
2	EDO	B	301	4/4	0.80	0.19	59,60,66,68	0
2	EDO	B	302	4/4	0.82	0.22	49,57,62,63	0
2	EDO	C	303	4/4	0.84	0.24	61,66,66,66	0
2	EDO	A	303	4/4	0.85	0.19	46,55,55,57	0
2	EDO	A	302	4/4	0.92	0.14	68,71,72,72	0
2	EDO	C	302	4/4	0.92	0.16	62,64,64,66	0
2	EDO	A	301	4/4	0.95	0.33	44,45,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.