



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2018 – 01:54 pm GMT

PDB ID : 1TQ2
Title : Crystal Structure of IIGP1: a paradigm for interferon inducible p47 resistance GTPases
Authors : Ghosh, A.; Uthaiiah, R.; Howard, J.; Herrmann, C.; Wolf, E.
Deposited on : 2004-06-16
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MPD	B	1000	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6573 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called interferon-inducible GTPase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	0	0
			3200	2065	524	597	14			
1	B	392	Total	C	N	O	S	0	0	0
			3181	2053	520	594	14			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	412	LYS	-	CLONING ARTIFACT	UNP Q9QZ85
A	413	LEU	-	CLONING ARTIFACT	UNP Q9QZ85
A	414	GLY	-	CLONING ARTIFACT	UNP Q9QZ85
A	415	ARG	-	CLONING ARTIFACT	UNP Q9QZ85
A	416	LEU	-	CLONING ARTIFACT	UNP Q9QZ85
A	417	GLU	-	CLONING ARTIFACT	UNP Q9QZ85
A	418	ARG	-	CLONING ARTIFACT	UNP Q9QZ85
A	419	PRO	-	CLONING ARTIFACT	UNP Q9QZ85
A	420	HIS	-	CLONING ARTIFACT	UNP Q9QZ85
A	421	ARG	-	CLONING ARTIFACT	UNP Q9QZ85
A	422	ASP	-	CLONING ARTIFACT	UNP Q9QZ85
B	412	LYS	-	CLONING ARTIFACT	UNP Q9QZ85
B	413	LEU	-	CLONING ARTIFACT	UNP Q9QZ85
B	414	GLY	-	CLONING ARTIFACT	UNP Q9QZ85
B	415	ARG	-	CLONING ARTIFACT	UNP Q9QZ85
B	416	LEU	-	CLONING ARTIFACT	UNP Q9QZ85
B	417	GLU	-	CLONING ARTIFACT	UNP Q9QZ85
B	418	ARG	-	CLONING ARTIFACT	UNP Q9QZ85
B	419	PRO	-	CLONING ARTIFACT	UNP Q9QZ85
B	420	HIS	-	CLONING ARTIFACT	UNP Q9QZ85
B	421	ARG	-	CLONING ARTIFACT	UNP Q9QZ85
B	422	ASP	-	CLONING ARTIFACT	UNP Q9QZ85

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: $C_{10}H_{17}N_6O_{13}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
2	B	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			8	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	50	Total	O	0	0
			50	50		
5	B	69	Total	O	0	0
			69	69		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: interferon-inducible GTPase



• Molecule 1: interferon-inducible GTPase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.86Å 92.67Å 142.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70 19.96 – 2.70	Depositor EDS
% Data completeness (in resolution range)	93.1 (20.00-2.70) 93.4 (19.96-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.20 (at 2.71Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.222 , 0.280 0.222 , 0.278	Depositor DCC
R_{free} test set	1419 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtriage
Anisotropy	0.386	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6573	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MPD, GNP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3264	0.61	0/4404
1	B	0.41	0/3245	0.60	0/4379
All	All	0.40	0/6509	0.60	0/8783

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3200	0	3236	172	0
1	B	3181	0	3216	152	0
2	A	32	0	13	2	0
2	B	32	0	13	2	0
3	B	1	0	0	0	0
4	B	8	0	14	7	0
5	A	50	0	0	10	0
5	B	69	0	0	11	0
All	All	6573	0	6492	317	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 317 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:105:VAL:HG12	1:B:106:GLU:H	1.24	1.01
1:A:340:GLN:HG3	5:A:506:HOH:O	1.68	0.90
1:A:45:ARG:HD3	1:A:53:LEU:HB3	1.56	0.88
1:B:159:ARG:HB3	1:B:216:THR:HG21	1.57	0.86
1:A:87:ASN:HD21	1:A:100:ALA:H	1.27	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/422 (93%)	345 (88%)	31 (8%)	15 (4%)	3	8
1	B	388/422 (92%)	343 (88%)	33 (8%)	12 (3%)	4	11
All	All	779/844 (92%)	688 (88%)	64 (8%)	27 (4%)	4	9

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	GLU
1	A	203	GLU
1	A	235	VAL
1	A	302	LEU
1	B	107	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/386 (93%)	344 (96%)	15 (4%)	32	62
1	B	358/386 (93%)	342 (96%)	16 (4%)	30	60
All	All	717/772 (93%)	686 (96%)	31 (4%)	32	61

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	390	LEU
1	B	109	MET
1	B	359	ILE
1	B	101	LYS
1	B	224	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	50	ASN
1	B	118	ASN
1	B	368	GLN
1	B	94	ASN
1	B	134	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GNP	A	500	-	27,34,34	4.36	15 (55%)	30,54,54	2.63	13 (43%)
4	MPD	B	1000	-	7,7,7	0.57	0	9,10,10	0.41	0
2	GNP	B	1500	3	27,34,34	4.29	15 (55%)	30,54,54	2.61	15 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GNP	A	500	-	-	0/16/38/38	0/3/3/3
4	MPD	B	1000	-	-	0/5/5/5	0/0/0/0
2	GNP	B	1500	3	-	0/16/38/38	0/3/3/3

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	GNP	C4-N9	-9.47	1.35	1.47
2	B	1500	GNP	C4-N9	-9.05	1.35	1.47
2	B	1500	GNP	PB-O2B	-8.49	1.33	1.56
2	A	500	GNP	PB-O2B	-8.41	1.33	1.56
2	A	500	GNP	C5-C6	-6.58	1.41	1.52

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1500	GNP	O3'-C3'-C4'	-4.84	97.05	111.06
2	A	500	GNP	O3'-C3'-C4'	-4.83	97.07	111.06
2	A	500	GNP	O1G-PG-N3B	-4.59	104.92	111.79
2	B	1500	GNP	O1G-PG-N3B	-4.28	105.39	111.79
2	A	500	GNP	O6-C6-C5	-3.53	112.53	119.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	GNP	2	0
4	B	1000	MPD	7	0
2	B	1500	GNP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	395/422 (93%)	0.10	24 (6%)	21 20	25, 55, 98, 109	0
1	B	392/422 (92%)	0.01	14 (3%)	42 42	27, 52, 93, 103	0
All	All	787/844 (93%)	0.05	38 (4%)	30 28	25, 53, 96, 109	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	104	VAL	6.5
1	A	414	GLY	5.7
1	B	108	THR	5.5
1	B	105	VAL	5.0
1	A	107	VAL	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GNP	A	500	32/32	0.79	0.32	109,109,114,115	32
4	MPD	B	1000	8/8	0.82	0.31	66,68,72,73	0
2	GNP	B	1500	32/32	0.89	0.21	68,71,72,74	32
3	MG	B	1501	1/1	0.94	0.36	55,55,55,55	1

6.5 Other polymers [i](#)

There are no such residues in this entry.