



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 10, 2018 – 09:00 PM EDT

PDB ID : 2TSR  
Title : THYMIDYLATE SYNTHASE FROM RAT IN TERNARY COMPLEX  
WITH DUMP AND TOMUDEX  
Authors : Sotelo-Mundo, R.R.; Ciesla, J.; Dzik, J.M.; Rode, W.; Maley, F.; Maley, G.;  
Hardy, L.W.; Montfort, W.R.  
Deposited on : 1998-06-19  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031021  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031021


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	
1	B	307	
1	C	307	
1	D	307	

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	0	0	0
			2271	1453	394	412	12			
1	B	281	Total	C	N	O	S	0	0	0
			2271	1453	394	412	12			
1	C	281	Total	C	N	O	S	0	0	0
			2271	1453	394	412	12			
1	D	281	Total	C	N	O	S	0	0	0
			2271	1453	394	412	12			

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula:  $C_9H_{13}N_2O_8P$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	D	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is TOMUDEX (three-letter code: D16) (formula: C<sub>21</sub>H<sub>22</sub>N<sub>4</sub>O<sub>6</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			32	21	4	6	1		
3	B	1	Total	C	N	O	S	0	0
			32	21	4	6	1		
3	C	1	Total	C	N	O	S	0	0
			32	21	4	6	1		
3	D	1	Total	C	N	O	S	0	0
			32	21	4	6	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	24	Total	O	0	0
			24	24		
4	B	26	Total	O	0	0
			26	26		
4	C	9	Total	O	0	0
			9	9		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	16	Total	O	0	0
			16	16		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: THYMIDYLATE SYNTHASE




#### • Molecule 1: THYMIDYLATE SYNTHASE



#### • Molecule 1: THYMIDYLATE SYNTHASE



#### • Molecule 1: THYMIDYLATE SYNTHASE



N296	P299	T300	T301	LYS	MET	GLU	MET	ALA	VAL
------	------	------	------	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.70Å 101.60Å 140.80Å 90.00° 101.40° 90.00°	Depositor
Resolution (Å)	8.00 – 2.60 59.41 – 2.59	Depositor EDS
% Data completeness (in resolution range)	82.9 (8.00-2.60) 81.9 (59.41-2.59)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.54 (at 2.58Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.162 , 0.222 0.165 , 0.221	Depositor DCC
$R_{free}$ test set	1760 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.4	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 39.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9367	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: D16, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/2330	0.58	0/3152
1	B	0.30	0/2330	0.59	0/3152
1	C	0.31	0/2330	0.59	0/3152
1	D	0.30	0/2330	0.59	0/3152
All	All	0.30	0/9320	0.59	0/12608

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2271	0	2230	11	0
1	B	2271	0	2230	9	0
1	C	2271	0	2230	10	0
1	D	2271	0	2230	9	0
2	A	20	0	11	0	0
2	B	20	0	11	2	0
2	C	20	0	11	2	0
2	D	20	0	11	3	0
3	A	32	0	20	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	32	0	20	1	0
3	C	32	0	20	1	0
3	D	32	0	20	1	0
4	A	24	0	0	0	0
4	B	26	0	0	0	0
4	C	9	0	0	1	0
4	D	16	0	0	0	0
All	All	9367	0	9044	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ARG:HH11	1:B:170:ARG:NH2	1.85	0.74
1:A:68:LEU:HD12	1:A:218:PRO:HB3	1.71	0.72
1:D:190:HIS:HB3	1:D:206:LEU:HD11	1.73	0.71
1:D:212:ASP:HB2	2:D:608:UMP:H2''	1.76	0.67
1:A:186:LEU:HD12	1:A:187:PRO:HD2	1.77	0.66
1:C:190:HIS:HB2	1:C:206:LEU:HD11	1.77	0.66
2:D:608:UMP:H1'	3:D:609:D16:C4	2.26	0.66
1:A:44:ARG:HH11	1:B:170:ARG:HH21	1.44	0.63
1:B:21:GLN:HG3	1:B:25:LEU:HD12	1.82	0.62
1:A:190:HIS:HB3	1:A:206:LEU:HD11	1.83	0.61
2:B:408:UMP:H1'	3:B:409:D16:C4	2.31	0.61
2:C:508:UMP:H1'	3:C:509:D16:C4	2.34	0.57
1:C:254:ASN:ND2	1:C:254:ASN:H	2.06	0.53
1:D:68:LEU:HD22	1:D:218:PRO:HB3	1.90	0.53
1:A:102:ILE:HG13	1:A:103:TRP:CD1	2.46	0.51
2:C:508:UMP:O2	2:C:508:UMP:H2'	2.12	0.49
1:D:268:ARG:HD2	1:D:296:ASN:O	2.15	0.46
1:D:155:LEU:O	1:D:158:VAL:HG22	2.16	0.45
1:B:141:LYS:HG2	1:B:142:ASP:N	2.31	0.45
1:C:257:GLU:HB2	1:C:258:PRO:HD3	1.98	0.45
2:D:608:UMP:O2	2:D:608:UMP:H2'	2.16	0.44
1:C:163:LYS:HG3	1:C:235:THR:HG22	2.00	0.44
1:A:136:PHE:CE1	1:B:178:PRO:HD2	2.54	0.43
1:C:178:PRO:HD2	1:D:136:PHE:CE1	2.53	0.43
1:C:155:LEU:O	1:C:158:VAL:HG22	2.19	0.42
1:C:39:PHE:HD2	4:C:511:HOH:O	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:266:GLU:HA	1:D:267:PRO:HD3	1.90	0.42
1:A:153:ASP:O	1:A:157:LYS:HB2	2.19	0.42
1:B:125:LEU:HD13	1:B:186:LEU:HD21	2.01	0.42
1:B:212:ASP:HB2	2:B:408:UMP:H2"	2.02	0.42
1:B:268:ARG:HD2	1:B:296:ASN:O	2.18	0.42
1:C:254:ASN:N	1:C:254:ASN:HD22	2.18	0.41
1:C:201:GLU:HA	1:C:238:GLN:O	2.20	0.41
1:D:72:ARG:O	1:D:300:THR:HG23	2.21	0.41
1:A:109:ARG:NH2	1:A:120:ARG:O	2.54	0.41
1:B:24:GLU:HG3	1:B:68:LEU:HD22	2.03	0.41
1:C:187:PRO:HA	1:C:188:PRO:HD3	1.90	0.40
1:D:265:ARG:NH2	1:D:299:PRO:O	2.54	0.40
1:A:65:PHE:HA	1:A:66:PRO:HD3	1.84	0.40
1:A:187:PRO:HA	1:A:188:PRO:HD3	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/307 (91%)	264 (95%)	14 (5%)	1 (0%)	36	60
1	B	279/307 (91%)	267 (96%)	11 (4%)	1 (0%)	36	60
1	C	279/307 (91%)	264 (95%)	13 (5%)	2 (1%)	24	46
1	D	279/307 (91%)	268 (96%)	11 (4%)	0	100	100
All	All	1116/1228 (91%)	1063 (95%)	49 (4%)	4 (0%)	36	60

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	129	TYR

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	129	TYR
1	B	129	TYR
1	C	74	PHE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/266 (92%)	239 (98%)	6 (2%)	52	77
1	B	245/266 (92%)	228 (93%)	17 (7%)	17	34
1	C	245/266 (92%)	230 (94%)	15 (6%)	20	41
1	D	245/266 (92%)	237 (97%)	8 (3%)	41	68
All	All	980/1064 (92%)	934 (95%)	46 (5%)	29	54

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	GLU
1	A	78	VAL
1	A	157	LYS
1	A	192	LEU
1	A	266	GLU
1	A	268	ARG
1	B	40	LYS
1	B	44	ARG
1	B	63	ASP
1	B	93	LYS
1	B	98	LYS
1	B	101	ARG
1	B	110	ASP
1	B	145	SER
1	B	181	LEU
1	B	190	HIS
1	B	198	VAL
1	B	208	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	264	GLN
1	B	266	GLU
1	B	277	ARG
1	B	292	ILE
1	B	300	THR
1	C	63	ASP
1	C	68	LEU
1	C	72	ARG
1	C	79	LEU
1	C	109	ARG
1	C	173	MET
1	C	181	LEU
1	C	183	LEU
1	C	186	LEU
1	C	215	LEU
1	C	238	GLN
1	C	254	ASN
1	C	274	ARG
1	C	279	VAL
1	C	301	ILE
1	D	63	ASP
1	D	98	LYS
1	D	101	ARG
1	D	179	LYS
1	D	190	HIS
1	D	274	ARG
1	D	277	ARG
1	D	300	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	205	GLN
1	B	56	GLN
1	B	205	GLN
1	B	298	HIS
1	C	56	GLN
1	C	205	GLN
1	C	254	ASN
1	D	56	GLN
1	D	165	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	205	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	UMP	A	308	-	17,21,21	1.14	2 (11%)	23,31,31	3.29	2 (8%)
3	D16	A	309	-	23,34,34	3.18	9 (39%)	26,48,48	3.55	10 (38%)
2	UMP	B	408	-	17,21,21	1.13	2 (11%)	23,31,31	3.35	3 (13%)
3	D16	B	409	-	23,34,34	3.08	10 (43%)	26,48,48	3.53	10 (38%)
2	UMP	C	508	-	17,21,21	1.14	2 (11%)	23,31,31	3.37	3 (13%)
3	D16	C	509	-	23,34,34	3.02	9 (39%)	26,48,48	3.53	10 (38%)
2	UMP	D	608	-	17,21,21	1.15	2 (11%)	23,31,31	3.39	3 (13%)
3	D16	D	609	-	23,34,34	2.93	10 (43%)	26,48,48	3.51	10 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMP	A	308	-	-	0/6/22/22	0/2/2/2
3	D16	A	309	-	-	0/13/25/25	0/3/3/3
2	UMP	B	408	-	-	0/6/22/22	0/2/2/2
3	D16	B	409	-	-	0/13/25/25	0/3/3/3
2	UMP	C	508	-	-	0/6/22/22	0/2/2/2
3	D16	C	509	-	-	0/13/25/25	0/3/3/3
2	UMP	D	608	-	-	0/6/22/22	0/2/2/2
3	D16	D	609	-	-	0/13/25/25	0/3/3/3

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	508	UMP	C6-C5	-2.14	1.33	1.38
2	D	608	UMP	C6-C5	-2.00	1.33	1.38
2	A	308	UMP	C6-N1	2.04	1.38	1.35
2	B	408	UMP	C6-N1	2.06	1.38	1.35
3	D	609	D16	C-N	2.13	1.38	1.34
3	B	409	D16	C-N	2.23	1.39	1.34
3	B	409	D16	C5-C6	2.73	1.43	1.37
3	B	409	D16	C7-C6	2.77	1.44	1.38
3	D	609	D16	C7-C6	2.81	1.44	1.38
3	C	509	D16	C7-C6	2.88	1.44	1.38
3	A	309	D16	C5-C6	2.96	1.44	1.37
3	A	309	D16	C7-C6	3.05	1.45	1.38
3	D	609	D16	C8-C7	3.10	1.43	1.36
3	C	509	D16	C5-C6	3.16	1.44	1.37
3	D	609	D16	C5-C6	3.16	1.44	1.37
2	B	408	UMP	C4-N3	3.27	1.38	1.33
2	C	508	UMP	C4-N3	3.28	1.38	1.33
3	C	509	D16	C8-C7	3.34	1.43	1.36
3	A	309	D16	C8-C7	3.36	1.43	1.36
2	A	308	UMP	C4-N3	3.36	1.39	1.33
2	D	608	UMP	C4-N3	3.37	1.39	1.33
3	B	409	D16	C8-C7	3.54	1.44	1.36
3	B	409	D16	C9-C6	3.60	1.57	1.51
3	C	509	D16	C9-C6	4.35	1.59	1.51
3	D	609	D16	C9-C6	4.35	1.59	1.51
3	A	309	D16	C9-C6	4.48	1.59	1.51
3	D	609	D16	C2-N3	4.57	1.42	1.34
3	C	509	D16	CB-CA	4.59	1.59	1.53
3	D	609	D16	CB-CA	4.70	1.59	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	509	D16	C2-N3	4.94	1.43	1.34
3	A	309	D16	C2-N3	4.95	1.43	1.34
3	B	409	D16	C2-N3	4.99	1.43	1.34
3	D	609	D16	C4-N3	5.13	1.42	1.33
3	A	309	D16	C4-N3	5.40	1.42	1.33
3	C	509	D16	C4-N3	5.41	1.42	1.33
3	D	609	D16	C2-N1	5.44	1.43	1.34
3	B	409	D16	C4-N3	5.63	1.43	1.33
3	B	409	D16	CB-CA	5.63	1.60	1.53
3	C	509	D16	C2-N1	5.75	1.44	1.34
3	A	309	D16	CB-CA	5.80	1.60	1.53
3	D	609	D16	C9-N10	5.91	1.56	1.46
3	B	409	D16	C2-N1	5.93	1.44	1.34
3	B	409	D16	C9-N10	5.97	1.56	1.46
3	A	309	D16	C2-N1	5.99	1.44	1.34
3	C	509	D16	C9-N10	6.14	1.56	1.46
3	A	309	D16	C9-N10	6.49	1.57	1.46

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	409	D16	C4A-C4-N3	-12.54	115.78	124.45
3	D	609	D16	C4A-C4-N3	-12.23	116.00	124.45
3	A	309	D16	C4A-C4-N3	-12.17	116.03	124.45
3	C	509	D16	C4A-C4-N3	-12.03	116.13	124.45
3	A	309	D16	C4A-C8A-N1	-8.05	119.08	123.67
3	B	409	D16	C4A-C8A-N1	-8.02	119.10	123.67
3	C	509	D16	C4A-C8A-N1	-7.83	119.20	123.67
3	D	609	D16	C4A-C8A-N1	-7.58	119.35	123.67
2	A	308	UMP	C5-C4-N3	-3.68	114.63	123.17
2	C	508	UMP	C5-C4-N3	-3.64	114.72	123.17
2	D	608	UMP	C5-C4-N3	-3.63	114.74	123.17
2	B	408	UMP	C5-C4-N3	-3.56	114.91	123.17
3	A	309	D16	CA-N-C	-3.55	117.51	122.25
3	A	309	D16	N1-C2-N3	-3.35	118.89	125.61
3	D	609	D16	N1-C2-N3	-3.34	118.93	125.61
3	C	509	D16	N1-C2-N3	-3.33	118.94	125.61
3	B	409	D16	N1-C2-N3	-3.27	119.07	125.61
3	B	409	D16	CA-N-C	-3.22	117.94	122.25
3	C	509	D16	CA-N-C	-3.02	118.22	122.25
3	D	609	D16	CA-N-C	-2.75	118.58	122.25
3	D	609	D16	C4-C4A-C8A	2.14	120.44	118.57

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	408	UMP	O4'-C1'-C2'	2.20	110.45	106.25
3	B	409	D16	CM2-C2-N1	2.25	120.95	117.21
3	C	509	D16	C4-C4A-C8A	2.26	120.54	118.57
3	D	609	D16	CM2-C2-N1	2.28	121.01	117.21
3	B	409	D16	CB-CA-N	2.29	113.55	110.19
3	C	509	D16	CM2-C2-N1	2.32	121.08	117.21
3	C	509	D16	CB-CA-N	2.38	113.68	110.19
3	A	309	D16	CM2-C2-N1	2.39	121.19	117.21
3	B	409	D16	C6-C9-N10	2.53	116.46	113.04
2	C	508	UMP	C2'-C1'-N1	2.55	120.21	114.27
3	A	309	D16	C4-C4A-C8A	2.57	120.81	118.57
3	D	609	D16	CB-CA-N	2.67	114.10	110.19
2	D	608	UMP	C2'-C1'-N1	2.73	120.64	114.27
3	B	409	D16	C4-C4A-C8A	2.79	121.01	118.57
3	A	309	D16	CB-CA-N	2.83	114.34	110.19
3	B	409	D16	CB-CA-CT	3.55	117.39	112.11
3	A	309	D16	C6-C9-N10	3.58	117.88	113.04
3	C	509	D16	CB-CA-CT	3.65	117.54	112.11
3	D	609	D16	CB-CA-CT	3.75	117.69	112.11
3	A	309	D16	CB-CA-CT	3.88	117.88	112.11
3	D	609	D16	C6-C9-N10	4.27	118.80	113.04
3	A	309	D16	CG-CB-CA	4.55	122.27	113.19
3	C	509	D16	C6-C9-N10	4.66	119.33	113.04
3	D	609	D16	CG-CB-CA	4.90	122.96	113.19
3	B	409	D16	CG-CB-CA	4.93	123.03	113.19
3	C	509	D16	CG-CB-CA	5.13	123.44	113.19
2	D	608	UMP	C4-N3-C2	15.14	127.17	114.14
2	B	408	UMP	C4-N3-C2	15.15	127.17	114.14
2	A	308	UMP	C4-N3-C2	15.17	127.19	114.14
2	C	508	UMP	C4-N3-C2	15.21	127.23	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	408	UMP	2	0
3	B	409	D16	1	0
2	C	508	UMP	2	0
3	C	509	D16	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	608	UMP	3	0
3	D	609	D16	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	281/307 (91%)	-0.68	0 <b>100</b> <b>100</b>	10, 27, 61, 89	0
1	B	281/307 (91%)	-0.55	4 (1%) 75 71	11, 26, 63, 97	0
1	C	281/307 (91%)	-0.35	6 (2%) 63 58	15, 40, 75, 95	0
1	D	281/307 (91%)	-0.60	4 (1%) 75 71	17, 33, 62, 96	0
All	All	1124/1228 (91%)	-0.55	14 (1%) 79 75	10, 32, 67, 97	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	43	ASP	3.8
1	C	43	ASP	3.5
1	B	44	ARG	3.1
1	D	44	ARG	3.1
1	C	44	ARG	2.5
1	C	45	THR	2.5
1	D	21	GLN	2.4
1	C	253	LEU	2.4
1	D	43	ASP	2.3
1	C	301	ILE	2.2
1	B	45	THR	2.2
1	C	21	GLN	2.1
1	B	47	THR	2.0
1	D	45	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	D16	B	409	32/32	0.88	0.24	60,71,85,89	0
3	D16	A	309	32/32	0.89	0.25	65,72,96,99	0
2	UMP	D	608	20/20	0.91	0.23	46,53,64,65	0
3	D16	D	609	32/32	0.92	0.21	48,59,72,72	0
2	UMP	B	408	20/20	0.92	0.20	43,67,70,70	0
2	UMP	A	308	20/20	0.94	0.17	33,50,63,67	0
2	UMP	C	508	20/20	0.94	0.16	40,52,63,64	0
3	D16	C	509	32/32	0.95	0.22	47,54,82,86	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.