



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 12:50 pm GMT

PDB ID : 3TSQ
Title : Crystal structure of E. coli HypF with ATP and Carbamoyl phosphate
Authors : Petkun, S.; Shi, R.; Li, Y.; Cygler, M.
Deposited on : 2011-09-13
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3481 (2.40-2.40)
Clashscore	122126	3956 (2.40-2.40)
Ramachandran outliers	120053	3897 (2.40-2.40)
Sidechain outliers	120020	3898 (2.40-2.40)
RSRZ outliers	108989	3386 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	658	 83% 14% ..

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulatory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	647	Total	C	N	O	S	0	1	0
			4935	3118	886	893	38			


There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	GLY	-	EXPRESSION TAG	UNP Q7ABC4
A	91	SER	-	EXPRESSION TAG	UNP Q7ABC4
A	571	ALA	GLN	CONFLICT	UNP Q7ABC4
A	572	ALA	GLN	CONFLICT	UNP Q7ABC4
A	573	ALA	GLN	CONFLICT	UNP Q7ABC4

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		

- Molecule 3 is 5'-O-[(S)-(carbamoyloxy)(hydroxy)phosphoryl]adenosine (three-letter code: CA0) (formula: C₁₁H₁₅N₆O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			26	11	6	8	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	378	Total	O	0	0
			378	378		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional regulatory protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.53Å 78.29Å 200.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.41 – 2.40 35.73 – 2.40	Depositor EDS
% Data completeness (in resolution range)	91.8 (100.41-2.40) 91.8 (35.73-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	13.36 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.154 , 0.233 0.154 , 0.233	Depositor DCC
R_{free} test set	1351 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.594	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5343	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA0, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	5/5060 (0.1%)	0.90	9/6909 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	525	CYS	CB-SG	-7.34	1.69	1.82
1	A	406	CYS	CB-SG	-6.96	1.70	1.82
1	A	296	GLU	CG-CD	5.94	1.60	1.51
1	A	414	VAL	CB-CG1	-5.74	1.40	1.52
1	A	297	VAL	CB-CG2	-5.20	1.42	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	707	ARG	NE-CZ-NH2	-8.26	116.17	120.30
1	A	707	ARG	NE-CZ-NH1	6.21	123.40	120.30
1	A	432	ARG	NE-CZ-NH1	-6.03	117.28	120.30
1	A	741	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	A	254	LEU	CB-CG-CD1	-5.66	101.38	111.00
1	A	705	ARG	NE-CZ-NH1	5.43	123.02	120.30
1	A	432	ARG	NE-CZ-NH2	5.24	122.92	120.30
1	A	705	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	353	ARG	NE-CZ-NH1	5.07	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4935	0	4849	60	0
2	A	3	0	0	0	0
3	A	26	0	14	1	0
4	A	1	0	0	0	0
5	A	378	0	0	15	0
All	All	5343	0	4863	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:443:ASN:HB3	5:A:983:HOH:O	1.77	0.84
1:A:640:ARG:HD2	1:A:679:LEU:HD12	1.60	0.84
1:A:212:MET:HE1	5:A:1031:HOH:O	1.82	0.79
1:A:406:CYS:SG	1:A:413:VAL:CG2	2.72	0.77
1:A:449:VAL:HG13	1:A:470:THR:HG22	1.67	0.75
1:A:449:VAL:CG1	1:A:470:THR:HG22	2.15	0.75
1:A:285:GLU:HB2	5:A:4:HOH:O	1.85	0.75
1:A:153:MET:CE	1:A:353:ARG:HG3	2.20	0.71
1:A:399:ALA:HB2	1:A:730:LEU:HD22	1.75	0.68
1:A:406:CYS:SG	1:A:413:VAL:HG23	2.33	0.67
1:A:634:HIS:HD2	1:A:636:VAL:H	1.43	0.66
1:A:451:HIS:HD2	1:A:460[A]:CYS:SG	2.18	0.66
1:A:153:MET:HE3	1:A:353:ARG:HG3	1.77	0.65
1:A:424:ASP:O	1:A:427:ILE:HG22	1.97	0.64
1:A:308:HIS:HD2	5:A:801:HOH:O	1.82	0.62
1:A:443:ASN:CB	5:A:983:HOH:O	2.46	0.58
1:A:369:GLU:OE2	1:A:705:ARG:NH2	2.38	0.57
1:A:153:MET:HE2	1:A:156:PHE:CD2	2.40	0.56
1:A:640:ARG:CD	1:A:679:LEU:HD12	2.33	0.56
1:A:153:MET:HE1	1:A:353:ARG:HG3	1.88	0.56
1:A:562:GLN:NE2	5:A:20:HOH:O	2.37	0.56
1:A:472:THR:HG22	5:A:43:HOH:O	2.05	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:748:CA0:O2A	3:A:748:CA0:O1B	2.26	0.53
1:A:308:HIS:HE1	5:A:751:HOH:O	1.91	0.53
1:A:640:ARG:HB2	5:A:1034:HOH:O	2.07	0.53
1:A:467:ASN:ND2	5:A:919:HOH:O	2.44	0.51
1:A:280:LYS:HE3	1:A:294:LEU:O	2.11	0.51
1:A:652:GLN:NE2	5:A:1019:HOH:O	2.28	0.51
1:A:451:HIS:CD2	1:A:460[A]:CYS:SG	3.03	0.51
1:A:396:CYS:HB2	1:A:406:CYS:HB3	1.93	0.50
1:A:518:LEU:HD23	1:A:527:HIS:HA	1.94	0.50
1:A:372:ARG:NH1	5:A:1045:HOH:O	2.34	0.49
1:A:351:HIS:HD2	1:A:353:ARG:H	1.59	0.49
1:A:640:ARG:HG3	1:A:679:LEU:CD1	2.43	0.49
1:A:399:ALA:HB2	1:A:730:LEU:CD2	2.43	0.48
1:A:406:CYS:SG	1:A:413:VAL:HG21	2.52	0.48
1:A:397:LEU:HD12	1:A:463:ALA:HB2	1.97	0.47
1:A:397:LEU:CD1	1:A:463:ALA:HB2	2.44	0.47
1:A:694:PHE:O	1:A:720:PRO:HD3	2.15	0.47
1:A:407:LEU:HB2	1:A:414:VAL:HG12	1.97	0.46
1:A:360:ASP:O	1:A:374:SER:HB2	2.16	0.46
1:A:419:LEU:O	1:A:427:ILE:HD11	2.16	0.45
1:A:723:LEU:HD11	1:A:732:LEU:HD22	1.98	0.45
1:A:393:PRO:HB2	1:A:446:PRO:HA	1.98	0.45
1:A:335:GLU:HG3	5:A:753:HOH:O	2.15	0.45
1:A:523:ARG:NE	1:A:746:GLU:OE2	2.45	0.45
1:A:634:HIS:HE1	1:A:670:ASP:OD2	2.00	0.44
1:A:425:ASP:HB2	5:A:996:HOH:O	2.17	0.43
1:A:224:PHE:HZ	1:A:334:ASN:HD22	1.67	0.43
1:A:451:HIS:O	1:A:472:THR:HA	2.19	0.42
1:A:140:ILE:HB	1:A:152:VAL:HG22	2.01	0.42
1:A:126:TYR:HA	1:A:127:PRO:HD3	1.90	0.42
1:A:138:PHE:CE1	1:A:357:GLN:HG3	2.55	0.41
1:A:252:VAL:HG11	1:A:286:LEU:HD21	2.03	0.41
1:A:195:GLU:N	1:A:195:GLU:CD	2.74	0.41
1:A:640:ARG:CG	1:A:679:LEU:HD12	2.49	0.41
1:A:714:ASP:N	1:A:714:ASP:OD1	2.49	0.41
1:A:372:ARG:CD	5:A:1045:HOH:O	2.69	0.41
1:A:168:ASP:HA	1:A:169:PRO:HD3	1.85	0.40
1:A:230:ALA:HB3	1:A:317:PRO:HB2	2.04	0.40
1:A:631:GLY:HA2	1:A:666:TRP:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	646/658 (98%)	620 (96%)	22 (3%)	4 (1%)	27	39

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	514	GLY
1	A	642	ASP
1	A	539	GLY
1	A	503	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	509/514 (99%)	487 (96%)	22 (4%)	32	49

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	GLN
1	A	102	ILE
1	A	183	GLU
1	A	195	GLU
1	A	266	GLN
1	A	278	VAL
1	A	288	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	335	GLU
1	A	358	ARG
1	A	371	LEU
1	A	421	ASP
1	A	425	ASP
1	A	472	THR
1	A	504	ILE
1	A	506	MET
1	A	566	GLU
1	A	615	GLU
1	A	635	PRO
1	A	642	ASP
1	A	702	ARG
1	A	718	LEU
1	A	747	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	A	193	HIS
1	A	204	GLN
1	A	214	ASN
1	A	295	ASN
1	A	308	HIS
1	A	324	ASN
1	A	334	ASN
1	A	336	GLN
1	A	351	HIS
1	A	430	GLN
1	A	443	ASN
1	A	451	HIS
1	A	587	ASN
1	A	634	HIS
1	A	700	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CA0	A	748	-	24,28,28	2.10	4 (16%)	24,42,42	1.83	6 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CA0	A	748	-	-	0/8/31/31	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	748	CA0	O4'-C1'	4.00	1.46	1.41
3	A	748	CA0	C5-C4	4.03	1.49	1.40
3	A	748	CA0	C2-N3	4.26	1.39	1.32
3	A	748	CA0	CB-N2B	6.43	1.45	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	748	CA0	N3-C2-N1	-5.07	124.52	128.86
3	A	748	CA0	C4-C5-N7	-3.42	106.10	109.41
3	A	748	CA0	O3A-PA-O2A	-2.62	101.06	108.71
3	A	748	CA0	O1A-PA-O3A	2.08	110.67	103.72
3	A	748	CA0	O1A-PA-O2A	2.41	124.38	112.14
3	A	748	CA0	C2'-C3'-C4'	2.54	107.50	102.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	748	CA0	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	647/658 (98%)	-0.65	3 (0%) 90 89	9, 22, 42, 55	2 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	642	ASP	2.9
1	A	101	GLN	2.5
1	A	193	HIS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CAO	A	748	26/26	0.86	0.16	42,56,69,69	0
2	ZN	A	3	1/1	0.95	0.04	58,58,58,58	0
4	MG	A	749	1/1	0.97	0.07	20,20,20,20	0
2	ZN	A	2	1/1	0.99	0.04	55,55,55,55	0
2	ZN	A	1	1/1	0.99	0.04	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.