



# Full wwPDB NMR Structure Validation Report ⓘ

Feb 16, 2018 – 10:38 pm GMT

PDB ID : 1U3M  
Title : NMR structure of the chicken prion protein fragment 128-242  
Authors : Lysek, D.A.; Calzolari, L.; Guntert, P.; Wuthrich, K.  
Deposited on : 2004-07-22

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk30686
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30686


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 80%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136279	12091
Ramachandran outliers	132675	10835
Sidechain outliers	132484	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	117	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 8 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:135-A:199, A:212-A:240 (94)	0.52	8

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 8, 9, 16
2	11, 12
3	10, 13
4	2, 19
Single-model clusters	14; 15; 17; 18; 20

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1819 atoms, of which 884 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called prion-like protein.

Mol	Chain	Residues	Atoms						Trace
1	A	117	Total	C	H	N	O	S	0
			1819	585	884	163	180	7	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	126	GLY	-	CLONING ARTIFACT	UNP P27177
A	127	SER	-	CLONING ARTIFACT	UNP P27177

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: prion-like protein




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: prion-like protein



#### 4.2.2 Score per residue for model 2


- Molecule 1: prion-like protein






### 4.2.3 Score per residue for model 3

- Molecule 1: prion-like protein




### 4.2.4 Score per residue for model 4

- Molecule 1: prion-like protein



### 4.2.5 Score per residue for model 5

- Molecule 1: prion-like protein



### 4.2.6 Score per residue for model 6


- Molecule 1: prion-like protein






#### 4.2.7 Score per residue for model 7

- Molecule 1: prion-like protein




#### 4.2.8 Score per residue for model 8 (medoid)

- Molecule 1: prion-like protein




#### 4.2.9 Score per residue for model 9

- Molecule 1: prion-like protein




#### 4.2.10 Score per residue for model 10

- Molecule 1: prion-like protein




#### 4.2.11 Score per residue for model 11

- Molecule 1: prion-like protein




#### 4.2.12 Score per residue for model 12

- Molecule 1: prion-like protein




#### 4.2.13 Score per residue for model 13

- Molecule 1: prion-like protein




## 4.2.14 Score per residue for model 14

- Molecule 1: prion-like protein




## 4.2.15 Score per residue for model 15

- Molecule 1: prion-like protein




## 4.2.16 Score per residue for model 16

- Molecule 1: prion-like protein




## 4.2.17 Score per residue for model 17

- Molecule 1: prion-like protein




### 4.2.18 Score per residue for model 18

- Molecule 1: prion-like protein




### 4.2.19 Score per residue for model 19

- Molecule 1: prion-like protein



### 4.2.20 Score per residue for model 20

- Molecule 1: prion-like protein



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	6.0
DYANA	refinement	6.0

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6269
Number of chemical shift lists	1
Total number of shifts	1294
Number of shifts mapped to atoms	1294
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	80%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.71±0.01	0±0/819 (0.0±0.0%)	1.16±0.05	2±2/1113 (0.2±0.1%)
All	All	0.71	0/16380 (0.0%)	1.16	34/22260 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.1±0.9
All	All	0	43

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	223	VAL	CA-CB-CG2	8.61	123.81	110.90	19	1
1	A	163	ARG	NE-CZ-NH2	-7.46	116.57	120.30	15	1
1	A	226	VAL	CA-CB-CG2	6.68	120.92	110.90	16	1
1	A	228	ARG	CD-NE-CZ	6.66	132.93	123.60	9	1
1	A	154	TYR	CB-CG-CD2	-6.51	117.09	121.00	18	3
1	A	168	VAL	CG1-CB-CG2	-6.22	100.95	110.90	7	1
1	A	191	VAL	CA-CB-CG2	5.96	119.84	110.90	4	4
1	A	164	TYR	CB-CG-CD2	-5.93	117.44	121.00	15	1
1	A	228	ARG	NE-CZ-NH1	5.77	123.18	120.30	9	1
1	A	185	ASP	CB-CG-OD2	-5.76	113.11	118.30	20	1
1	A	146	TYR	CB-CG-CD2	-5.65	117.61	121.00	13	1
1	A	192	THR	CA-CB-CG2	5.61	120.26	112.40	6	1
1	A	239	ARG	NE-CZ-NH2	-5.57	117.52	120.30	3	3
1	A	177	VAL	CG1-CB-CG2	5.49	119.68	110.90	13	1
1	A	177	VAL	CB-CA-C	5.49	121.83	111.40	13	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	187	PHE	CB-CG-CD2	-5.46	116.98	120.80	19	1
1	A	139	ARG	NE-CZ-NH2	-5.20	117.70	120.30	10	1
1	A	238	TYR	CB-CG-CD2	-5.18	117.89	121.00	15	1
1	A	191	VAL	CA-CB-CG1	5.18	118.66	110.90	8	1
1	A	150	ARG	CD-NE-CZ	5.15	130.81	123.60	6	1
1	A	163	ARG	NE-CZ-NH1	5.13	122.87	120.30	20	1
1	A	167	ARG	NE-CZ-NH1	5.13	122.87	120.30	3	1
1	A	226	VAL	CA-CB-CG1	5.12	118.58	110.90	9	3
1	A	139	ARG	NE-CZ-NH1	5.08	122.84	120.30	10	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	155	ARG	Sidechain	4
1	A	228	ARG	Sidechain	4
1	A	171	ARG	Sidechain	3
1	A	239	ARG	Sidechain	3
1	A	236	ARG	Sidechain	3
1	A	194	TYR	Sidechain	3
1	A	163	ARG	Sidechain	3
1	A	173	TYR	Sidechain	3
1	A	146	TYR	Sidechain	2
1	A	150	ARG	Sidechain	2
1	A	170	TYR	Sidechain	2
1	A	169	TYR	Sidechain	2
1	A	154	TYR	Sidechain	2
1	A	167	ARG	Sidechain	2
1	A	235	TYR	Sidechain	1
1	A	164	TYR	Sidechain	1
1	A	142	SER	Peptide	1
1	A	238	TYR	Sidechain	1
1	A	135	TYR	Sidechain	1

## 6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	796	741	741	3±1
All	All	15920	14820	14820	51

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:187:PHE:CE1	1:A:224:THR:HG23	0.61	2.30	4	3
1:A:187:PHE:CZ	1:A:224:THR:HG22	0.60	2.31	19	1
1:A:226:VAL:HG12	1:A:230:MET:CE	0.56	2.31	17	2
1:A:220:ASN:O	1:A:224:THR:HG23	0.56	2.00	17	9
1:A:177:VAL:HG22	1:A:181:VAL:HG21	0.54	1.78	3	1
1:A:148:PHE:CE2	1:A:156:TRP:CE3	0.53	2.96	2	1
1:A:191:VAL:HG23	1:A:199:ALA:CB	0.52	2.35	9	1
1:A:228:ARG:O	1:A:232:VAL:HG23	0.50	2.06	9	3
1:A:183:VAL:HG23	1:A:231:CYS:HB2	0.49	1.84	18	1
1:A:187:PHE:CE1	1:A:191:VAL:HG11	0.48	2.43	19	1
1:A:226:VAL:HG12	1:A:230:MET:HE2	0.48	1.86	16	1
1:A:187:PHE:CE1	1:A:224:THR:HG22	0.47	2.44	19	1
1:A:164:TYR:CE1	1:A:226:VAL:HG21	0.46	2.45	19	1
1:A:220:ASN:HA	1:A:223:VAL:HG22	0.46	1.86	19	1
1:A:212:ASN:O	1:A:216:VAL:HG13	0.46	2.11	8	1
1:A:177:VAL:HG22	1:A:181:VAL:HG11	0.45	1.87	18	1
1:A:218:MET:HE2	1:A:222:VAL:HG23	0.44	1.90	16	1
1:A:168:VAL:HG21	1:A:230:MET:CB	0.44	2.43	15	1
1:A:187:PHE:HA	1:A:227:ILE:CG2	0.43	2.43	17	3
1:A:140:VAL:HG13	1:A:166:ASN:ND2	0.43	2.28	9	1
1:A:218:MET:CE	1:A:218:MET:HA	0.43	2.43	14	1
1:A:218:MET:O	1:A:222:VAL:HG12	0.43	2.13	20	1
1:A:187:PHE:CZ	1:A:224:THR:HG23	0.43	2.48	7	3
1:A:191:VAL:HG12	1:A:196:ILE:HD12	0.43	1.90	19	1
1:A:218:MET:O	1:A:218:MET:HE2	0.43	2.14	16	1
1:A:218:MET:HA	1:A:218:MET:CE	0.42	2.44	15	1
1:A:173:TYR:CG	1:A:177:VAL:HG11	0.42	2.49	14	1
1:A:183:VAL:HG13	1:A:231:CYS:HB2	0.42	1.90	4	1
1:A:141:MET:SD	1:A:230:MET:HG2	0.41	2.55	9	1
1:A:146:TYR:CD1	1:A:148:PHE:CZ	0.41	3.09	6	1
1:A:183:VAL:HG21	1:A:232:VAL:HG22	0.41	1.92	11	1
1:A:148:PHE:CZ	1:A:156:TRP:CE3	0.41	3.09	10	1
1:A:223:VAL:HA	1:A:226:VAL:HG12	0.41	1.93	13	1
1:A:223:VAL:O	1:A:227:ILE:HD12	0.40	2.16	17	1

## 6.3 Torsion angles

### 6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/117 (80%)	81±2 (86±2%)	9±2 (9±2%)	4±1 (5±1%)	5	29
All	All	1880/2340 (80%)	1623 (86%)	172 (9%)	85 (5%)	5	29

All 16 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	175	SER	20
1	A	166	ASN	13
1	A	212	ASN	9
1	A	148	PHE	7
1	A	146	TYR	6
1	A	164	TYR	5
1	A	163	ARG	4
1	A	147	HIS	4
1	A	144	MET	4
1	A	145	ASN	4
1	A	172	ASP	2
1	A	149	ASP	2
1	A	135	TYR	2
1	A	177	VAL	1
1	A	176	PRO	1
1	A	165	PRO	1

### 6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/99 (88%)	67±3 (77±4%)	20±3 (23±4%)	3	28

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1740/1980 (88%)	1337 (77%)	403 (23%)	<b>3</b> 28

All 60 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	220	ASN	20
1	A	177	VAL	20
1	A	155	ARG	19
1	A	179	GLN	17
1	A	214	THR	16
1	A	238	TYR	15
1	A	157	TRP	14
1	A	226	VAL	14
1	A	218	MET	14
1	A	167	ARG	13
1	A	156	TRP	12
1	A	221	LYS	11
1	A	160	ASN	11
1	A	212	ASN	10
1	A	187	PHE	10
1	A	228	ARG	10
1	A	139	ARG	10
1	A	219	GLU	9
1	A	240	LEU	9
1	A	223	VAL	9
1	A	170	TYR	9
1	A	180	ASP	8
1	A	154	TYR	8
1	A	213	GLN	7
1	A	233	GLN	6
1	A	234	GLN	6
1	A	142	SER	6
1	A	192	THR	6
1	A	163	ARG	6
1	A	158	SER	6
1	A	230	MET	5
1	A	237	GLU	4
1	A	215	GLU	4
1	A	153	GLU	4
1	A	152	ASP	4
1	A	164	TYR	4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Models (Total)
1	A	145	ASN	4
1	A	147	HIS	3
1	A	188	ASN	3
1	A	183	VAL	3
1	A	150	ARG	3
1	A	231	CYS	3
1	A	193	GLU	3
1	A	225	LYS	3
1	A	159	GLU	2
1	A	172	ASP	2
1	A	239	ARG	2
1	A	166	ASN	2
1	A	191	VAL	2
1	A	149	ASP	2
1	A	189	ILE	1
1	A	140	VAL	1
1	A	174	SER	1
1	A	229	GLU	1
1	A	196	ILE	1
1	A	185	ASP	1
1	A	235	TYR	1
1	A	175	SER	1
1	A	186	CYS	1
1	A	144	MET	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 80% for the well-defined parts and 81% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 6269

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1294
Number of shifts mapped to atoms	1294
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	113	$0.29 \pm 0.16$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	101	$0.68 \pm 0.11$	Should be applied
$^{13}\text{C}'$	0	—	None (insufficient data)
$^{15}\text{N}$	110	$-1.31 \pm 0.30$	Should be applied

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 80%, i.e. 984 atoms were assigned a chemical shift out of a possible 1227. 12 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	358/460 (78%)	180/183 (98%)	90/188 (48%)	88/89 (99%)
Sidechain	526/629 (84%)	336/373 (90%)	168/217 (77%)	22/39 (56%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	100/138 (72%)	64/71 (90%)	34/64 (53%)	2/3 (67%)
Overall	984/1227 (80%)	580/627 (93%)	292/469 (62%)	112/131 (85%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 81%, i.e. 1159 atoms were assigned a chemical shift out of a possible 1434. 15 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	448/575 (78%)	225/229 (98%)	113/234 (48%)	110/112 (98%)
Sidechain	611/721 (85%)	387/425 (91%)	201/254 (79%)	23/42 (55%)
Aromatic	100/138 (72%)	64/71 (90%)	34/64 (53%)	2/3 (67%)
Overall	1159/1434 (81%)	676/725 (93%)	348/552 (63%)	135/157 (86%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	157	TRP	NE1	139.20	139.19 – 119.59	5.0

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

