



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2018 – 08:41 am GMT

PDB ID : 3V0B  
Title : 3.9 angstrom crystal structure of BoNT/Ai in complex with NTNHA  
Authors : Gu, S.; Rumpel, S.; Zhou, J.; Strotmeier, J.; Bigalke, H.; Perry, K.; Shoemaker, C.B.; Rummel, A.; Jin, R.  
Deposited on : 2011-12-07  
Resolution : 3.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1145 (4.20-3.60)
Clashscore	122126	1225 (4.20-3.60)
Ramachandran outliers	120053	1184 (4.20-3.60)
Sidechain outliers	120020	1175 (4.20-3.60)
RSRZ outliers	108989	1046 (4.20-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1296	<div> <div>6%</div> <div>79%</div> <div>19%</div> <div>..</div> </div>
2	B	1196	<div> <div>6%</div> <div>71%</div> <div>22%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ZN	A	1297	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 19830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BoNT/A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1280	Total	C	N	O	S	36	0	0
			10418	6681	1721	1983	33			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	224	GLN	GLU	ENGINEERED MUTATION	UNP Q7B8V4
A	363	ALA	ARG	ENGINEERED MUTATION	UNP Q7B8V4
A	366	PHE	TYR	ENGINEERED MUTATION	UNP Q7B8V4
A	1158	ALA	THR	CONFLICT	UNP Q7B8V4

- Molecule 2 is a protein called NTN1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1150	Total	C	N	O	S	46	0	0
			9410	6022	1515	1841	32			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	EXPRESSION TAG	UNP Q45914
B	0	SER	-	EXPRESSION TAG	UNP Q45914
B	1194	PRO	-	EXPRESSION TAG	UNP Q45914

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: BoNT/A



#### • Molecule 2: NTNH



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	282.23Å 282.23Å 374.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.47 – 3.90 48.88 – 3.90	Depositor EDS
% Data completeness (in resolution range)	92.1 (48.47-3.90) 92.1 (48.88-3.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.63 (at 3.88Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, $R_{free}$	0.252 , 0.280 0.250 , 0.276	Depositor DCC
$R_{free}$ test set	3737 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	139.8	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 121.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	19830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	179.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/10639	0.49	0/14405
2	B	0.32	0/9605	0.51	0/13015
All	All	0.31	0/20244	0.50	0/27420

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10418	0	10297	190	30
2	B	9410	0	9202	214	31
3	A	1	0	0	0	0
4	A	1	0	0	0	0
All	All	19830	0	19499	382	32

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 382 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:528:ASP:HB2	2:B:530:SER:H	1.07	1.09
2:B:426:THR:HB	2:B:427:VAL:HA	1.02	1.02
2:B:528:ASP:HB2	2:B:530:SER:N	1.76	1.00
2:B:426:THR:HB	2:B:427:VAL:CA	1.91	0.99
2:B:426:THR:CB	2:B:427:VAL:HA	1.93	0.98

The worst 5 of 32 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:832:GLY:CA	2:B:925:GLU:OE1[4_655]	1.34	0.86
1:A:832:GLY:C	2:B:925:GLU:OE1[4_655]	1.35	0.85
1:A:1187:LYS:N	2:B:793:ASP:OD1[10_555]	1.42	0.78
1:A:832:GLY:N	2:B:925:GLU:OE2[4_655]	1.53	0.67
1:A:825:ASP:N	2:B:718:ASP:CB[4_655]	1.53	0.67

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1276/1296 (98%)	1242 (97%)	34 (3%)	0	100	100
2	B	1144/1196 (96%)	1105 (97%)	38 (3%)	1 (0%)	53	86
All	All	2420/2492 (97%)	2347 (97%)	72 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	429	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1162/1176 (99%)	1126 (97%)	36 (3%)	43	71
2	B	1076/1114 (97%)	1007 (94%)	69 (6%)	19	53
All	All	2238/2290 (98%)	2133 (95%)	105 (5%)	29	62

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	320	LEU
2	B	509	THR
2	B	1115	SER
2	B	326	ILE
2	B	469	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	913	GLN
1	A	1265	ASN
2	B	961	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1280/1296 (98%)	0.34	77 (6%)	22 16	106, 184, 268, 448	10 (0%)
2	B	1150/1196 (96%)	0.35	75 (6%)	19 13	92, 159, 251, 587	14 (1%)
All	All	2430/2492 (97%)	0.34	152 (6%)	20 14	92, 173, 263, 587	24 (0%)

The worst 5 of 152 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	MET	12.1
2	B	451	ASP	8.7
2	B	150	TYR	7.6
1	A	1227	ASN	7.3
2	B	43	PRO	6.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	A	1297	1/1	0.75	0.43	550,550,550,550	0
4	CA	A	1298	1/1	0.84	0.12	169,169,169,169	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.