



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2019 – 10:38 AM EDT

PDB ID : 1V1P
Title : The structure SSL from Staphylococcus Aureus from an orthorhombic crystal form
Authors : Briggs, D.C.; Naylor, C.E.
Deposited on : 2004-04-21
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.4
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	211	<div> <div>7%</div> <div>65%</div> <div>22%</div> <div>•</div> <div>9%</div> </div>
2	B	213	<div> <div>6%</div> <div>68%</div> <div>15%</div> <div>6%</div> <div>10%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exotoxin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	0	0
			1529	958	265	305	1			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q9ZFS5
A	2	ARG	-	expression tag	UNP Q9ZFS5
A	3	GLY	-	expression tag	UNP Q9ZFS5
A	4	SER	-	expression tag	UNP Q9ZFS5
A	5	HIS	-	expression tag	UNP Q9ZFS5
A	6	HIS	-	expression tag	UNP Q9ZFS5
A	7	HIS	-	expression tag	UNP Q9ZFS5
A	8	HIS	-	expression tag	UNP Q9ZFS5
A	9	HIS	-	expression tag	UNP Q9ZFS5
A	10	HIS	-	expression tag	UNP Q9ZFS5
A	11	GLY	-	expression tag	UNP Q9ZFS5
A	12	SER	ALA	conflict	UNP Q9ZFS5
A	?	-	LYS	deletion	UNP Q9ZFS5
A	?	-	ASP	deletion	UNP Q9ZFS5

- Molecule 2 is a protein called Exotoxin 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	191	Total	C	N	O	S	0	0	0
			1514	948	261	304	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP Q9ZFS5
B	2	ARG	-	expression tag	UNP Q9ZFS5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	GLY	-	expression tag	UNP Q9ZFS5
B	4	SER	-	expression tag	UNP Q9ZFS5
B	5	HIS	-	expression tag	UNP Q9ZFS5
B	6	HIS	-	expression tag	UNP Q9ZFS5
B	7	HIS	-	expression tag	UNP Q9ZFS5
B	8	HIS	-	expression tag	UNP Q9ZFS5
B	9	HIS	-	expression tag	UNP Q9ZFS5
B	10	HIS	-	expression tag	UNP Q9ZFS5
B	11	GLY	-	expression tag	UNP Q9ZFS5
B	12	SER	ALA	conflict	UNP Q9ZFS5

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	16	Total O 16 16	0	0
3	B	8	Total O 8 8	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Exotoxin 1

Chain A: 



• Molecule 2: Exotoxin 1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.65Å 71.60Å 103.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.72 – 2.70 31.07 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (58.72-2.70) 100.0 (31.07-2.70)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.68Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.232 , 0.301 0.240 , 0.244	Depositor DCC
R_{free} test set	527 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	43.5	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3067	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	1/1550 (0.1%)	0.79	5/2085 (0.2%)
2	B	0.50	1/1535 (0.1%)	0.84	9/2068 (0.4%)
All	All	0.50	2/3085 (0.1%)	0.82	14/4153 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	2
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	211	ILE	C-OXT	5.20	1.33	1.23
2	B	213	ILE	C-OXT	5.16	1.33	1.23

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	182	ASP	CB-CG-OD2	8.08	125.57	118.30
1	A	70	ASP	CB-CG-OD2	7.51	125.06	118.30
2	B	72	ASP	CB-CG-OD2	6.91	124.52	118.30
2	B	116	THR	O-C-N	-6.72	108.33	121.10
2	B	187	ASP	CB-CG-OD2	6.55	124.19	118.30
1	A	24	ASP	CB-CG-OD2	6.48	124.13	118.30
2	B	27	ASP	CB-CG-OD2	6.13	123.81	118.30
2	B	127	ASP	CB-CG-OD2	5.79	123.51	118.30
1	A	200	ASP	CB-CG-OD2	5.75	123.47	118.30
2	B	177	ASP	CB-CG-OD2	5.54	123.29	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	129	ASP	CB-CG-OD2	5.36	123.12	118.30
2	B	184	ASP	CB-CG-OD2	5.22	123.00	118.30
1	A	27	ASP	CB-CG-OD2	5.20	122.98	118.30
2	B	202	ASP	CB-CG-OD2	5.10	122.89	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	104	LYS	Peptide
2	B	116	THR	Mainchain
2	B	48	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1529	0	1497	18	0
2	B	1514	0	1469	20	0
3	A	16	0	0	1	0
3	B	8	0	0	0	0
All	All	3067	0	2966	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:ASN:C	1:A:172:ASN:HD22	1.71	0.93
2:B:84:ASN:ND2	2:B:109:ASN:HD22	1.90	0.69
2:B:106:LYS:O	2:B:107:LYS:HB2	1.94	0.68
1:A:172:ASN:C	1:A:172:ASN:ND2	2.47	0.67
2:B:106:LYS:O	2:B:107:LYS:CB	2.49	0.61
1:A:105:LYS:HA	1:A:107:ASN:N	2.16	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:VAL:HG21	2:B:123:VAL:HG21	1.85	0.58
2:B:98:LEU:C	2:B:98:LEU:HD12	2.26	0.55
1:A:172:ASN:HD21	1:A:175:ASP:H	1.54	0.54
1:A:105:LYS:HA	1:A:106:ASN:C	2.28	0.54
2:B:84:ASN:HD22	2:B:109:ASN:HD22	1.57	0.51
1:A:126:LEU:HD11	2:B:209:THR:HB	1.93	0.50
2:B:80:LEU:CD2	2:B:83:GLN:HG2	2.42	0.49
2:B:58:ASN:HB2	2:B:63:ASN:HA	1.95	0.49
1:A:71:LYS:NZ	3:A:2004:HOH:O	2.46	0.47
1:A:121:VAL:HG11	2:B:123:VAL:HG11	1.98	0.46
1:A:45:LYS:O	1:A:46:VAL:HB	2.15	0.45
1:A:142:LEU:HD11	1:A:181:ILE:HD11	1.98	0.45
2:B:175:LEU:O	2:B:176:LYS:C	2.55	0.45
2:B:146:GLU:O	2:B:150:LYS:HG2	2.16	0.45
1:A:171:ILE:HB	1:A:179:VAL:HB	1.99	0.45
1:A:57:PHE:O	1:A:62:HIS:HB2	2.18	0.44
2:B:177:ASP:HA	2:B:178:GLU:HG2	1.99	0.44
2:B:176:LYS:O	2:B:177:ASP:C	2.56	0.44
2:B:175:LEU:O	2:B:178:GLU:HA	2.18	0.43
1:A:163:THR:OG1	1:A:210:GLN:NE2	2.52	0.43
1:A:54:VAL:HG12	1:A:55:VAL:N	2.34	0.42
1:A:145:LEU:HD23	1:A:171:ILE:HD13	2.01	0.42
2:B:89:GLN:HG3	2:B:98:LEU:HD13	2.02	0.42
2:B:171:ILE:HG12	2:B:208:VAL:HG22	2.02	0.41
2:B:174:ASN:C	2:B:174:ASN:HD22	2.23	0.41
1:A:86:VAL:HG23	1:A:143:LYS:HE3	2.03	0.41
2:B:58:ASN:ND2	2:B:59:PRO:O	2.54	0.41
2:B:83:GLN:HB2	2:B:105:THR:O	2.21	0.40
1:A:62:HIS:CD2	1:A:96:LEU:HG	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	187/211 (89%)	166 (89%)	16 (9%)	5 (3%)	5	14
2	B	189/213 (89%)	169 (89%)	14 (7%)	6 (3%)	4	10
All	All	376/424 (89%)	335 (89%)	30 (8%)	11 (3%)	5	12

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	46	VAL
1	A	105	LYS
1	A	174	LYS
2	B	50	ASN
2	B	107	LYS
1	A	50	ASN
2	B	48	ASN
2	B	81	GLN
2	B	177	ASP
2	B	49	TYR
1	A	24	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/192 (90%)	146 (85%)	26 (15%)	3	8
2	B	168/194 (87%)	146 (87%)	22 (13%)	4	11
All	All	340/386 (88%)	292 (86%)	48 (14%)	4	9

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	HIS
1	A	38	GLU
1	A	40	SER
1	A	41	ASN
1	A	47	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	56	ARG
1	A	58	ASN
1	A	73	GLN
1	A	75	LYS
1	A	79	GLN
1	A	96	LEU
1	A	104	LYS
1	A	105	LYS
1	A	110	SER
1	A	122	ASN
1	A	126	LEU
1	A	129	SER
1	A	130	ILE
1	A	136	GLN
1	A	137	LYS
1	A	138	GLU
1	A	172	ASN
1	A	185	ASP
1	A	188	GLN
1	A	191	ARG
1	A	209	ASN
2	B	26	ARG
2	B	36	SER
2	B	38	GLU
2	B	49	TYR
2	B	50	ASN
2	B	58	ASN
2	B	77	LYS
2	B	83	GLN
2	B	93	ASP
2	B	98	LEU
2	B	105	THR
2	B	124	ASN
2	B	128	LEU
2	B	134	SER
2	B	136	LEU
2	B	155	LEU
2	B	174	ASN
2	B	175	LEU
2	B	176	LYS
2	B	177	ASP
2	B	178	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	211	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	60	GLN
1	A	61	ASN
1	A	62	HIS
1	A	63	GLN
1	A	122	ASN
1	A	136	GLN
1	A	155	ASN
1	A	172	ASN
1	A	209	ASN
1	A	210	GLN
2	B	53	ASN
2	B	58	ASN
2	B	84	ASN
2	B	124	ASN
2	B	158	ASN
2	B	174	ASN
2	B	211	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	59:PRO	C	60:GLN	N	4.13

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	191/211 (90%)	0.37	15 (7%)	12 10	26, 40, 62, 67	0
2	B	191/213 (89%)	0.33	12 (6%)	20 19	26, 40, 62, 67	0
All	All	382/424 (90%)	0.35	27 (7%)	16 14	26, 40, 62, 67	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	213	ILE	5.7
2	B	108	ASN	4.8
2	B	51	GLY	4.5
2	B	50	ASN	4.2
1	A	22	LEU	4.1
1	A	93	ASN	4.0
2	B	43	SER	4.0
1	A	58	ASN	3.9
1	A	60	GLN	3.7
1	A	39	TYR	3.1
2	B	60	LYS	3.0
1	A	61	ASN	2.9
1	A	52	SER	2.9
1	A	62	HIS	2.6
1	A	21	HIS	2.5
1	A	57	PHE	2.4
2	B	41	ASN	2.4
1	A	211	ILE	2.3
2	B	58	ASN	2.3
1	A	95	ARG	2.2
2	B	52	SER	2.2
1	A	94	GLY	2.2
2	B	49	TYR	2.2
2	B	62	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	106	ASN	2.1
2	B	95	ASN	2.1
1	A	56	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.