



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 07:27 am GMT

PDB ID : 2V4Q
Title : Post-insertion complex of the Y-family DNA polymerase Dpo4 with M1dG containing template DNA
Authors : Eoff, R.L.; Stafford, J.B.; Szekely, J.; Rizzo, C.J.; Egli, M.; Guengerich, F.P.; Marnett, L.J.
Deposited on : 2008-09-26
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix)	:	1.13
EDS	:	trunk31020
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 6%, orange 0%, yellow 65%, green 29%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 6% 65% 29% • • </div> </div>
2	P	14	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 57%, yellow 43%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 57% 43% </div> </div>
3	T	18	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 6%, orange 0%, yellow 33%, green 56%, orange 6%, grey 6%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 6% 33% 56% 6% 6% </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	1	1
			2755	1766	477	505	7			

- Molecule 2 is a DNA chain called 5'-D(*GP*GP*GP*GP*GP*AP*AP*GP*GP*AP*TP*TP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	14	Total	C	N	O	P	0	0	0
			292	138	60	81	13			

- Molecule 3 is a DNA chain called 5'-D(*TP*CP*AP*C M1GP*GP*AP*AP*TP*CP*CP *TP*TP*CP*CP*CP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	17	Total	C	N	O	P	0	0	0
			338	164	58	100	16			

- Molecule 4 is 2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (three-letter code: DGT) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Ca 3 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	60	Total O 60 60	0	0
6	P	16	Total O 16 16	0	0
6	T	10	Total O 10 10	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.61Å 103.14Å 52.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.72 – 2.60 28.72 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (28.72-2.60) 99.6 (28.72-2.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 2.61Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.205 , 0.254 0.204 , 0.250	Depositor DCC
R_{free} test set	771 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	58.1	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3505	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, M1G, DGT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2794	0.62	1/3752 (0.0%)
2	P	0.63	0/329	0.67	0/508
3	T	1.50	6/347 (1.7%)	0.85	0/528
All	All	0.64	6/3470 (0.2%)	0.65	1/4788 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	T	0	2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	2	DC	C4-N4	15.11	1.47	1.33
3	T	2	DC	C2-N3	9.29	1.43	1.35
3	T	2	DC	C5-C6	9.02	1.41	1.34
3	T	2	DC	O5'-C5'	7.95	1.62	1.42
3	T	2	DC	C4-C5	7.25	1.48	1.43
3	T	2	DC	N3-C4	6.41	1.38	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	ARG	NE-CZ-NH1	5.22	122.91	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	T	10	DC	Sidechain
3	T	7	DA	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2755	0	2901	94	0
2	P	292	0	158	7	0
3	T	338	0	192	15	0
4	A	31	0	12	0	0
5	A	3	0	0	0	0
6	A	60	0	0	8	0
6	P	16	0	0	3	0
6	T	10	0	0	1	0
All	All	3505	0	3263	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ASP:OD1	1:A:169:GLU:HG2	1.69	0.91
3:T:3:DA:H1'	6:T:2002:HOH:O	1.71	0.89
1:A:193:LYS:HD3	1:A:216:MET:HE2	1.55	0.89
1:A:17:GLU:HB3	1:A:24:LYS:NZ	1.92	0.84
1:A:256:ARG:HH11	1:A:256:ARG:HB3	1.44	0.80
1:A:257:ASN:HB3	1:A:260:GLU:HG2	1.64	0.80
1:A:256:ARG:NH1	1:A:256:ARG:HB3	1.96	0.79
1:A:39:ASP:OD1	1:A:65:LYS:HE3	1.84	0.78
1:A:247:ARG:HB2	6:A:2045:HOH:O	1.87	0.75
3:T:3:DA:H2'	3:T:3:DA:N3	2.02	0.74
1:A:254:ASN:HD22	1:A:291:GLU:HB3	1.54	0.73
1:A:254:ASN:ND2	1:A:291:GLU:HB3	2.03	0.72
1:A:38:GLU:O	1:A:39:ASP:HB2	1.89	0.71
2:P:3:DG:H2''	2:P:4:DG:H5'	1.72	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:GLU:O	1:A:137:LYS:HE2	1.92	0.70
1:A:210:PHE:CE2	1:A:214:LYS:HD2	2.27	0.69
2:P:13:DC:H4'	2:P:14:DC:H5''	1.75	0.69
1:A:113:ASP:O	1:A:114:LYS:HG2	1.93	0.68
1:A:17:GLU:HB3	1:A:24:LYS:HZ1	1.57	0.67
1:A:119:ARG:NH1	6:A:2024:HOH:O	2.27	0.66
1:A:157:MET:CE	1:A:166:ILE:HD11	2.27	0.65
1:A:331:ARG:HD2	6:A:2056:HOH:O	1.97	0.65
1:A:157:MET:HE2	1:A:166:ILE:HD11	1.78	0.65
1:A:17:GLU:HB3	1:A:24:LYS:HZ2	1.60	0.64
1:A:257:ASN:HB3	1:A:260:GLU:CG	2.28	0.64
3:T:2:DC:C2'	3:T:3:DA:H5'	2.28	0.64
1:A:270:GLU:OE2	1:A:308:LYS:HD3	2.00	0.61
1:A:167:ASP:O	1:A:171:VAL:HG23	2.00	0.61
1:A:318:LEU:O	1:A:322:ILE:HG13	2.00	0.61
3:T:14:DC:H2''	3:T:15:DC:H5'	1.83	0.61
1:A:298:ARG:NH1	2:P:8:DG:H3'	2.17	0.59
1:A:331:ARG:NH2	3:T:4:DC:OP1	2.33	0.58
1:A:233:TYR:CE1	1:A:235:GLU:HG2	2.38	0.58
1:A:36:ARG:NH1	1:A:331:ARG:NH1	2.52	0.57
1:A:293:LEU:HD22	1:A:293:LEU:N	2.19	0.56
1:A:247:ARG:HG3	1:A:247:ARG:HH11	1.71	0.56
1:A:119:ARG:HD3	6:A:2024:HOH:O	2.04	0.56
1:A:256:ARG:CB	1:A:256:ARG:HH11	2.15	0.56
1:A:99:ILE:HD12	1:A:109:LEU:HD21	1.89	0.55
1:A:40:SER:HB2	3:T:3:DA:OP2	2.07	0.55
1:A:31:CYS:HB3	1:A:61:ILE:HD11	1.89	0.54
1:A:169:GLU:CG	1:A:170:GLU:N	2.71	0.54
1:A:171:VAL:O	1:A:175:ILE:HG13	2.07	0.54
1:A:169:GLU:HG3	1:A:170:GLU:N	2.23	0.53
1:A:256:ARG:HG3	1:A:329:LYS:HG3	1.91	0.53
1:A:84:VAL:O	1:A:88:ILE:HG13	2.09	0.52
1:A:170:GLU:HA	1:A:173:ARG:HG3	1.91	0.52
1:A:76:MET:CE	1:A:78:LYS:HG2	2.40	0.52
1:A:6:VAL:O	1:A:106:GLU:HA	2.11	0.51
3:T:14:DC:H1'	3:T:15:DC:H5''	1.92	0.51
3:T:2:DC:H2'	3:T:3:DA:H5'	1.91	0.51
1:A:79:GLU:O	1:A:82:GLN:HB3	2.11	0.51
3:T:3:DA:C2'	3:T:3:DA:N3	2.71	0.51
1:A:145:SER:OG	1:A:146:LYS:N	2.43	0.50
1:A:242:ARG:HA	1:A:242:ARG:HH11	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:ILE:HG12	1:A:104:ILE:O	2.12	0.49
1:A:262:LYS:HB2	1:A:263:PRO:HD3	1.95	0.49
3:T:14:DC:H2''	3:T:15:DC:C5'	2.42	0.49
1:A:248:ILE:HA	1:A:334:GLY:HA3	1.94	0.49
2:P:4:DG:H2''	2:P:5:DG:OP2	2.11	0.49
1:A:199:ILE:HG23	1:A:204:ASP:HB2	1.94	0.49
1:A:8:PHE:N	1:A:8:PHE:CD1	2.77	0.49
3:T:11:DC:H2'	3:T:12:DT:H71	1.95	0.48
1:A:116:ARG:HG3	6:A:2025:HOH:O	2.13	0.47
1:A:153:ILE:HG22	1:A:157:MET:HE2	1.96	0.47
1:A:244:SER:O	1:A:245:ILE:HG13	2.14	0.47
1:A:93:ARG:HG3	1:A:93:ARG:HH11	1.81	0.46
1:A:189:ILE:O	1:A:193:LYS:HG3	2.15	0.46
1:A:166:ILE:HD13	1:A:174:LEU:HD11	1.98	0.46
1:A:289:VAL:HB	1:A:332:ARG:HB2	1.97	0.46
1:A:147:ASN:HD21	1:A:150:PHE:HD1	1.63	0.45
1:A:32:VAL:HG11	3:T:5:M1G:H5'	1.99	0.45
1:A:275:LYS:NZ	6:A:2045:HOH:O	2.48	0.45
2:P:14:DC:OP1	6:P:2015:HOH:O	2.21	0.45
1:A:179:ASP:HB3	1:A:182:ASP:OD2	2.16	0.45
3:T:2:DC:H2''	3:T:3:DA:OP1	2.16	0.45
1:A:195:LYS:HA	1:A:195:LYS:HD3	1.84	0.45
1:A:98:LYS:HB2	1:A:110:ASP:HB3	1.98	0.44
1:A:115:VAL:HG13	1:A:120:GLU:HB3	1.99	0.44
1:A:257:ASN:CB	1:A:260:GLU:HG2	2.43	0.44
1:A:276:LEU:O	1:A:279:ARG:HB2	2.18	0.44
1:A:193:LYS:HD3	1:A:216:MET:CE	2.38	0.43
1:A:62:VAL:HG23	6:A:2008:HOH:O	2.18	0.43
1:A:60:PRO:HB3	3:T:3:DA:H5''	1.99	0.43
1:A:233:TYR:CZ	1:A:235:GLU:HG2	2.53	0.43
1:A:244:SER:HB2	3:T:8:DA:OP2	2.19	0.43
1:A:255:SER:N	1:A:329:LYS:HG2	2.33	0.43
1:A:161:ASN:O	1:A:161:ASN:OD1	2.37	0.43
1:A:189:ILE:HG22	1:A:193:LYS:NZ	2.34	0.42
1:A:27:PRO:HG2	1:A:71:ALA:HA	2.01	0.42
1:A:136:GLU:O	1:A:137:LYS:HB2	2.19	0.42
1:A:15:VAL:HG11	1:A:81:TYR:HD1	1.84	0.42
1:A:169:GLU:CG	1:A:170:GLU:H	2.32	0.42
1:A:133:LEU:O	1:A:137:LYS:HA	2.19	0.42
1:A:115:VAL:HG12	1:A:116:ARG:N	2.34	0.42
1:A:329:LYS:HB2	1:A:329:LYS:HE3	1.87	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:14:DC:P	6:P:2016:HOH:O	2.76	0.42
1:A:124:LEU:O	1:A:128:ILE:HG13	2.18	0.42
1:A:247:ARG:HG3	1:A:247:ARG:NH1	2.35	0.42
1:A:76:MET:HE3	1:A:78:LYS:HG2	2.01	0.42
2:P:14:DC:OP2	6:P:2014:HOH:O	2.21	0.42
1:A:127:GLU:C	1:A:127:GLU:OE2	2.58	0.41
1:A:256:ARG:CG	1:A:329:LYS:HA	2.49	0.41
1:A:9:ASP:O	1:A:10:TYR:C	2.58	0.41
1:A:331:ARG:NH1	6:A:2056:HOH:O	2.53	0.41
1:A:20:ASN:ND2	1:A:22:SER:OG	2.53	0.41
1:A:1:MET:HA	1:A:112:SER:OG	2.20	0.41
1:A:293:LEU:CD2	1:A:293:LEU:N	2.84	0.40
1:A:251:MET:HG2	1:A:264:TYR:CD2	2.56	0.40
1:A:78:LYS:HE3	1:A:104:ILE:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	341/358 (95%)	314 (92%)	23 (7%)	4 (1%)	14 30

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	TYR
1	A	38	GLU
1	A	39	ASP
1	A	277	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	301/315 (96%)	293 (97%)	8 (3%)	48 74

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ASP
1	A	8	PHE
1	A	20	ASN
1	A	130	ASN
1	A	244	SER
1	A	247	ARG
1	A	273	TYR
1	A	300	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	123	ASN
1	A	161	ASN
1	A	234	ASN
1	A	285	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	M1G	T	5	3	20,28,29	1.01	1 (5%)	19,41,44	1.51	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	M1G	T	5	3	-	0/3/21/22	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	5	M1G	C8-N9	-2.73	1.33	1.36

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	5	M1G	O4'-C1'-C2'	-2.39	101.70	106.25
3	T	5	M1G	C4-C5-N7	-2.24	107.24	109.41
3	T	5	M1G	C4'-O4'-C1'	-2.04	104.49	109.43
3	T	5	M1G	C6A-N2-C2	3.66	121.24	116.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	T	5	M1G	1	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	DGT	A	594	5	27,33,33	1.36	3 (11%)	31,52,52	1.94	9 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DGT	A	594	5	-	0/18/34/34	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	594	DGT	C8-N9	-2.76	1.33	1.36
4	A	594	DGT	C2-N1	3.04	1.40	1.35
4	A	594	DGT	C6-N1	4.23	1.40	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	594	DGT	N3-C2-N1	-4.71	120.50	127.41
4	A	594	DGT	PB-O3B-PG	-4.28	118.23	132.63
4	A	594	DGT	C5-C6-N1	-3.22	118.89	123.47
4	A	594	DGT	PA-O3A-PB	-2.84	123.08	132.63
4	A	594	DGT	C2'-C3'-C4'	-2.43	97.61	102.76
4	A	594	DGT	C4-C5-N7	-2.06	107.42	109.41
4	A	594	DGT	N2-C2-N1	2.11	120.57	117.25
4	A	594	DGT	C6-N1-C2	2.15	119.16	116.06
4	A	594	DGT	C2-N3-C4	4.95	120.94	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/358 (95%)	-0.15	4 (1%) 79 75	32, 56, 76, 84	35 (10%)
2	P	14/14 (100%)	-0.20	0 100 100	48, 59, 78, 81	1 (7%)
3	T	16/18 (88%)	0.07	1 (6%) 20 15	46, 67, 89, 90	0
All	All	372/390 (95%)	-0.14	5 (1%) 77 73	32, 57, 77, 90	36 (9%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	T	2	DC	3.1
1	A	192	GLU	2.9
1	A	119	ARG	2.7
1	A	216	MET	2.3
1	A	2	ILE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	M1G	T	5	25/26	0.97	0.15	39,43,50,51	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	CA	A	1001	1/1	0.94	0.25	57,57,57,57	0
5	CA	A	1002	1/1	0.95	0.14	77,77,77,77	0
5	CA	A	1000	1/1	0.95	0.19	43,43,43,43	0
4	DGT	A	594	31/31	0.97	0.16	38,43,52,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.