



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 09:53 pm GMT

PDB ID : 1V5I
Title : Crystal structure of serine protease inhibitor POIA1 in complex with subtilisin BPN'
Authors : Lee, W.C.; Kikkawa, M.; Kojima, S.; Miura, K.; Tanokura, M.
Deposited on : 2003-11-24
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	2727 (1.50-1.50)
Ramachandran outliers	120053	2661 (1.50-1.50)
Sidechain outliers	120020	2659 (1.50-1.50)
RSRZ outliers	108989	2481 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
2	B	76	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	2001	-	X	-	-
5	GOL	A	2003	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	2005	-	X	-	-
5	GOL	A	2006	-	X	-	-
5	GOL	A	2007	-	X	-	-
5	GOL	A	2008	-	X	-	-
5	GOL	A	2009	-	X	-	-
5	GOL	A	2010	-	X	-	-
5	GOL	A	2011	-	X	-	-
5	GOL	A	2012	-	X	-	-
5	GOL	A	2013	-	X	-	-
5	GOL	A	2016	-	X	-	-
5	GOL	A	2018	-	X	-	X
5	GOL	B	2002	-	X	-	-
5	GOL	B	2004	-	X	-	-
5	GOL	B	2014	-	X	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3039 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Subtilisin BPN'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			1940	1204	335	395	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	221	CSD	SER	ENGINEERED MUTATION	UNP P00782

- Molecule 2 is a protein called IA-1=serine proteinase inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	76	Total	C	N	O	S	0	0	0
			587	368	95	122	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	71	HIS	GLY	ENGINEERED MUTATION	UNP Q7M4T6
B	72	VAL	ILE	ENGINEERED MUTATION	UNP Q7M4T6
B	73	ALA	VAL	ENGINEERED MUTATION	UNP Q7M4T6
B	74	HIS	THR	ENGINEERED MUTATION	UNP Q7M4T6
B	75	ALA	THR	ENGINEERED MUTATION	UNP Q7M4T6
B	76	TYR	GLN	ENGINEERED MUTATION	UNP Q7M4T6

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	318	Total O 318 318	0	0
6	B	87	Total O 87 87	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Subtilisin BPN'



- Molecule 2: IA-1=serine proteinase inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.70Å 71.80Å 94.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.90 – 1.50 35.93 – 1.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (35.90-1.50) 93.2 (35.93-1.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 1.30Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.180 , 0.192 0.183 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3039	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, SO4, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/1970	0.60	0/2689
2	B	0.30	0/596	0.58	0/800
All	All	0.27	0/2566	0.60	0/3489

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1940	0	1898	16	0
2	B	587	0	568	6	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	1	0	0	0	0
5	A	78	0	39	4	0
5	B	18	0	9	1	0
6	A	318	0	0	3	0
6	B	87	0	0	1	0
All	All	3039	0	2514	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:ASN:HD22	1:A:64:HIS:H	1.32	0.75
1:A:240:ASN:HB2	6:A:3301:HOH:O	1.91	0.69
1:A:38:SER:HB2	5:A:2018:GOL:H12	1.90	0.53
1:A:43:LYS:HG3	1:A:75:LEU:HD11	1.92	0.52
1:A:62:ASN:ND2	1:A:64:HIS:HB2	2.26	0.51
2:B:63:LEU:HD22	5:B:2002:GOL:H11	1.93	0.50
1:A:62:ASN:O	1:A:63:SER:CB	2.61	0.49
1:A:96:LEU:HD13	2:B:75:ALA:HB2	1.95	0.48
1:A:29:ALA:HB2	1:A:119:MET:HG3	1.98	0.45
1:A:127:GLY:HA2	1:A:167:TYR:O	2.18	0.44
2:B:47:GLU:HG3	6:B:2054:HOH:O	2.17	0.43
1:A:182:SER:OG	5:A:2007:GOL:H11	2.18	0.43
1:A:240:ASN:HB2	6:A:3255:HOH:O	2.19	0.43
1:A:96:LEU:CD1	2:B:75:ALA:HB2	2.49	0.42
1:A:13:ALA:N	1:A:14:PRO:CD	2.81	0.42
1:A:113:TRP:HE1	5:A:2009:GOL:H12	1.85	0.41
1:A:251:GLU:HB3	1:A:265:LYS:HD2	2.01	0.41
1:A:56:ASN:CG	1:A:59:GLN:HG3	2.40	0.41
5:A:2012:GOL:H12	6:A:3141:HOH:O	2.20	0.41
2:B:7:VAL:CG1	2:B:64:ILE:HD12	2.50	0.41
2:B:10:LYS:HE2	2:B:65:ASP:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	272/275 (99%)	262 (96%)	9 (3%)	1 (0%)	36 14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	74/76 (97%)	72 (97%)	2 (3%)	0	100	100
All	All	346/351 (99%)	334 (96%)	11 (3%)	1 (0%)	43	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	63	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/204 (100%)	203 (100%)	1 (0%)	90	79
2	B	63/63 (100%)	62 (98%)	1 (2%)	65	38
All	All	267/267 (100%)	265 (99%)	2 (1%)	85	71

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	222	MET
2	B	37	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	ASN
1	A	118	ASN
1	A	206	GLN
1	A	212	ASN
1	A	240	ASN
2	B	34	ASN
2	B	37	ASN
2	B	71	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	74	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CSD	A	221	1	4,7,8	2.10	1 (25%)	2,8,10	4.23	2 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	221	1	-	0/2/6/8	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	CSD	CA-C	3.99	1.55	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	221	CSD	OD1-SG-CB	-5.25	95.54	105.54
1	A	221	CSD	O-C-CA	-2.85	117.12	124.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 1 is monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	1001	-	4,4,4	0.19	0	6,6,6	1.26	1 (16%)
5	GOL	A	2001	-	5,5,5	4.81	5 (100%)	5,5,5	5.71	3 (60%)
5	GOL	A	2003	-	5,5,5	5.03	5 (100%)	5,5,5	5.68	3 (60%)
5	GOL	A	2005	-	5,5,5	4.85	5 (100%)	5,5,5	5.73	3 (60%)
5	GOL	A	2006	-	5,5,5	4.92	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	A	2007	-	5,5,5	4.86	5 (100%)	5,5,5	5.71	3 (60%)
5	GOL	A	2008	-	5,5,5	4.88	5 (100%)	5,5,5	5.70	3 (60%)
5	GOL	A	2009	-	5,5,5	4.85	5 (100%)	5,5,5	5.71	3 (60%)
5	GOL	A	2010	-	5,5,5	4.88	5 (100%)	5,5,5	5.71	3 (60%)
5	GOL	A	2011	-	5,5,5	4.89	5 (100%)	5,5,5	5.71	3 (60%)
5	GOL	A	2012	-	5,5,5	4.88	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	A	2013	-	5,5,5	4.87	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	A	2016	-	5,5,5	4.93	5 (100%)	5,5,5	5.69	3 (60%)
5	GOL	A	2018	-	5,5,5	4.83	5 (100%)	5,5,5	5.77	3 (60%)
3	SO4	B	1002	-	4,4,4	0.24	0	6,6,6	1.23	1 (16%)
5	GOL	B	2002	-	5,5,5	4.86	5 (100%)	5,5,5	5.74	3 (60%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	2004	-	5,5,5	4.85	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	B	2014	-	5,5,5	4.86	5 (100%)	5,5,5	5.72	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	1001	-	-	0/0/0/0	0/0/0/0
5	GOL	A	2001	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2003	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2005	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2006	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2007	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2008	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2009	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2010	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2011	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2012	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2013	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2016	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2018	-	-	0/4/4/4	0/0/0/0
3	SO4	B	1002	-	-	0/0/0/0	0/0/0/0
5	GOL	B	2002	-	-	0/4/4/4	0/0/0/0
5	GOL	B	2004	-	-	0/4/4/4	0/0/0/0
5	GOL	B	2014	-	-	0/4/4/4	0/0/0/0

All (80) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	2016	GOL	C3-C2	-8.03	1.21	1.52
5	A	2003	GOL	C3-C2	-7.98	1.21	1.52
5	A	2006	GOL	C3-C2	-7.97	1.21	1.52
5	A	2008	GOL	C3-C2	-7.95	1.21	1.52
5	A	2013	GOL	C3-C2	-7.94	1.21	1.52
5	A	2012	GOL	C3-C2	-7.93	1.21	1.52
5	A	2011	GOL	C3-C2	-7.90	1.21	1.52
5	A	2010	GOL	C3-C2	-7.90	1.21	1.52
5	A	2005	GOL	C3-C2	-7.88	1.21	1.52
5	A	2007	GOL	C3-C2	-7.88	1.21	1.52
5	B	2014	GOL	C3-C2	-7.87	1.21	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	2002	GOL	C3-C2	-7.87	1.21	1.52
5	B	2004	GOL	C3-C2	-7.86	1.21	1.52
5	A	2009	GOL	C3-C2	-7.86	1.21	1.52
5	A	2001	GOL	C3-C2	-7.85	1.21	1.52
5	A	2018	GOL	C3-C2	-7.81	1.22	1.52
5	A	2016	GOL	O3-C3	-4.28	1.24	1.42
5	A	2006	GOL	O3-C3	-4.28	1.24	1.42
5	A	2018	GOL	O3-C3	-4.28	1.24	1.42
5	A	2011	GOL	O3-C3	-4.25	1.24	1.42
5	A	2012	GOL	O3-C3	-4.24	1.24	1.42
5	A	2008	GOL	O3-C3	-4.24	1.24	1.42
5	A	2010	GOL	O3-C3	-4.23	1.24	1.42
5	A	2013	GOL	O3-C3	-4.23	1.24	1.42
5	A	2007	GOL	O3-C3	-4.22	1.24	1.42
5	A	2005	GOL	O3-C3	-4.22	1.24	1.42
5	B	2002	GOL	O3-C3	-4.21	1.24	1.42
5	A	2009	GOL	O3-C3	-4.21	1.24	1.42
5	B	2014	GOL	O3-C3	-4.21	1.24	1.42
5	B	2004	GOL	O3-C3	-4.20	1.24	1.42
5	A	2003	GOL	O3-C3	-4.13	1.25	1.42
5	A	2001	GOL	O3-C3	-4.09	1.25	1.42
5	A	2003	GOL	C1-C2	-3.46	1.38	1.52
5	A	2001	GOL	C1-C2	-3.15	1.40	1.52
5	A	2016	GOL	C1-C2	-3.11	1.40	1.52
5	A	2007	GOL	C1-C2	-3.10	1.40	1.52
5	A	2013	GOL	C1-C2	-3.07	1.40	1.52
5	A	2012	GOL	C1-C2	-3.06	1.40	1.52
5	A	2003	GOL	O2-C2	-3.06	1.34	1.43
5	A	2011	GOL	O2-C2	-3.05	1.34	1.43
5	B	2002	GOL	C1-C2	-3.04	1.40	1.52
5	A	2010	GOL	C1-C2	-3.03	1.40	1.52
5	A	2008	GOL	C1-C2	-3.00	1.40	1.52
5	A	2006	GOL	C1-C2	-2.99	1.40	1.52
5	A	2006	GOL	O2-C2	-2.98	1.34	1.43
5	A	2011	GOL	C1-C2	-2.98	1.40	1.52
5	A	2009	GOL	C1-C2	-2.98	1.40	1.52
5	B	2014	GOL	C1-C2	-2.97	1.40	1.52
5	B	2004	GOL	C1-C2	-2.96	1.40	1.52
5	A	2007	GOL	O2-C2	-2.95	1.34	1.43
5	B	2014	GOL	O2-C2	-2.95	1.34	1.43
5	A	2005	GOL	C1-C2	-2.94	1.40	1.52
5	B	2002	GOL	O2-C2	-2.93	1.34	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	2009	GOL	O2-C2	-2.93	1.34	1.43
5	A	2008	GOL	O2-C2	-2.92	1.34	1.43
5	A	2012	GOL	O2-C2	-2.92	1.34	1.43
5	B	2004	GOL	O2-C2	-2.92	1.34	1.43
5	A	2018	GOL	C1-C2	-2.91	1.40	1.52
5	A	2001	GOL	O2-C2	-2.91	1.34	1.43
5	A	2010	GOL	O2-C2	-2.90	1.34	1.43
5	A	2005	GOL	O2-C2	-2.88	1.34	1.43
5	A	2016	GOL	O2-C2	-2.88	1.34	1.43
5	A	2013	GOL	O2-C2	-2.86	1.35	1.43
5	A	2018	GOL	O2-C2	-2.72	1.35	1.43
5	A	2001	GOL	O1-C1	4.37	1.60	1.42
5	A	2013	GOL	O1-C1	4.48	1.61	1.42
5	A	2012	GOL	O1-C1	4.48	1.61	1.42
5	A	2007	GOL	O1-C1	4.49	1.61	1.42
5	A	2008	GOL	O1-C1	4.51	1.61	1.42
5	A	2016	GOL	O1-C1	4.54	1.61	1.42
5	B	2002	GOL	O1-C1	4.55	1.61	1.42
5	B	2014	GOL	O1-C1	4.56	1.61	1.42
5	A	2005	GOL	O1-C1	4.56	1.61	1.42
5	A	2011	GOL	O1-C1	4.57	1.61	1.42
5	A	2009	GOL	O1-C1	4.57	1.61	1.42
5	B	2004	GOL	O1-C1	4.58	1.61	1.42
5	A	2006	GOL	O1-C1	4.59	1.61	1.42
5	A	2010	GOL	O1-C1	4.60	1.61	1.42
5	A	2018	GOL	O1-C1	4.62	1.61	1.42
5	A	2003	GOL	O1-C1	4.97	1.63	1.42

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1002	SO4	O4-S-O3	2.11	118.20	108.83
3	A	1001	SO4	O4-S-O3	2.14	118.37	108.83
5	A	2003	GOL	O1-C1-C2	3.00	124.69	110.11
5	A	2001	GOL	O1-C1-C2	3.09	125.10	110.11
5	A	2011	GOL	O1-C1-C2	3.23	125.76	110.11
5	B	2014	GOL	O1-C1-C2	3.23	125.79	110.11
5	B	2004	GOL	O1-C1-C2	3.23	125.80	110.11
5	A	2010	GOL	O1-C1-C2	3.23	125.80	110.11
5	A	2006	GOL	O1-C1-C2	3.25	125.90	110.11
5	B	2002	GOL	O1-C1-C2	3.26	125.90	110.11
5	A	2013	GOL	O1-C1-C2	3.26	125.91	110.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2008	GOL	O1-C1-C2	3.26	125.92	110.11
5	A	2005	GOL	O1-C1-C2	3.26	125.94	110.11
5	A	2007	GOL	O1-C1-C2	3.28	126.02	110.11
5	A	2012	GOL	O1-C1-C2	3.28	126.03	110.11
5	A	2009	GOL	O1-C1-C2	3.30	126.12	110.11
5	A	2016	GOL	O1-C1-C2	3.31	126.15	110.11
5	A	2018	GOL	O1-C1-C2	3.51	127.13	110.11
5	A	2003	GOL	O2-C2-C3	6.34	137.94	109.00
5	A	2018	GOL	O2-C2-C3	6.43	138.36	109.00
5	A	2011	GOL	O2-C2-C3	6.44	138.42	109.00
5	B	2004	GOL	O2-C2-C3	6.44	138.43	109.00
5	A	2005	GOL	O2-C2-C3	6.45	138.47	109.00
5	B	2014	GOL	O2-C2-C3	6.46	138.50	109.00
5	A	2006	GOL	O2-C2-C3	6.46	138.51	109.00
5	A	2009	GOL	O2-C2-C3	6.47	138.54	109.00
5	A	2010	GOL	O2-C2-C3	6.47	138.54	109.00
5	B	2002	GOL	O2-C2-C3	6.49	138.62	109.00
5	A	2007	GOL	O2-C2-C3	6.50	138.69	109.00
5	A	2008	GOL	O2-C2-C3	6.50	138.71	109.00
5	A	2013	GOL	O2-C2-C3	6.51	138.72	109.00
5	A	2012	GOL	O2-C2-C3	6.51	138.73	109.00
5	A	2016	GOL	O2-C2-C3	6.54	138.87	109.00
5	A	2001	GOL	O2-C2-C3	6.60	139.13	109.00
5	A	2016	GOL	O3-C3-C2	10.38	160.47	110.11
5	A	2008	GOL	O3-C3-C2	10.45	160.82	110.11
5	A	2001	GOL	O3-C3-C2	10.47	160.89	110.11
5	A	2009	GOL	O3-C3-C2	10.48	160.94	110.11
5	A	2012	GOL	O3-C3-C2	10.48	160.94	110.11
5	A	2007	GOL	O3-C3-C2	10.48	160.95	110.11
5	A	2013	GOL	O3-C3-C2	10.51	161.08	110.11
5	A	2006	GOL	O3-C3-C2	10.51	161.12	110.11
5	A	2010	GOL	O3-C3-C2	10.51	161.13	110.11
5	A	2011	GOL	O3-C3-C2	10.52	161.17	110.11
5	B	2014	GOL	O3-C3-C2	10.54	161.24	110.11
5	B	2004	GOL	O3-C3-C2	10.55	161.28	110.11
5	A	2005	GOL	O3-C3-C2	10.55	161.30	110.11
5	B	2002	GOL	O3-C3-C2	10.56	161.35	110.11
5	A	2003	GOL	O3-C3-C2	10.59	161.47	110.11
5	A	2018	GOL	O3-C3-C2	10.62	161.65	110.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	2007	GOL	1	0
5	A	2009	GOL	1	0
5	A	2012	GOL	1	0
5	A	2018	GOL	1	0
5	B	2002	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/275 (99%)	-0.00	7 (2%) 56 62	9, 13, 22, 30	8 (2%)
2	B	76/76 (100%)	0.04	3 (3%) 39 44	12, 17, 27, 30	0
All	All	350/351 (99%)	0.01	10 (2%) 51 57	9, 14, 23, 30	8 (2%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	257	LEU	4.3
1	A	160	GLY	3.4
1	A	275	GLN	3.3
2	B	19	ARG	3.1
1	A	1	ALA	3.0
1	A	159	SER	2.4
1	A	161	SER	2.3
1	A	259	ASP	2.2
2	B	16	ASP	2.2
2	B	39	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSD	A	221	8/9	0.95	0.08	9,11,16,17	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	A	2013	6/6	0.26	0.40	46,46,46,47	0
5	GOL	B	2002	6/6	0.32	0.30	50,50,51,51	0
5	GOL	B	2004	6/6	0.32	0.24	47,48,48,49	0
5	GOL	A	2011	6/6	0.33	0.22	46,46,47,47	0
5	GOL	A	2016	6/6	0.36	0.28	41,41,42,42	0
5	GOL	A	2010	6/6	0.37	0.40	41,45,46,47	0
5	GOL	A	2003	6/6	0.38	0.38	42,46,47,47	0
5	GOL	A	2018	6/6	0.49	0.40	39,40,40,41	0
5	GOL	B	2014	6/6	0.53	0.30	48,48,48,48	0
5	GOL	A	2012	6/6	0.56	0.38	50,50,51,51	0
5	GOL	A	2005	6/6	0.57	0.28	48,49,50,51	0
5	GOL	A	2007	6/6	0.58	0.28	41,47,48,50	0
5	GOL	A	2009	6/6	0.62	0.17	47,47,47,48	0
5	GOL	A	2006	6/6	0.69	0.36	44,47,47,47	0
5	GOL	A	2008	6/6	0.77	0.17	25,33,34,37	0
5	GOL	A	2001	6/6	0.83	0.13	22,23,24,26	0
3	SO4	A	1001	5/5	0.93	0.15	28,28,31,31	0
3	SO4	B	1002	5/5	0.97	0.12	17,21,24,25	0
4	CA	A	3001	1/1	0.99	0.03	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.