



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 01:15 pm GMT

PDB ID : 1V83
Title : Crystal structure of human GlcAT-P in complex with Udp and Mn²⁺
Authors : Kakuda, S.; Shiba, T.; Ishiguro, M.; Tagawa, H.; Oka, S.; Kajihara, Y.;
Kawasaki, T.; Wakatsuki, S.; Kato, R.
Deposited on : 2003-12-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5502 (1.90-1.90)
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	<div> <div>8%</div> <div> <div></div> <div>78%</div> <div>17%</div> <div>• •</div> </div> </div>
1	B	253	<div> <div>10%</div> <div> <div></div> <div>75%</div> <div>17%</div> <div>• 5%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4413 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	0	0
			1990	1269	368	348	5			
1	B	241	Total	C	N	O	S	0	0	0
			1947	1241	359	342	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	ALA	-	CLONING ARTIFACT	UNP Q9P2W7
B	82	ALA	-	CLONING ARTIFACT	UNP Q9P2W7

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
4	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	177	Total 177	O 177	0	0
5	B	237	Total 237	O 237	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1



- Molecule 1: Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.74Å 85.70Å 122.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.90 49.94 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.2 (40.00-1.90) 98.2 (49.94-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.79 (at 1.90Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.201 , 0.229 0.195 , 0.201	Depositor DCC
R_{free} test set	2600 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.932	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4413	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TLA, MN, UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/2039	0.57	0/2771
1	B	0.31	0/1995	0.57	0/2713
All	All	0.31	0/4034	0.57	0/5484

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1990	0	2023	45	0
1	B	1947	0	1976	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	10	0	4	0	0
4	A	25	0	11	2	0
4	B	25	0	11	0	0
5	A	177	0	0	2	0
5	B	237	0	0	1	0
All	All	4413	0	4025	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 11.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:182:PRO:HG2	1:B:185:SER:HB3	1.59	0.84
1:A:171:ASN:HD21	1:A:273:PHE:H	1.28	0.81
1:B:171:ASN:HD21	1:B:273:PHE:H	1.28	0.80
1:A:132:ARG:HD2	5:A:642:HOH:O	1.87	0.73
1:B:182:PRO:HD3	5:B:604:HOH:O	1.88	0.72
1:B:261:ASN:HD22	1:B:263:ARG:H	1.35	0.72
1:A:313:ARG:NH1	1:A:313:ARG:HB3	2.07	0.70
1:A:111:HIS:HE1	1:B:107:ASN:O	1.77	0.67
1:A:231:ARG:NH2	1:A:240:ARG:CB	2.59	0.66
1:B:324:LYS:HA	1:B:324:LYS:HE3	1.81	0.63
1:B:276:ARG:HA	1:B:276:ARG:HE	1.64	0.63
1:A:146:VAL:HB	1:A:172:LEU:HD22	1.81	0.63
1:B:246:ASP:O	1:B:249:ARG:HG3	1.98	0.62
1:A:233:ASN:C	1:A:233:ASN:HD22	2.02	0.62
1:A:148:THR:HB	1:A:153:LYS:HE3	1.81	0.62
1:B:83:LEU:HG	1:B:209:ARG:HD3	1.81	0.62
1:A:218:PRO:HB2	1:A:307:ILE:HD13	1.82	0.61
1:A:240:ARG:HE	1:A:248:HIS:HD2	1.48	0.60
1:A:133:LEU:O	1:A:137:THR:HG23	2.02	0.59
1:A:307:ILE:HG13	1:B:334:ILE:HG22	1.83	0.59
1:B:222:VAL:HG23	1:B:228:GLU:HG3	1.84	0.58
1:A:292:VAL:HG22	1:A:293:THR:N	2.19	0.58
1:A:313:ARG:HB3	1:A:313:ARG:HH11	1.69	0.57
1:B:149:PRO:HD2	1:B:168:MET:SD	2.46	0.56
1:B:261:ASN:ND2	1:B:263:ARG:H	2.04	0.55
1:A:240:ARG:HH21	1:A:240:ARG:HG2	1.72	0.55
1:B:163:ILE:HD13	1:B:163:ILE:H	1.73	0.54
1:B:276:ARG:HA	1:B:276:ARG:NE	2.23	0.53
1:A:222:VAL:HG23	1:A:228:GLU:HG3	1.91	0.53
1:A:116:HIS:HE1	1:A:142:THR:OG1	1.90	0.53
1:A:240:ARG:NH2	1:A:240:ARG:HG2	2.23	0.52
1:B:313:ARG:HH11	1:B:313:ARG:HG3	1.73	0.52
1:A:165:ARG:HE	4:A:503:UDP:H5	1.58	0.51
1:B:133:LEU:O	1:B:137:THR:HG23	2.10	0.51
1:B:116:HIS:HE1	1:B:142:THR:OG1	1.95	0.49
1:A:261:ASN:ND2	1:A:263:ARG:HB3	2.26	0.49
1:B:146:VAL:HB	1:B:172:LEU:HD22	1.94	0.49
1:B:275:LEU:N	1:B:275:LEU:HD23	2.28	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:ASN:HB2	1:A:200:TYR:CZ	2.49	0.48
1:B:129:LEU:C	1:B:129:LEU:HD23	2.33	0.48
1:B:118:LEU:HD13	1:B:173:ALA:HB1	1.96	0.47
1:B:182:PRO:CG	1:B:185:SER:HB3	2.36	0.47
1:B:274:LYS:NZ	1:B:277:GLY:H	2.13	0.47
1:A:233:ASN:C	1:A:233:ASN:ND2	2.68	0.47
1:A:263:ARG:O	1:A:267:GLN:HG2	2.15	0.47
1:B:240:ARG:HH11	1:B:240:ARG:HG2	1.79	0.47
1:A:231:ARG:NH2	1:A:240:ARG:HB2	2.30	0.46
1:B:163:ILE:HG22	1:B:275:LEU:O	2.15	0.46
1:A:126:ARG:HB3	1:A:143:HIS:CE1	2.51	0.46
1:A:233:ASN:ND2	1:A:237:LYS:H	2.12	0.46
1:A:231:ARG:NH2	1:A:240:ARG:HG3	2.30	0.46
1:A:171:ASN:HD21	1:A:273:PHE:N	2.07	0.45
1:A:315:GLU:HB2	1:B:313:ARG:HB2	1.99	0.45
1:B:171:ASN:HD21	1:B:273:PHE:N	2.06	0.45
1:A:329:ASP:HB3	1:A:332:VAL:HG23	1.98	0.44
1:B:329:ASP:HB3	1:B:332:VAL:HG23	1.97	0.44
1:A:164:PRO:HD3	1:A:278:VAL:HG11	1.99	0.44
1:A:87:HIS:ND1	1:A:116:HIS:HD2	2.15	0.44
1:B:276:ARG:HB3	1:B:277:GLY:H	1.58	0.44
1:B:307:ILE:N	1:B:307:ILE:HD12	2.34	0.43
1:B:133:LEU:HD23	1:B:133:LEU:C	2.39	0.43
1:B:87:HIS:ND1	1:B:116:HIS:HD2	2.16	0.42
1:A:110:LEU:HA	1:A:139:LEU:HD21	2.02	0.42
1:A:283:GLN:NE2	5:A:576:HOH:O	2.51	0.42
1:A:132:ARG:HG3	1:A:135:ARG:NH1	2.35	0.42
1:A:292:VAL:HG22	1:A:293:THR:O	2.19	0.42
1:B:183:ARG:NH1	1:B:267:GLN:HA	2.34	0.42
1:A:261:ASN:HD21	1:A:263:ARG:HB3	1.83	0.42
1:B:198:ASN:HB2	1:B:200:TYR:CZ	2.54	0.42
1:B:275:LEU:H	1:B:275:LEU:HD23	1.84	0.42
1:A:164:PRO:HB2	1:A:283:GLN:NE2	2.34	0.42
1:A:240:ARG:HE	1:A:248:HIS:CD2	2.31	0.42
1:B:184:ASN:H	1:B:184:ASN:ND2	2.17	0.42
1:A:231:ARG:NH2	1:A:240:ARG:HB3	2.31	0.42
1:A:116:HIS:CE1	1:A:142:THR:OG1	2.71	0.42
1:A:253:ILE:HD11	1:A:257:GLY:O	2.21	0.41
1:B:109:LEU:HG	1:B:115:LEU:HD11	2.02	0.41
1:B:123:ALA:HB1	1:B:124:PRO:HD2	2.02	0.41
1:B:292:VAL:HG22	1:B:293:THR:N	2.36	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:VAL:HG22	1:A:293:THR:H	1.83	0.41
1:B:116:HIS:CE1	1:B:142:THR:OG1	2.73	0.41
1:A:292:VAL:CG2	1:A:293:THR:N	2.85	0.40
1:A:165:ARG:HB2	4:A:503:UDP:H5'2	2.02	0.40
1:B:246:ASP:HB3	1:B:249:ARG:HH11	1.86	0.40
1:B:246:ASP:N	1:B:247:PRO:HD3	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/253 (95%)	230 (95%)	9 (4%)	2 (1%)	21	10
1	B	237/253 (94%)	223 (94%)	13 (6%)	1 (0%)	36	25
All	All	478/506 (94%)	453 (95%)	22 (5%)	3 (1%)	27	15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	276	ARG
1	A	240	ARG
1	A	278	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/222 (98%)	210 (97%)	7 (3%)	42	33
1	B	212/222 (96%)	204 (96%)	8 (4%)	36	26
All	All	429/444 (97%)	414 (96%)	15 (4%)	39	29

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	ARG
1	A	109	LEU
1	A	140	ASN
1	A	233	ASN
1	A	240	ARG
1	A	283	GLN
1	A	313	ARG
1	B	95	ARG
1	B	109	LEU
1	B	163	ILE
1	B	184	ASN
1	B	261	ASN
1	B	275	LEU
1	B	276	ARG
1	B	324	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	ASN
1	A	111	HIS
1	A	116	HIS
1	A	140	ASN
1	A	151	ASN
1	A	171	ASN
1	A	187	GLN
1	A	198	ASN
1	A	233	ASN
1	A	261	ASN
1	A	283	GLN
1	B	107	ASN
1	B	114	ASN
1	B	116	HIS
1	B	140	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	169	GLN
1	B	171	ASN
1	B	184	ASN
1	B	198	ASN
1	B	261	ASN
1	B	267	GLN
1	B	283	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TLA	A	500	-	3,9,9	0.56	0	6,12,12	1.24	1 (16%)
4	UDP	A	503	2	20,26,26	2.58	4 (20%)	23,40,40	3.82	6 (26%)
4	UDP	B	504	2	20,26,26	2.56	4 (20%)	23,40,40	3.81	5 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TLA	A	500	-	-	0/4/12/12	0/0/0/0
4	UDP	A	503	2	-	0/12/32/32	0/2/2/2
4	UDP	B	504	2	-	0/12/32/32	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	504	UDP	PB-O1B	2.48	1.59	1.50
4	A	503	UDP	PB-O1B	2.98	1.60	1.50
4	A	503	UDP	C4-N3	5.52	1.43	1.33
4	B	504	UDP	C4-N3	5.75	1.43	1.33
4	B	504	UDP	C6-N1	5.94	1.43	1.35
4	A	503	UDP	C6-N1	6.13	1.44	1.35
4	A	503	UDP	C6-C5	6.22	1.51	1.38
4	B	504	UDP	C6-C5	6.28	1.51	1.38

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	504	UDP	C5-C4-N3	-3.17	115.81	123.17
4	A	503	UDP	C5-C4-N3	-3.12	115.92	123.17
3	A	500	TLA	C1-C2-C3	-2.13	108.53	113.11
4	A	503	UDP	O2A-PA-O1A	2.02	122.42	112.14
4	A	503	UDP	PA-O5'-C5'	2.06	133.73	121.68
4	B	504	UDP	PA-O5'-C5'	2.06	133.74	121.68
4	B	504	UDP	O3B-PB-O2B	2.68	118.20	107.59
4	A	503	UDP	O3B-PB-O2B	2.69	118.23	107.59
4	A	503	UDP	O5'-PA-O1A	3.47	122.62	109.07
4	B	504	UDP	O5'-PA-O1A	3.59	123.08	109.07
4	B	504	UDP	C4-N3-C2	16.76	128.56	114.14
4	A	503	UDP	C4-N3-C2	16.87	128.65	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	UDP	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/253 (96%)	0.42	19 (7%) 13 15	13, 23, 47, 66	0
1	B	241/253 (95%)	0.60	25 (10%) 6 7	14, 22, 46, 65	0
All	All	486/506 (96%)	0.51	44 (9%) 9 10	13, 23, 47, 66	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	186	SER	10.2
1	B	278	VAL	9.1
1	A	186	SER	8.7
1	A	185	SER	7.4
1	B	184	ASN	7.4
1	B	277	GLY	7.0
1	B	185	SER	6.7
1	B	162	ARG	6.6
1	B	275	LEU	6.5
1	B	276	ARG	6.4
1	B	82	ALA	6.0
1	B	163	ILE	6.0
1	A	278	VAL	4.6
1	A	187	GLN	4.6
1	B	240	ARG	4.6
1	A	184	ASN	4.4
1	A	324	LYS	4.3
1	B	279	LYS	4.2
1	B	183	ARG	4.2
1	A	240	ARG	4.2
1	B	282	TYR	4.1
1	A	277	GLY	3.7
1	B	280	GLY	3.6
1	A	183	ARG	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	248	HIS	3.5
1	A	325	LYS	3.3
1	A	327	PHE	3.2
1	A	326	GLY	3.2
1	A	162	ARG	3.0
1	B	149	PRO	3.0
1	B	182	PRO	3.0
1	A	212	ARG	2.9
1	A	154	LEU	2.7
1	A	263	ARG	2.6
1	B	245	PHE	2.5
1	B	148	THR	2.5
1	B	274	LYS	2.4
1	B	324	LYS	2.4
1	B	187	GLN	2.3
1	A	182	PRO	2.2
1	A	181	PHE	2.1
1	B	281	GLY	2.1
1	B	246	ASP	2.1
1	A	188	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	UDP	B	504	25/25	0.92	0.15	18,24,35,40	0
4	UDP	A	503	25/25	0.93	0.15	16,22,32,41	0
2	MN	B	502	1/1	0.97	0.10	19,19,19,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TLA	A	500	10/10	0.97	0.09	22,25,27,27	0
2	MN	A	501	1/1	0.99	0.14	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.