



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 10:15 pm GMT

PDB ID : 1VB9  
Title : Crystal structure of Thermoactinomyces vulgaris R-47 alpha-amylase II (TVA II) complexed with transglycosylated product  
Authors : Mizuno, M.; Tono-zuka, T.; Uechi, A.; Ohtaki, A.; Ichikawa, K.; Kamitori, S.; Nishikawa, A.; Sakano, Y.  
Deposited on : 2004-02-25  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	<div> <div>5%</div> <div> <div></div> <div>72%</div> <div>25%</div> <div>••</div> </div> </div>
1	B	585	<div> <div>5%</div> <div> <div></div> <div>75%</div> <div>22%</div> <div>•</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10196 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called alpha-amylase II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	585	Total	C	N	O	S	0	0	0
			4776	3056	832	873	15			
1	B	585	Total	C	N	O	S	0	0	0
			4776	3056	832	873	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	325	ASN	ASP	ENGINEERED	UNP Q08751
B	325	ASN	ASP	ENGINEERED	UNP Q08751

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 11 6 5	0	0
2	A	1	Total C O 11 6 5	0	0
2	A	1	Total C O 11 6 5	0	0
2	A	1	Total C O 12 6 6	0	0
2	A	1	Total C O 11 6 5	0	0
2	B	1	Total C O 11 6 5	0	0
2	B	1	Total C O 11 6 5	0	0
2	B	1	Total C O 11 6 5	0	0
2	B	1	Total C O 11 6 5	0	0
2	B	1	Total C O 12 6 6	0	0
2	B	1	Total C O 11 6 5	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Ca 1 1	0	0
3	A	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	249	Total O 249 249	0	0
4	B	259	Total O 259 259	0	0


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha-amylase II



- Molecule 1: alpha-amylase II





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.05Å 118.27Å 112.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.99 – 2.20 33.98 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (33.99-2.20) 99.9 (33.98-2.20)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.29 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.194 , 0.233 0.202 , 0.240	Depositor DCC
$R_{free}$ test set	7737 reflections (10.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.2	Xtriage
Anisotropy	0.397	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.008 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10196	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/4906	0.62	3/6641 (0.0%)
1	B	0.35	0/4906	0.60	1/6641 (0.0%)
All	All	0.35	0/9812	0.61	4/13282 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	277	LYS	N-CA-C	-7.84	89.83	111.00
1	A	276	SER	N-CA-C	5.99	127.18	111.00
1	A	279	SER	N-CA-C	-5.39	96.45	111.00
1	B	323	ARG	N-CA-C	-5.08	97.29	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4776	0	4609	143	0
1	B	4776	0	4609	129	0
2	A	67	0	57	4	0
2	B	67	0	57	4	0
3	A	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	A	249	0	0	6	0
4	B	259	0	0	14	0
All	All	10196	0	9332	269	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 269 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:VAL:HG11	1:A:351:ILE:HD11	1.46	0.97
1:B:293:MET:HE1	2:B:702:GLC:H62	1.45	0.97
1:A:293:MET:HE1	2:A:702:GLC:H62	1.48	0.94
1:A:585:ARG:HH11	1:A:585:ARG:HB3	1.35	0.89
1:B:198:SER:HB3	1:B:203:LYS:HG3	1.57	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/585 (100%)	556 (95%)	24 (4%)	3 (0%)	31	33
1	B	583/585 (100%)	556 (95%)	22 (4%)	5 (1%)	19	17
All	All	1166/1170 (100%)	1112 (95%)	46 (4%)	8 (1%)	24	23

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	279	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	275	VAL
1	B	278	THR
1	A	275	VAL
1	B	281	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/493 (100%)	468 (95%)	25 (5%)	26	31
1	B	493/493 (100%)	474 (96%)	19 (4%)	35	44
All	All	986/986 (100%)	942 (96%)	44 (4%)	30	37

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	483	LEU
1	A	585	ARG
1	B	542	LEU
1	A	504	ASN
1	A	573	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	563	HIS
1	B	244	HIS
1	B	524	GLN
1	B	135	GLN
1	B	257	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	A	701	2	11,11,12	0.46	0	15,15,17	0.55	0
2	GLC	A	702	2	11,11,12	0.50	0	15,15,17	0.66	0
2	GLC	A	703	2	11,11,12	0.70	0	15,15,17	1.54	2 (13%)
2	GLC	A	704	2	11,11,12	0.65	0	15,15,17	0.66	1 (6%)
2	GLC	A	705	2	12,12,12	0.45	0	17,17,17	0.36	0
2	GLC	A	706	2	11,11,12	0.49	0	15,15,17	0.53	0
2	GLC	B	701	2	11,11,12	0.46	0	15,15,17	0.50	0
2	GLC	B	702	2	11,11,12	0.54	0	15,15,17	0.59	0
2	GLC	B	703	2	11,11,12	0.49	0	15,15,17	1.12	2 (13%)
2	GLC	B	704	2	11,11,12	0.57	0	15,15,17	0.70	1 (6%)
2	GLC	B	705	2	12,12,12	0.39	0	17,17,17	0.34	0
2	GLC	B	706	2	11,11,12	0.48	0	15,15,17	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	701	2	-	0/2/19/22	0/1/1/1
2	GLC	A	702	2	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	703	2	-	0/2/19/22	0/1/1/1
2	GLC	A	704	2	-	0/2/19/22	0/1/1/1
2	GLC	A	705	2	-	0/2/22/22	0/1/1/1
2	GLC	A	706	2	-	0/2/19/22	0/1/1/1
2	GLC	B	701	2	-	0/2/19/22	0/1/1/1
2	GLC	B	702	2	-	0/2/19/22	0/1/1/1
2	GLC	B	703	2	-	0/2/19/22	0/1/1/1
2	GLC	B	704	2	-	0/2/19/22	0/1/1/1
2	GLC	B	705	2	-	0/2/22/22	0/1/1/1
2	GLC	B	706	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	703	GLC	C1-C2-C3	-2.74	106.19	109.66
2	B	703	GLC	C1-C2-C3	-2.35	106.69	109.66
2	B	704	GLC	C1-O5-C5	2.09	115.07	112.19
2	A	704	GLC	C1-O5-C5	2.14	115.13	112.19
2	B	703	GLC	C1-O5-C5	2.58	115.74	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	702	GLC	3	0
2	A	703	GLC	1	0
2	B	702	GLC	3	0
2	B	703	GLC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	585/585 (100%)	-0.01	28 (4%)	30 29	18, 28, 48, 96	0
1	B	585/585 (100%)	0.02	32 (5%)	25 24	19, 29, 51, 106	0
All	All	1170/1170 (100%)	0.00	60 (5%)	28 26	18, 28, 49, 106	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	SER	16.3
1	B	279	SER	14.5
1	B	273	PHE	11.5
1	B	280	ARG	9.4
1	B	277	LYS	8.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	B	706	11/12	0.68	0.26	56,59,60,61	0
2	GLC	B	705	12/12	0.76	0.20	46,51,54,54	0
2	GLC	A	706	11/12	0.85	0.23	48,52,54,55	0
2	GLC	A	703	11/12	0.86	0.18	24,28,31,42	0
2	GLC	B	704	11/12	0.87	0.16	37,43,47,52	0
2	GLC	A	705	12/12	0.87	0.17	36,42,45,47	0
2	GLC	B	703	11/12	0.89	0.16	28,32,36,46	0
2	GLC	A	701	11/12	0.91	0.15	30,35,40,46	0
2	GLC	B	701	11/12	0.93	0.14	37,39,42,43	0
2	GLC	A	704	11/12	0.94	0.10	31,34,37,43	0
2	GLC	B	702	11/12	0.95	0.09	30,30,32,36	0
2	GLC	A	702	11/12	0.97	0.08	26,28,30,31	0
3	CA	B	602	1/1	0.99	0.04	34,34,34,34	0
3	CA	A	601	1/1	0.99	0.06	26,26,26,26	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.