



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 11, 2019 – 06:53 PM EST

PDB ID : 5W3Z  
Title : Crystal structure of SsoPox AsC6 mutant (L72I-Y99F-I122L-L228M-F229S-W263L)  
Authors : Hiblot, J.; Gotthard, G.; Jacquet, P.; Daude, D.; Bergonzi, C.; Chabriere, E.; Elias, M.  
Deposited on : 2017-06-08  
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.4  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1053 (2.56-2.52)
Clashscore	122126	1098 (2.56-2.52)
Ramachandran outliers	120053	1088 (2.56-2.52)
Sidechain outliers	120020	1088 (2.56-2.52)
RSRZ outliers	108989	1043 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	<div> <div> <div></div> <div>75%</div> <div>16%</div> <div>• 7%</div> </div> </div>
1	B	314	<div> <div>5%</div> <div>73%</div> <div>19%</div> <div>• • •</div> </div>
1	C	314	<div> <div>4%</div> <div>70%</div> <div>22%</div> <div>• 6%</div> </div>
1	D	314	<div> <div>14%</div> <div>69%</div> <div>25%</div> <div>5%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9811 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aryldialkylphosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	0
			2333	1490	397	440	6			
1	B	302	Total	C	N	O	S	0	0	0
			2403	1535	407	453	8			
1	C	296	Total	C	N	O	S	0	0	0
			2364	1510	404	443	7			
1	D	314	Total	C	N	O	S	0	0	0
			2501	1600	425	468	8			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	72	ILE	LEU	engineered mutation	UNP Q97VT7
A	99	PHE	TYR	engineered mutation	UNP Q97VT7
A	122	LEU	ILE	engineered mutation	UNP Q97VT7
A	228	MET	LEU	engineered mutation	UNP Q97VT7
A	229	SER	PHE	engineered mutation	UNP Q97VT7
A	263	LEU	TRP	engineered mutation	UNP Q97VT7
B	72	ILE	LEU	engineered mutation	UNP Q97VT7
B	99	PHE	TYR	engineered mutation	UNP Q97VT7
B	122	LEU	ILE	engineered mutation	UNP Q97VT7
B	228	MET	LEU	engineered mutation	UNP Q97VT7
B	229	SER	PHE	engineered mutation	UNP Q97VT7
B	263	LEU	TRP	engineered mutation	UNP Q97VT7
C	72	ILE	LEU	engineered mutation	UNP Q97VT7
C	99	PHE	TYR	engineered mutation	UNP Q97VT7
C	122	LEU	ILE	engineered mutation	UNP Q97VT7
C	228	MET	LEU	engineered mutation	UNP Q97VT7
C	229	SER	PHE	engineered mutation	UNP Q97VT7
C	263	LEU	TRP	engineered mutation	UNP Q97VT7
D	72	ILE	LEU	engineered mutation	UNP Q97VT7
D	99	PHE	TYR	engineered mutation	UNP Q97VT7
D	122	LEU	ILE	engineered mutation	UNP Q97VT7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	228	MET	LEU	engineered mutation	UNP Q97VT7
D	229	SER	PHE	engineered mutation	UNP Q97VT7
D	263	LEU	TRP	engineered mutation	UNP Q97VT7


- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Fe 1 1	0	0
2	A	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Co 1 1	0	0
3	A	1	Total Co 1 1	0	0
3	D	1	Total Co 1 1	0	0
3	C	1	Total Co 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	52	Total	O	0	0
			52	52		
6	B	33	Total	O	0	0
			33	33		
6	C	39	Total	O	0	0
			39	39		
6	D	30	Total	O	0	0
			30	30		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Aryldialkylphosphatase




#### • Molecule 1: Aryldialkylphosphatase




#### • Molecule 1: Aryldialkylphosphatase





● Molecule 1: Aryldialkylphosphatase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.65Å 105.80Å 154.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.31 – 2.55 46.31 – 2.55	Depositor EDS
% Data completeness (in resolution range)	97.3 (46.31-2.55) 97.3 (46.31-2.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.99 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.189 , 0.252 0.204 , 0.263	Depositor DCC
$R_{free}$ test set	2161 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.8	Xtriage
Anisotropy	0.244	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 64.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9811	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, KCX, CO, EDO, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.99	2/2361 (0.1%)	1.04	4/3182 (0.1%)
1	B	0.92	1/2433 (0.0%)	1.07	11/3281 (0.3%)
1	C	0.98	3/2394 (0.1%)	1.06	7/3227 (0.2%)
1	D	0.89	2/2536 (0.1%)	0.99	5/3422 (0.1%)
All	All	0.94	8/9724 (0.1%)	1.04	27/13112 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	307	GLU	CD-OE1	6.45	1.32	1.25
1	D	162	GLU	CD-OE1	6.21	1.32	1.25
1	D	278	TRP	CB-CG	5.48	1.60	1.50
1	C	117	LEU	C-O	5.32	1.33	1.23
1	B	110	SER	CB-OG	-5.23	1.35	1.42

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	121	ASP	CB-CG-OD1	11.55	128.69	118.30
1	C	121	ASP	CB-CG-OD2	-9.92	109.37	118.30
1	D	102	LEU	CA-CB-CG	7.71	133.03	115.30
1	C	74	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	B	77	ARG	NE-CZ-NH1	6.83	123.72	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	221	LEU	Peptide
1	B	221	LEU	Peptide
1	C	221	LEU	Peptide
1	D	221	LEU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2333	0	2356	38	0
1	B	2403	0	2426	40	0
1	C	2364	0	2384	45	0
1	D	2501	0	2536	53	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	1	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	4	0	6	1	0
4	B	4	0	6	0	0
4	C	8	0	12	1	0
4	D	8	0	12	1	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
5	C	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	6	0	8	0	0
6	A	52	0	0	1	0
6	B	33	0	0	4	0
6	C	39	0	0	2	0
6	D	30	0	0	7	0
All	All	9811	0	9770	177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:LEU:HD12	1:B:133:ALA:HB3	1.61	0.82
1:D:94:THR:O	1:D:136:VAL:HA	1.80	0.81
1:B:257:TYR:O	1:B:259:CYS:N	2.16	0.79
1:C:14:LYS:HA	1:C:310:LYS:HD2	1.64	0.79
1:C:256:ASP:O	1:C:257:TYR:HB2	1.87	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/314 (92%)	275 (96%)	11 (4%)	2 (1%)	24	34
1	B	297/314 (95%)	267 (90%)	24 (8%)	6 (2%)	8	9
1	C	291/314 (93%)	263 (90%)	23 (8%)	5 (2%)	10	13
1	D	311/314 (99%)	268 (86%)	29 (9%)	14 (4%)	3	1
All	All	1187/1256 (94%)	1073 (90%)	87 (7%)	27 (2%)	7	7

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	GLU
1	B	258	CYS
1	B	260	THR
1	C	205	ASN
1	C	237	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/272 (93%)	240 (94%)	14 (6%)	24	33
1	B	262/272 (96%)	247 (94%)	15 (6%)	23	31
1	C	257/272 (94%)	246 (96%)	11 (4%)	32	44
1	D	272/272 (100%)	251 (92%)	21 (8%)	14	19
All	All	1045/1088 (96%)	984 (94%)	61 (6%)	22	30

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	297	ASN
1	C	186	THR
1	D	270	TYR
1	B	300	VAL
1	C	109	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	131	ASN
1	D	38	HIS
1	D	182	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	KCX	A	137	1,2	8,11,12	0.91	0	5,12,14	1.95	2 (40%)
1	KCX	B	137	1,3,2	8,11,12	1.58	2 (25%)	5,12,14	1.38	1 (20%)
1	KCX	C	137	1,3,2	8,11,12	0.50	0	5,12,14	2.51	2 (40%)
1	KCX	D	137	1,3,2	8,11,12	0.72	0	5,12,14	1.50	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	137	1,2	-	2/6/10/12	-
1	KCX	B	137	1,3,2	-	3/6/10/12	-
1	KCX	C	137	1,3,2	-	3/6/10/12	-
1	KCX	D	137	1,3,2	-	3/6/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	137	KCX	CB-CA	2.91	1.57	1.53
1	B	137	KCX	CA-C	2.80	1.53	1.50

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	137	KCX	CD-CE-NZ	3.95	122.25	111.50
1	C	137	KCX	CE-NZ-CX	-3.54	118.75	123.28
1	A	137	KCX	CD-CE-NZ	3.43	120.85	111.50
1	A	137	KCX	O-C-CA	-2.43	118.47	124.98
1	B	137	KCX	O-C-CA	-2.39	118.57	124.98

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	137	KCX	N-CA-CB-CG
1	A	137	KCX	C-CA-CB-CG
1	B	137	KCX	C-CA-CB-CG
1	C	137	KCX	N-CA-CB-CG
1	C	137	KCX	C-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	137	KCX	1	0
1	D	137	KCX	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 8 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	A	403	-	3,3,3	1.25	0	2,2,2	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	404	-	5,5,5	0.40	0	5,5,5	0.72	0
4	EDO	B	403	-	3,3,3	1.01	0	2,2,2	0.41	0
5	GOL	B	404	-	5,5,5	0.33	0	5,5,5	0.52	0
4	EDO	C	403	-	3,3,3	0.74	0	2,2,2	0.66	0
4	EDO	C	404	-	3,3,3	0.70	0	2,2,2	0.29	0
5	GOL	C	405	-	5,5,5	0.32	0	5,5,5	0.46	0
4	EDO	D	403	-	3,3,3	0.55	0	2,2,2	0.48	0
5	GOL	D	404	-	5,5,5	0.28	0	5,5,5	0.25	0
4	EDO	D	405	-	3,3,3	0.85	0	2,2,2	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	403	-	-	1/1/1/1	-
5	GOL	A	404	-	-	2/4/4/4	-
4	EDO	B	403	-	-	1/1/1/1	-
5	GOL	B	404	-	-	2/4/4/4	-
4	EDO	C	403	-	-	1/1/1/1	-
4	EDO	C	404	-	-	0/1/1/1	-
5	GOL	C	405	-	-	4/4/4/4	-
4	EDO	D	403	-	-	1/1/1/1	-
5	GOL	D	404	-	-	2/4/4/4	-
4	EDO	D	405	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	404	GOL	C1-C2-C3-O3
5	D	404	GOL	O1-C1-C2-O2
5	D	404	GOL	O1-C1-C2-C3
5	B	404	GOL	C1-C2-C3-O3
5	A	404	GOL	O2-C2-C3-O3

There are no ring outliers.



3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	EDO	1	0
4	C	404	EDO	1	0
4	D	403	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/314 (92%)	0.33	3 (1%) 82 87	34, 61, 98, 135	0
1	B	301/314 (95%)	0.51	17 (5%) 24 28	34, 69, 119, 168	0
1	C	295/314 (93%)	0.43	14 (4%) 31 37	39, 76, 108, 135	0
1	D	313/314 (99%)	1.10	44 (14%) 2 3	44, 94, 137, 187	0
All	All	1201/1256 (95%)	0.60	78 (6%) 19 22	34, 75, 124, 187	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	272	PRO	15.3
1	D	270	TYR	11.9
1	D	275	ALA	11.2
1	D	278	TRP	10.9
1	D	269	GLU	10.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	B	137	12/13	0.96	0.15	36,38,40,41	0
1	KCX	A	137	12/13	0.97	0.16	35,35,36,36	0
1	KCX	C	137	12/13	0.97	0.17	36,48,55,57	0
1	KCX	D	137	12/13	0.98	0.15	66,74,75,78	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	D	405	4/4	0.83	0.30	31,32,33,33	0
5	GOL	C	405	6/6	0.85	0.32	49,53,55,55	0
4	EDO	C	403	4/4	0.88	0.31	47,52,53,54	0
4	EDO	D	403	4/4	0.90	0.20	48,56,58,61	0
5	GOL	D	404	6/6	0.91	0.21	35,37,38,40	0
4	EDO	C	404	4/4	0.93	0.12	28,30,32,36	0
5	GOL	B	404	6/6	0.93	0.19	35,37,37,38	0
4	EDO	B	403	4/4	0.94	0.31	34,35,37,38	0
4	EDO	A	403	4/4	0.94	0.12	24,27,28,29	0
5	GOL	A	404	6/6	0.95	0.16	27,30,31,31	0
3	CO	D	402	1/1	0.96	0.07	39,39,39,39	0
2	FE	D	401	1/1	0.98	0.07	30,30,30,30	0
2	FE	C	401	1/1	0.99	0.07	20,20,20,20	0
2	FE	A	401	1/1	0.99	0.08	15,15,15,15	0
3	CO	B	402	1/1	0.99	0.06	22,22,22,22	0
3	CO	C	402	1/1	0.99	0.06	14,14,14,14	0
2	FE	B	401	1/1	0.99	0.09	16,16,16,16	0
3	CO	A	402	1/1	1.00	0.05	16,16,16,16	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.