



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2018 – 02:21 pm GMT

PDB ID : 2WS3
Title : Crystal structure of the E. coli succinate:quinone oxidoreductase (SQR) SdhD Tyr83Phe mutant
Authors : Ruprecht, J.; Yankovskaya, V.; Maklashina, E.; Iwata, S.; Cecchini, G.
Deposited on : 2009-09-03
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1121 (3.22-3.18)
Clashscore	122126	1091 (3.20-3.20)
Ramachandran outliers	120053	1074 (3.20-3.20)
Sidechain outliers	120020	1073 (3.20-3.20)
RSRZ outliers	108989	1083 (3.22-3.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 87%, yellow 87%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 87% 12% </div> </div>
1	E	588	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 4%, green 4%, green 88%, yellow 88%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 4% 88% 12% </div> </div>
1	I	588	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 21%, green 21%, green 88%, yellow 88%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 21% 88% 11% </div> </div>
2	B	238	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, green 3%, green 87%, yellow 87%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 87% 12% </div> </div>
2	F	238	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, green 3%, green 87%, yellow 87%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 87% 13% </div> </div>
2	J	238	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 13%, green 13%, green 88%, yellow 88%, yellow 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 13% 88% 12% </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	129	
3	G	129	
3	K	129	
4	D	115	
4	H	115	
4	L	115	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	F3S	F	304	-	-	X	-
7	TEO	E	1590	-	-	X	-
7	TEO	I	1590	-	-	X	-
9	SF4	J	303	-	-	X	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 24900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUB-UNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	588	Total	C	N	O	S	0	0	0
			4522	2812	821	861	28			
1	E	588	Total	C	N	O	S	0	0	0
			4522	2812	821	861	28			
1	I	588	Total	C	N	O	S	0	0	0
			4522	2812	821	861	28			

- Molecule 2 is a protein called SUCCINATE DEHYDROGENASE IRON-SULFUR SUB-UNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	238	Total	C	N	O	S	0	0	0
			1869	1172	329	348	20			
2	F	238	Total	C	N	O	S	0	0	0
			1869	1172	329	348	20			
2	J	238	Total	C	N	O	S	0	0	0
			1869	1172	329	348	20			

- Molecule 3 is a protein called SUCCINATE DEHYDROGENASE CYTOCHROME B-556 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	121	Total	C	N	O	S	0	0	0
			933	619	151	158	5			
3	G	121	Total	C	N	O	S	0	0	0
			933	619	151	158	5			
3	K	121	Total	C	N	O	S	0	0	0
			933	619	151	158	5			


- Molecule 4 is a protein called SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	105	Total	C	N	O	S	0	0	0
			835	577	123	132	3			
4	H	105	Total	C	N	O	S	0	0	0
			835	577	123	132	3			
4	L	105	Total	C	N	O	S	0	0	0
			835	577	123	132	3			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	83	PHE	TYR	engineered mutation	UNP P0AC44
H	83	PHE	TYR	engineered mutation	UNP P0AC44
L	83	PHE	TYR	engineered mutation	UNP P0AC44

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
5	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
5	I	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	I	1	Total	Na	0	0
			1	1		
6	A	1	Total	Na	0	0
			1	1		
6	E	1	Total	Na	0	0
			1	1		

- Molecule 7 is MALATE LIKE INTERMEDIATE (three-letter code: TEO) (formula: $C_4H_4O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			9	4	5		
7	E	1	Total	C	O	0	0
			9	4	5		
7	I	1	Total	C	O	0	0
			9	4	5		

- Molecule 8 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			4	2	2		
8	F	1	Total	Fe	S	0	0
			4	2	2		
8	J	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 9 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			8	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	F	1	Total	Fe	S	0	0
			8	4	4		
9	J	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 10 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	Fe	S	0	0
			7	3	4		
10	F	1	Total	Fe	S	0	0
			7	3	4		
10	J	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 11 is 2-METHYL-N-PHENYL-5,6-DIHYDRO-1,4-OXATHIINE-3-CARBOXAMIDE (three-letter code: CBE) (formula: $\text{C}_{12}\text{H}_{13}\text{NO}_2\text{S}$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	C	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
11	G	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
11	K	1	Total	C	N	O	S	0	0
			16	12	1	2	1		

- Molecule 12 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
12	G	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
12	K	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUBUNIT




• Molecule 1: SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUBUNIT





• Molecule 1: SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUBUNIT









• Molecule 2: SUCCINATE DEHYDROGENASE IRON-SULFUR SUBUNIT





• Molecule 2: SUCCINATE DEHYDROGENASE IRON-SULFUR SUBUNIT



• Molecule 2: SUCCINATE DEHYDROGENASE IRON-SULFUR SUBUNIT



• Molecule 3: SUCCINATE DEHYDROGENASE CYTOCHROME B-556 SUBUNIT





• Molecule 3: SUCCINATE DEHYDROGENASE CYTOCHROME B-556 SUBUNIT







• Molecule 3: SUCCINATE DEHYDROGENASE CYTOCHROME B-556 SUBUNIT





• Molecule 4: SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR SUBUNIT



• Molecule 4: SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR SUBUNIT



• Molecule 4: SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	119.85Å 184.71Å 203.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.03 – 3.20 49.01 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.03-3.20) 99.8 (49.01-3.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.219 , 0.253 0.210 , 0.243	Depositor DCC
R_{free} test set	3797 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	84.8	Xtriage
Anisotropy	0.360	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	24900	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TEO, NA, SF4, CBE, F3S, FES, HEM, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	1/4611 (0.0%)	0.67	0/6237
1	E	0.59	0/4611	0.64	0/6237
1	I	0.46	0/4611	0.58	0/6237
2	B	0.72	2/1908 (0.1%)	0.69	0/2578
2	F	0.61	0/1908	0.68	0/2578
2	J	0.51	0/1908	0.61	0/2578
3	C	0.66	0/953	0.62	0/1293
3	G	0.62	0/953	0.61	1/1293 (0.1%)
3	K	0.55	0/953	0.56	0/1293
4	D	0.68	0/858	0.59	0/1173
4	H	0.64	0/858	0.59	0/1173
4	L	0.61	0/858	0.58	0/1173
All	All	0.59	3/24990 (0.0%)	0.63	1/33843 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	154	CYS	CB-SG	-5.44	1.73	1.81
1	A	353	CYS	CB-SG	-5.42	1.73	1.81
2	B	216	CYS	CB-SG	-5.39	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	19	ARG	NE-CZ-NH1	5.54	123.07	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4522	0	4426	52	0
1	E	4522	0	4426	43	0
1	I	4522	0	4426	46	0
2	B	1869	0	1850	12	0
2	F	1869	0	1850	14	0
2	J	1869	0	1850	15	0
3	C	933	0	979	4	0
3	G	933	0	979	3	0
3	K	933	0	979	6	0
4	D	835	0	875	6	0
4	H	835	0	875	5	0
4	L	835	0	875	6	0
5	A	53	0	31	9	0
5	E	53	0	31	10	0
5	I	53	0	31	17	0
6	A	1	0	0	0	0
6	E	1	0	0	0	0
6	I	1	0	0	0	0
7	A	9	0	3	3	0
7	E	9	0	3	5	0
7	I	9	0	3	6	0
8	B	4	0	0	0	0
8	F	4	0	0	0	0
8	J	4	0	0	1	0
9	B	8	0	0	0	0
9	F	8	0	0	0	0
9	J	8	0	0	3	0
10	B	7	0	0	1	0
10	F	7	0	0	2	0
10	J	7	0	0	1	0
11	C	16	0	13	2	0
11	G	16	0	13	0	0
11	K	16	0	13	2	0
12	C	43	0	30	7	0
12	G	43	0	30	6	0
12	K	43	0	30	12	0
All	All	24900	0	24621	223	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 223 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:45:HIS:NE2	5:E:601:FAD:HM82	1.36	1.38
1:A:45:HIS:NE2	5:A:601:FAD:C8M	1.94	1.30
1:A:45:HIS:NE2	5:A:601:FAD:HM82	1.42	1.30
1:E:45:HIS:NE2	5:E:601:FAD:C8M	2.08	1.15
1:A:45:HIS:CE1	5:A:601:FAD:HM82	1.88	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	586/588 (100%)	560 (96%)	24 (4%)	2 (0%)	43	79
1	E	586/588 (100%)	558 (95%)	26 (4%)	2 (0%)	43	79
1	I	586/588 (100%)	561 (96%)	23 (4%)	2 (0%)	43	79
2	B	236/238 (99%)	223 (94%)	12 (5%)	1 (0%)	36	74
2	F	236/238 (99%)	220 (93%)	15 (6%)	1 (0%)	36	74
2	J	236/238 (99%)	223 (94%)	12 (5%)	1 (0%)	36	74
3	C	119/129 (92%)	114 (96%)	5 (4%)	0	100	100
3	G	119/129 (92%)	115 (97%)	4 (3%)	0	100	100
3	K	119/129 (92%)	116 (98%)	3 (2%)	0	100	100
4	D	103/115 (90%)	98 (95%)	5 (5%)	0	100	100
4	H	103/115 (90%)	98 (95%)	5 (5%)	0	100	100
4	L	103/115 (90%)	98 (95%)	5 (5%)	0	100	100
All	All	3132/3210 (98%)	2984 (95%)	139 (4%)	9 (0%)	43	79

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	GLY
1	E	92	GLY
1	I	92	GLY
1	A	472	SER
1	E	472	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	473/473 (100%)	464 (98%)	9 (2%)	60	84
1	E	473/473 (100%)	464 (98%)	9 (2%)	60	84
1	I	473/473 (100%)	465 (98%)	8 (2%)	63	86
2	B	208/208 (100%)	197 (95%)	11 (5%)	25	62
2	F	208/208 (100%)	198 (95%)	10 (5%)	28	65
2	J	208/208 (100%)	199 (96%)	9 (4%)	32	68
3	C	101/109 (93%)	99 (98%)	2 (2%)	58	83
3	G	101/109 (93%)	100 (99%)	1 (1%)	78	92
3	K	101/109 (93%)	100 (99%)	1 (1%)	78	92
4	D	88/96 (92%)	86 (98%)	2 (2%)	53	81
4	H	88/96 (92%)	86 (98%)	2 (2%)	53	81
4	L	88/96 (92%)	86 (98%)	2 (2%)	53	81
All	All	2610/2658 (98%)	2544 (98%)	66 (2%)	50	80

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	378	VAL
2	F	120	TYR
2	J	161	SER
1	E	451	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	24	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	78	GLN
4	L	78	GLN
4	H	78	GLN
4	D	14	HIS
1	E	449	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 3 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	TEO	A	1590	-	1,8,8	0.10	0	0,10,10	0.00	-
5	FAD	A	601	-	51,58,58	1.41	8 (15%)	57,89,89	2.29	11 (19%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	FES	B	302	2	0,4,4	0.00	-	0,4,4	0.00	-
9	SF4	B	303	2	0,12,12	0.00	-	0,24,24	0.00	-
10	F3S	B	304	2	0,9,9	0.00	-	0,15,15	0.00	-
11	CBE	C	1129	-	15,17,17	1.55	1 (6%)	15,22,22	2.66	2 (13%)
12	HEM	C	1130	3,4	27,50,50	2.18	5 (18%)	17,82,82	1.83	5 (29%)
7	TEO	E	1590	-	1,8,8	0.50	0	0,10,10	0.00	-
5	FAD	E	601	-	51,58,58	1.50	8 (15%)	57,89,89	2.24	13 (22%)
8	FES	F	302	2	0,4,4	0.00	-	0,4,4	0.00	-
9	SF4	F	303	2	0,12,12	0.00	-	0,24,24	0.00	-
10	F3S	F	304	2	0,9,9	0.00	-	0,15,15	0.00	-
11	CBE	G	1129	-	15,17,17	1.50	2 (13%)	15,22,22	2.56	3 (20%)
12	HEM	G	1130	3,4	27,50,50	2.37	8 (29%)	17,82,82	1.28	3 (17%)
7	TEO	I	1590	-	1,8,8	0.85	0	0,10,10	0.00	-
5	FAD	I	601	-	51,58,58	1.44	6 (11%)	57,89,89	2.07	7 (12%)
8	FES	J	302	2	0,4,4	0.00	-	0,4,4	0.00	-
9	SF4	J	303	2	0,12,12	0.00	-	0,24,24	0.00	-
10	F3S	J	304	2	0,9,9	0.00	-	0,15,15	0.00	-
11	CBE	K	1129	-	15,17,17	1.76	3 (20%)	15,22,22	2.21	2 (13%)
12	HEM	K	1130	3,4	27,50,50	2.41	7 (25%)	17,82,82	2.11	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	TEO	A	1590	-	-	0/2/8/8	0/0/0/0
5	FAD	A	601	-	-	0/28/50/50	0/6/6/6
8	FES	B	302	2	-	0/0/4/4	0/1/1/1
9	SF4	B	303	2	-	0/0/48/48	0/6/5/5
10	F3S	B	304	2	-	0/0/24/24	0/3/3/3
11	CBE	C	1129	-	-	0/6/19/19	0/1/2/2
12	HEM	C	1130	3,4	-	0/6/54/54	0/0/8/8
7	TEO	E	1590	-	-	0/2/8/8	0/0/0/0
5	FAD	E	601	-	-	0/28/50/50	0/6/6/6
8	FES	F	302	2	-	0/0/4/4	0/1/1/1
9	SF4	F	303	2	-	0/0/48/48	0/6/5/5
10	F3S	F	304	2	-	0/0/24/24	0/3/3/3
11	CBE	G	1129	-	-	0/6/19/19	0/1/2/2
12	HEM	G	1130	3,4	-	0/6/54/54	0/0/8/8
7	TEO	I	1590	-	-	0/2/8/8	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FAD	I	601	-	-	0/28/50/50	0/6/6/6
8	FES	J	302	2	-	0/0/4/4	0/1/1/1
9	SF4	J	303	2	-	0/0/48/48	0/6/5/5
10	F3S	J	304	2	-	0/0/24/24	0/3/3/3
11	CBE	K	1129	-	-	0/6/19/19	0/1/2/2
12	HEM	K	1130	3,4	-	0/6/54/54	0/0/8/8

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	K	1130	HEM	C3C-C2C	-5.17	1.33	1.40
12	C	1130	HEM	C3C-C2C	-4.95	1.33	1.40
12	C	1130	HEM	C3B-C2B	-4.86	1.33	1.40
12	G	1130	HEM	C3B-C2B	-4.76	1.33	1.40
12	K	1130	HEM	C3B-C2B	-4.68	1.33	1.40

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	601	FAD	N3A-C2A-N1A	-11.29	119.20	128.86
5	I	601	FAD	N3A-C2A-N1A	-10.65	119.75	128.86
5	E	601	FAD	N3A-C2A-N1A	-10.40	119.96	128.86
11	C	1129	CBE	C3-C8-N10	-4.62	107.47	115.94
12	K	1130	HEM	C1D-C2D-C3D	-4.00	104.21	107.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

16 monomers are involved in 81 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1590	TEO	3	0
5	A	601	FAD	9	0
10	B	304	F3S	1	0
11	C	1129	CBE	2	0
12	C	1130	HEM	7	0
7	E	1590	TEO	5	0
5	E	601	FAD	10	0
10	F	304	F3S	2	0
12	G	1130	HEM	6	0
7	I	1590	TEO	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	I	601	FAD	17	0
8	J	302	FES	1	0
9	J	303	SF4	3	0
10	J	304	F3S	1	0
11	K	1129	CBE	2	0
12	K	1130	HEM	12	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	588/588 (100%)	-0.05	8 (1%) 75 63	45, 63, 84, 97	0
1	E	588/588 (100%)	0.05	22 (3%) 41 27	53, 72, 95, 112	0
1	I	588/588 (100%)	1.09	125 (21%) 1 0	91, 124, 168, 190	0
2	B	238/238 (100%)	-0.01	8 (3%) 45 30	45, 61, 88, 107	0
2	F	238/238 (100%)	-0.00	6 (2%) 57 43	54, 72, 104, 123	0
2	J	238/238 (100%)	0.45	30 (12%) 3 2	75, 96, 165, 189	0
3	C	121/129 (93%)	-0.07	4 (3%) 46 30	62, 76, 107, 120	0
3	G	121/129 (93%)	0.18	3 (2%) 57 43	71, 94, 119, 126	0
3	K	121/129 (93%)	0.50	12 (9%) 7 4	93, 118, 135, 144	0
4	D	105/115 (91%)	-0.12	2 (1%) 66 53	55, 75, 109, 124	0
4	H	105/115 (91%)	0.14	6 (5%) 24 13	63, 79, 137, 155	0
4	L	105/115 (91%)	0.10	8 (7%) 14 7	73, 90, 142, 165	0
All	All	3156/3210 (98%)	0.26	234 (7%) 14 8	45, 80, 147, 190	0

The worst 5 of 234 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	48	SER	6.8
1	I	499	LEU	6.8
1	I	500	ASP	6.4
1	I	203	GLY	6.1
1	I	268	HIS	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	TEO	I	1590	9/9	0.88	0.31	109,112,114,114	0
6	NA	I	1589	1/1	0.91	0.47	53,53,53,53	0
11	CBE	K	1129	16/16	0.91	0.36	85,91,97,97	0
5	FAD	I	601	53/53	0.91	0.47	106,112,121,125	0
8	FES	J	302	4/4	0.94	0.26	108,110,116,117	0
6	NA	A	1589	1/1	0.94	0.27	27,27,27,27	0
5	FAD	E	601	53/53	0.95	0.33	46,61,71,73	0
11	CBE	G	1129	16/16	0.96	0.29	56,60,62,62	0
11	CBE	C	1129	16/16	0.96	0.20	53,55,56,56	0
12	HEM	K	1130	43/43	0.96	0.29	50,54,72,82	0
7	TEO	E	1590	9/9	0.97	0.16	50,52,53,55	0
12	HEM	G	1130	43/43	0.98	0.27	55,61,71,75	0
12	HEM	C	1130	43/43	0.98	0.25	40,48,62,67	0
9	SF4	J	303	8/8	0.98	0.18	83,87,90,91	0
6	NA	E	1589	1/1	0.98	0.33	45,45,45,45	0
7	TEO	A	1590	9/9	0.98	0.26	50,52,53,53	0
5	FAD	A	601	53/53	0.98	0.24	38,42,53,54	0
9	SF4	B	303	8/8	0.99	0.23	38,40,40,41	0
9	SF4	F	303	8/8	0.99	0.20	55,58,59,59	0
10	F3S	J	304	7/7	0.99	0.13	88,90,92,93	0
8	FES	F	302	4/4	0.99	0.25	54,55,56,58	0
10	F3S	F	304	7/7	0.99	0.13	64,69,72,75	0
8	FES	B	302	4/4	1.00	0.27	38,41,43,44	0
10	F3S	B	304	7/7	1.00	0.16	49,54,58,59	0

6.5 Other polymers ⓘ

There are no such residues in this entry.