



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2018 – 03:23 pm GMT

PDB ID : 4X2P  
Title : P. putida mandelate racemase in complex with 3-hydroxypyruvate  
Authors : Wyatt, B.N.; St.Maurice, M.  
Deposited on : 2014-11-26  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1521 (1.66-1.66)
Clashscore	122126	1616 (1.66-1.66)
Ramachandran outliers	120053	1584 (1.66-1.66)
Sidechain outliers	120020	1584 (1.66-1.66)
RSRZ outliers	108989	1487 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	383	<div> <div>2%</div> <div> <div></div> <div>73%</div> <div>17%</div> <div>• 7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3PY	A	401[A]	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mandelate racemase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2757	1772	469	502	14	0	12	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	initiating methionine	UNP P11444
A	-22	ALA	-	expression tag	UNP P11444
A	-21	SER	-	expression tag	UNP P11444
A	-20	TRP	-	expression tag	UNP P11444
A	-19	SER	-	expression tag	UNP P11444
A	-18	HIS	-	expression tag	UNP P11444
A	-17	PRO	-	expression tag	UNP P11444
A	-16	GLN	-	expression tag	UNP P11444
A	-15	PHE	-	expression tag	UNP P11444
A	-14	GLU	-	expression tag	UNP P11444
A	-13	LYS	-	expression tag	UNP P11444
A	-12	GLY	-	expression tag	UNP P11444
A	-11	ALA	-	expression tag	UNP P11444
A	-10	LEU	-	expression tag	UNP P11444
A	-9	GLU	-	expression tag	UNP P11444
A	-8	VAL	-	expression tag	UNP P11444
A	-7	LEU	-	expression tag	UNP P11444
A	-6	PHE	-	expression tag	UNP P11444
A	-5	GLN	-	expression tag	UNP P11444
A	-4	GLY	-	expression tag	UNP P11444
A	-3	PRO	-	expression tag	UNP P11444
A	-2	GLY	-	expression tag	UNP P11444
A	-1	TYR	-	expression tag	UNP P11444
A	0	HIS	-	expression tag	UNP P11444

- Molecule 2 is 3-HYDROXYPYRUVIC ACID (three-letter code: 3PY) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	1
			12	6	6		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	297	Total	O	0	0
			297	297		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mandelate racemase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.88Å 123.88Å 105.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.65 39.17 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.5 (40.00-1.65) 99.6 (39.17-1.65)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.88 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.143 , 0.177 0.143 , 0.176	Depositor DCC
$R_{free}$ test set	2481 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.2	Xtriage
Anisotropy	0.060	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 52.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3067	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3PY, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.74	28/2847 (1.0%)	1.60	35/3877 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	254[A]	GLU	CD-OE2	9.45	1.36	1.25
1	A	254[B]	GLU	CD-OE2	9.45	1.36	1.25
1	A	234	ARG	CZ-NH1	8.91	1.44	1.33
1	A	349	TRP	CE3-CZ3	8.29	1.52	1.38
1	A	215	GLU	CA-C	-6.98	1.34	1.52
1	A	216	GLY	C-O	6.85	1.34	1.23
1	A	326	ILE	CB-CG1	-6.83	1.34	1.54
1	A	254[A]	GLU	CD-OE1	6.78	1.33	1.25
1	A	254[B]	GLU	CD-OE1	6.78	1.33	1.25
1	A	275	GLY	N-CA	6.38	1.55	1.46
1	A	327	GLU	CG-CD	6.34	1.61	1.51
1	A	346	GLY	N-CA	6.04	1.55	1.46
1	A	174	GLN	CD-NE2	6.00	1.47	1.32
1	A	296	SER	CB-OG	5.99	1.50	1.42
1	A	357	TYR	CB-CG	-5.96	1.42	1.51
1	A	315	TRP	CE3-CZ3	5.76	1.48	1.38
1	A	237	SER	CB-OG	5.67	1.49	1.42
1	A	323	GLY	N-CA	5.55	1.54	1.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	169	TYR	CG-CD1	5.55	1.46	1.39
1	A	84	GLU	CD-OE2	5.38	1.31	1.25
1	A	257	LYS	CE-NZ	-5.35	1.35	1.49
1	A	123	LEU	N-CA	5.34	1.57	1.46
1	A	231	GLY	N-CA	5.31	1.54	1.46
1	A	118	VAL	C-O	5.23	1.33	1.23
1	A	50	TYR	CG-CD1	5.21	1.46	1.39
1	A	82	SER	CA-CB	5.19	1.60	1.52
1	A	150	GLU	CD-OE1	5.19	1.31	1.25
1	A	215	GLU	CB-CG	-5.04	1.42	1.52

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	175	ASP	CB-CG-OD1	10.88	128.10	118.30
1	A	183	ARG	NE-CZ-NH2	-10.04	115.28	120.30
1	A	326	ILE	CB-CG1-CD1	-9.96	86.01	113.90
1	A	125	LYS	CD-CE-NZ	-8.91	91.22	111.70
1	A	146	LYS	CD-CE-NZ	-6.89	95.85	111.70
1	A	52	PHE	CB-CG-CD2	-6.77	116.06	120.80
1	A	68	ASP	CB-CG-OD2	6.75	124.37	118.30
1	A	67	ASP	CB-CG-OD2	-6.61	112.35	118.30
1	A	229	TYR	CB-CG-CD2	-6.53	117.08	121.00
1	A	254[A]	GLU	CG-CD-OE2	-6.40	105.50	118.30
1	A	254[B]	GLU	CG-CD-OE2	-6.40	105.50	118.30
1	A	101	ARG	NE-CZ-NH1	6.37	123.49	120.30
1	A	78[A]	LEU	C-N-CA	6.34	137.56	121.70
1	A	78[B]	LEU	C-N-CA	6.34	137.56	121.70
1	A	257	LYS	CD-CE-NZ	-6.31	97.19	111.70
1	A	229	TYR	CB-CG-CD1	6.27	124.76	121.00
1	A	207	LYS	CD-CE-NZ	-6.27	97.29	111.70
1	A	99	LEU	CB-CG-CD1	-6.25	100.38	111.00
1	A	161	ARG	NE-CZ-NH1	6.23	123.41	120.30
1	A	188	ASP	CB-CG-OD1	6.19	123.87	118.30
1	A	221	GLU	OE1-CD-OE2	6.07	130.58	123.30
1	A	143	ASP	CB-CG-OD2	-6.02	112.88	118.30
1	A	256	PHE	CG-CD2-CE2	-5.97	114.24	120.80
1	A	65	LEU	CB-CG-CD2	-5.91	100.95	111.00
1	A	317	GLU	OE1-CD-OE2	-5.82	116.31	123.30
1	A	216	GLY	N-CA-C	-5.78	98.66	113.10
1	A	319	LEU	CB-CG-CD1	-5.67	101.37	111.00
1	A	3	GLU	N-CA-C	5.59	126.09	111.00

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	TYR	CB-CG-CD2	-5.47	117.72	121.00
1	A	10	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	A	176	LEU	CB-CG-CD1	-5.37	101.88	111.00
1	A	356	LYS	CD-CE-NZ	-5.22	99.68	111.70
1	A	318	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	169	TYR	CG-CD1-CE1	-5.16	117.18	121.30
1	A	29	VAL	CG1-CB-CG2	5.04	118.96	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	3	GLU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2757	0	2839	36	0
2	A	12	0	5	7	0
3	A	1	0	0	0	0
4	A	297	0	0	13	0
All	All	3067	0	2844	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:TYR:CD2	1:A:151[B]:ARG:NH2	1.94	1.32
1:A:20:TYR:CG	1:A:151[B]:ARG:NH2	2.02	1.26
1:A:227[A]:HIS:HD2	4:A:508:HOH:O	1.15	1.24
1:A:20:TYR:CE1	1:A:151[B]:ARG:HD3	1.83	1.13
1:A:227[A]:HIS:CD2	4:A:508:HOH:O	1.93	1.02

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:TYR:CE1	1:A:151[B]:ARG:CD	2.47	0.98
1:A:3:GLU:O	1:A:3:GLU:HG2	1.66	0.93
1:A:20:TYR:CZ	1:A:151[B]:ARG:HD2	2.04	0.92
2:A:401[A]:3PY:H31	4:A:626:HOH:O	1.81	0.79
1:A:166:LYS:HD3	1:A:197[A]:ASN:HD21	1.46	0.79
1:A:166:LYS:CD	1:A:197[A]:ASN:HD21	2.00	0.74
1:A:20:TYR:CZ	1:A:151[B]:ARG:CD	2.71	0.71
1:A:227[A]:HIS:ND1	4:A:502:HOH:O	2.24	0.71
2:A:401[B]:3PY:H31	4:A:626:HOH:O	1.91	0.69
2:A:401[A]:3PY:C2	4:A:750:HOH:O	2.41	0.68
1:A:166:LYS:HD2	1:A:197[A]:ASN:ND2	2.11	0.66
1:A:124:VAL:H	1:A:305:HIS:HD2	1.43	0.65
1:A:166:LYS:CD	1:A:197[A]:ASN:ND2	2.62	0.62
1:A:135:GLN:HA	1:A:336:ASN:HD22	1.63	0.62
1:A:60:LYS:HE3	1:A:359:VAL:CG1	2.32	0.59
1:A:20:TYR:CD1	1:A:151[B]:ARG:NH2	2.68	0.59
1:A:20:TYR:CD1	1:A:151[B]:ARG:HD3	2.37	0.58
1:A:60:LYS:HE3	1:A:359:VAL:HG11	1.87	0.57
1:A:112:TRP:CD1	1:A:277:VAL:HB	2.42	0.54
1:A:141:SER:HA	1:A:166:LYS:HG3	1.91	0.53
1:A:122:PRO:HD2	1:A:125:LYS:HD2	1.88	0.53
1:A:150:GLU:OE2	4:A:791:HOH:O	2.19	0.53
2:A:401[A]:3PY:C1	4:A:750:HOH:O	2.58	0.51
1:A:124:VAL:H	1:A:305:HIS:CD2	2.27	0.49
1:A:121[B]:THR:HG22	1:A:125:LYS:HB2	1.95	0.48
2:A:401[A]:3PY:C3	4:A:626:HOH:O	2.51	0.48
1:A:20:TYR:CB	1:A:151[B]:ARG:NH2	2.72	0.48
1:A:121[B]:THR:HG22	1:A:122:PRO:HD2	1.98	0.46
1:A:16:VAL:HB	1:A:324[A]:SER:OG	2.16	0.45
1:A:317:GLU:OE2	2:A:401[B]:3PY:O1	2.35	0.45
1:A:6:ILE:HD11	1:A:83[B]:LEU:HD11	2.00	0.44
1:A:166:LYS:HE3	4:A:752:HOH:O	2.16	0.44
1:A:298:LEU:HD21	1:A:319:LEU:HD13	1.99	0.44
1:A:20:TYR:CD1	1:A:151[B]:ARG:CD	2.98	0.44
2:A:401[B]:3PY:H32	4:A:752:HOH:O	2.17	0.42
1:A:254[B]:GLU:OE2	4:A:501:HOH:O	2.21	0.42
1:A:180:ARG:HD2	4:A:782:HOH:O	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/383 (96%)	356 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/301 (97%)	284 (98%)	7 (2%)	52	25

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	10	ARG
1	A	20	TYR
1	A	21	PRO
1	A	55	THR
1	A	226	GLN
1	A	326	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	226	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	305	HIS
1	A	336	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	3PY	A	401[A]	1,3	2,5,6	0.97	0	1,5,7	4.65	1 (100%)
2	3PY	A	401[B]	1,3	2,5,6	2.74	1 (50%)	1,5,7	3.27	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3PY	A	401[A]	1,3	-	0/1/3/6	0/0/0/0
2	3PY	A	401[B]	1,3	-	0/1/3/6	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401[B]	3PY	C2-C3	-3.66	1.31	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401[B]	3PY	O4-C3-C2	3.27	125.33	111.03
2	A	401[A]	3PY	O4-C3-C2	4.65	131.38	111.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401[A]	3PY	4	0
2	A	401[B]	3PY	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	357/383 (93%)	-0.48	6 (1%) 70 74	8, 14, 29, 63	3 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	26	VAL	5.5
1	A	25	ALA	4.0
1	A	359	VAL	3.7
1	A	27	GLY	2.6
1	A	28	THR	2.2
1	A	189	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	3PY	A	401[A]	6/7	0.86	0.16	14,26,31,32	6

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	3PY	A	401[B]	6/7	0.86	0.16	19,28,31,32	6
3	MG	A	402	1/1	1.00	0.16	2,2,2,2	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.