



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 08:24 pm GMT

PDB ID : 4Y1K
Title : PALMITOYLATED OPRM OUTER MEMBRANE FACTOR
Authors : Monlezun, L.; Phan, G.; Broutin, I.
Deposited on : 2015-02-07
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1028 (4.02-3.58)
Clashscore	122126	1061 (4.00-3.60)
Ramachandran outliers	120053	1025 (4.00-3.60)
Sidechain outliers	120020	1019 (4.00-3.60)
RSRZ outliers	108989	1021 (4.06-3.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	474	<div> <div>12%</div> <div>46%</div> <div>43%</div> <div>7%</div> <div>.</div> </div>
1	B	474	<div> <div>10%</div> <div>47%</div> <div>45%</div> <div>.</div> <div>.</div> </div>
1	C	474	<div> <div>12%</div> <div>48%</div> <div>42%</div> <div>6%</div> <div>.</div> </div>
1	D	474	<div> <div>27%</div> <div>44%</div> <div>45%</div> <div>7%</div> <div>.</div> </div>
1	E	474	<div> <div>26%</div> <div>46%</div> <div>46%</div> <div>.</div> <div>.</div> </div>
1	F	474	<div> <div>24%</div> <div>41%</div> <div>50%</div> <div>5%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLM	A	1001	-	-	-	X

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 21050 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer membrane protein OprM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	457	Total	C	N	O	S	0	0	0
			3501	2190	624	684	3			
1	B	457	Total	C	N	O	S	0	0	0
			3501	2190	624	684	3			
1	C	457	Total	C	N	O	S	0	0	0
			3503	2191	624	685	3			
1	D	457	Total	C	N	O	S	0	0	0
			3501	2190	624	684	3			
1	E	456	Total	C	N	O	S	0	0	0
			3492	2185	622	682	3			
1	F	457	Total	C	N	O	S	0	0	0
			3501	2190	624	684	3			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	469	HIS	-	expression tag	UNP Q51487
A	470	HIS	-	expression tag	UNP Q51487
A	471	HIS	-	expression tag	UNP Q51487
A	472	HIS	-	expression tag	UNP Q51487
A	473	HIS	-	expression tag	UNP Q51487
A	474	HIS	-	expression tag	UNP Q51487
B	469	HIS	-	expression tag	UNP Q51487
B	470	HIS	-	expression tag	UNP Q51487
B	471	HIS	-	expression tag	UNP Q51487
B	472	HIS	-	expression tag	UNP Q51487
B	473	HIS	-	expression tag	UNP Q51487
B	474	HIS	-	expression tag	UNP Q51487
C	469	HIS	-	expression tag	UNP Q51487
C	470	HIS	-	expression tag	UNP Q51487
C	471	HIS	-	expression tag	UNP Q51487
C	472	HIS	-	expression tag	UNP Q51487
C	473	HIS	-	expression tag	UNP Q51487

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	474	HIS	-	expression tag	UNP Q51487
D	469	HIS	-	expression tag	UNP Q51487
D	470	HIS	-	expression tag	UNP Q51487
D	471	HIS	-	expression tag	UNP Q51487
D	472	HIS	-	expression tag	UNP Q51487
D	473	HIS	-	expression tag	UNP Q51487
D	474	HIS	-	expression tag	UNP Q51487
E	469	HIS	-	expression tag	UNP Q51487
E	470	HIS	-	expression tag	UNP Q51487
E	471	HIS	-	expression tag	UNP Q51487
E	472	HIS	-	expression tag	UNP Q51487
E	473	HIS	-	expression tag	UNP Q51487
E	474	HIS	-	expression tag	UNP Q51487
F	469	HIS	-	expression tag	UNP Q51487
F	470	HIS	-	expression tag	UNP Q51487
F	471	HIS	-	expression tag	UNP Q51487
F	472	HIS	-	expression tag	UNP Q51487
F	473	HIS	-	expression tag	UNP Q51487
F	474	HIS	-	expression tag	UNP Q51487

- Molecule 2 is PALMITIC ACID (three-letter code: PLM) (formula: C₁₆H₃₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			17	16	1		
2	B	1	Total	C	O	0	0
			17	16	1		

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	C	1	17	16	1	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Outer membrane protein OprM






• Molecule 1: Outer membrane protein OprM



• Molecule 1: Outer membrane protein OprM





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	152.64Å 87.86Å 355.94Å 90.00° 98.94° 90.00°	Depositor
Resolution (Å)	87.91 – 3.80 87.90 – 3.80	Depositor EDS
% Data completeness (in resolution range)	86.8 (87.91-3.80) 84.4 (87.90-3.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 3.78Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.297 , 0.346 0.296 , 0.347	Depositor DCC
R_{free} test set	1992 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	96.7	Xtriage
Anisotropy	0.934	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 86.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.378 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-l$ 0.339 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-l$	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	21050	wwPDB-VP
Average B, all atoms (Å ²)	94.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/3557	0.73	0/4838
1	B	0.58	0/3557	0.72	0/4838
1	C	0.56	0/3559	0.72	0/4841
1	D	0.43	0/3557	0.63	1/4838 (0.0%)
1	E	0.44	0/3548	0.61	0/4826
1	F	0.43	0/3557	0.63	0/4838
All	All	0.50	0/21335	0.67	1/29019 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	442	LEU	CA-CB-CG	-5.06	103.67	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3501	0	3475	188	0
1	B	3501	0	3475	173	0
1	C	3503	0	3480	173	0
1	D	3501	0	3476	191	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	3492	0	3468	187	0
1	F	3501	0	3476	210	0
2	A	17	0	31	0	0
2	B	17	0	31	0	0
2	C	17	0	31	1	0
All	All	21050	0	20943	1038	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 1038 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:177:LEU:HD21	1:D:425:GLN:HB3	1.48	0.95
1:D:372:GLY:HA3	1:D:442:LEU:HD22	1.55	0.88
1:E:210:GLN:HE21	1:E:214:GLU:HG3	1.39	0.87
1:A:81:ARG:HA	1:A:136:ARG:HG3	1.57	0.87
1:E:98:ARG:HA	1:E:111:ILE:HG12	1.55	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/474 (96%)	363 (80%)	74 (16%)	18 (4%)	3	32
1	B	455/474 (96%)	368 (81%)	68 (15%)	19 (4%)	3	31
1	C	455/474 (96%)	365 (80%)	67 (15%)	23 (5%)	2	27
1	D	455/474 (96%)	371 (82%)	60 (13%)	24 (5%)	2	27
1	E	454/474 (96%)	380 (84%)	51 (11%)	23 (5%)	2	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	455/474 (96%)	385 (85%)	54 (12%)	16 (4%)	4	36
All	All	2729/2844 (96%)	2232 (82%)	374 (14%)	123 (4%)	3	29

5 of 123 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	23	ALA
1	A	69	ASN
1	A	314	SER
1	B	22	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/375 (96%)	320 (89%)	39 (11%)	7	34
1	B	359/375 (96%)	319 (89%)	40 (11%)	7	33
1	C	360/375 (96%)	321 (89%)	39 (11%)	7	34
1	D	359/375 (96%)	322 (90%)	37 (10%)	8	36
1	E	358/375 (96%)	333 (93%)	25 (7%)	16	53
1	F	359/375 (96%)	325 (90%)	34 (10%)	9	39
All	All	2154/2250 (96%)	1940 (90%)	214 (10%)	8	37

5 of 214 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	240	SER
1	D	58	GLU
1	F	256	LEU
1	C	256	LEU
1	C	344	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	210	GLN
1	B	451	ASN
1	E	20	GLN
1	B	193	GLN
1	D	357	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLM	A	1001	1	16,16,17	1.58	3 (18%)	14,15,17	1.12	2 (14%)
2	PLM	B	1001	1	16,16,17	0.94	1 (6%)	14,15,17	1.04	1 (7%)
2	PLM	C	1001	1	16,16,17	1.06	1 (6%)	14,15,17	1.13	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLM	A	1001	1	-	0/13/14/15	0/0/0/0
2	PLM	B	1001	1	-	0/13/14/15	0/0/0/0
2	PLM	C	1001	1	-	0/13/14/15	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1001	PLM	C2-C1	-4.04	1.38	1.49
2	B	1001	PLM	C2-C1	-3.24	1.40	1.49
2	A	1001	PLM	C5-C4	-2.36	1.38	1.51
2	A	1001	PLM	C7-C6	2.08	1.63	1.51
2	A	1001	PLM	C3-C2	4.87	1.71	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	PLM	O2-C1-C2	-2.86	111.76	125.63
2	C	1001	PLM	O2-C1-C2	-2.59	113.09	125.63
2	B	1001	PLM	O2-C1-C2	-2.44	113.78	125.63
2	A	1001	PLM	C4-C3-C2	2.20	122.11	113.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1001	PLM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	457/474 (96%)	0.83	55 (12%) 4 4	57, 72, 105, 105	0
1	B	457/474 (96%)	0.81	48 (10%) 6 5	54, 69, 102, 102	0
1	C	457/474 (96%)	0.86	57 (12%) 4 4	54, 69, 102, 102	0
1	D	457/474 (96%)	1.31	129 (28%) 0 0	81, 121, 121, 121	0
1	E	456/474 (96%)	1.44	124 (27%) 0 0	109, 120, 120, 120	0
1	F	457/474 (96%)	1.30	113 (24%) 0 0	79, 119, 119, 119	0
All	All	2741/2844 (96%)	1.09	526 (19%) 1 1	54, 105, 121, 121	0

The worst 5 of 526 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	203	ALA	18.0
1	F	210	GLN	16.4
1	E	219	THR	13.5
1	D	261	PRO	10.6
1	A	202	SER	10.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PLM	A	1001	17/18	0.65	0.44	72,72,76,76	0
2	PLM	B	1001	17/18	0.77	0.37	72,72,72,72	0
2	PLM	C	1001	17/18	0.80	0.30	58,58,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.