



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 28, 2019 – 12:24 PM EDT

PDB ID : 5Y9J
Title : BAFF in complex with belimumab
Authors : Heo, Y.-S.; Shin, W.
Deposited on : 2017-08-25
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.4
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1449 (2.04-2.04)
Clashscore	122126	1524 (2.04-2.04)
Ramachandran outliers	120053	1512 (2.04-2.04)
Sidechain outliers	120020	1512 (2.04-2.04)
RSRZ outliers	108989	1429 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	<div> <div>2%</div> <div>93%</div> <div>5%</div> <div>•</div> </div>
2	H	235	<div> <div>%</div> <div>89%</div> <div>7%</div> <div>•</div> </div>
3	A	152	<div> <div>3%</div> <div>90%</div> <div>5%</div> <div>5%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called belibumab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	210	Total	C	N	O	S	0	0	0
			1573	978	267	324	4			

- Molecule 2 is a protein called belimumab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	218	Total	C	N	O	S	0	0	0
			1614	1013	280	313	8			

- Molecule 3 is a protein called Tumor necrosis factor ligand superfamily member 13B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	144	Total	C	N	O	S	0	0	0
			1143	736	184	218	5			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	177	Total	O	0	0
			177	177		
4	H	187	Total	O	0	0
			187	187		
4	A	136	Total	O	0	0
			136	136		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: belibumab light chain



- Molecule 2: belimumab heavy chain



- Molecule 3: Tumor necrosis factor ligand superfamily member 13B



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	134.00Å 134.00Å 134.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.28 – 2.05 29.96 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.7 (26.28-2.05) 96.6 (29.96-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.76 (at 2.05Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.180 , 0.217 0.181 , 0.218	Depositor DCC
R_{free} test set	2458 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	30.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4830	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.34	0/1611	0.49	0/2201
2	H	0.37	0/1651	0.57	2/2246 (0.1%)
3	A	0.44	0/1165	0.57	0/1574
All	All	0.38	0/4427	0.54	2/6021 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	111	SER	C-N-CD	-7.37	104.39	120.60
2	H	112	PRO	N-CA-C	-5.84	96.91	112.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	111	SER	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1573	0	1504	11	0
2	H	1614	0	1592	5	0
3	A	1143	0	1147	4	0
4	A	136	0	0	2	2
4	H	187	0	0	2	0
4	L	177	0	0	8	0
All	All	4830	0	4243	20	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:260:GLN:OE1	4:A:301:HOH:O	1.86	0.93
1:L:158:LYS:NZ	4:L:302:HOH:O	2.23	0.70
2:H:2:VAL:N	4:H:303:HOH:O	2.29	0.65
2:H:98:ARG:HB3	2:H:112:PRO:HD2	1.80	0.64
2:H:171:SER:OG	4:H:301:HOH:O	2.15	0.63
1:L:159:ALA:N	4:L:307:HOH:O	2.33	0.61
1:L:9:ALA:N	4:L:305:HOH:O	2.28	0.59
2:H:202:GLN:OE1	2:H:203:THR:N	2.36	0.58
1:L:13:ALA:HB3	1:L:16:GLN:HG3	1.85	0.57
1:L:134:LEU:HD12	1:L:180:LEU:HD23	1.86	0.57
3:A:142:VAL:HG12	3:A:143:THR:HG23	1.88	0.56
1:L:38:LYS:NZ	4:L:309:HOH:O	2.38	0.56
1:L:167:SER:OG	4:L:301:HOH:O	2.14	0.56
2:H:33:ALA:HB2	2:H:52:ILE:HG12	1.90	0.53
3:A:216:LYS:HE3	3:A:219:VAL:HG12	1.92	0.51
1:L:9:ALA:C	4:L:305:HOH:O	2.52	0.48
3:A:186:LEU:HB3	4:A:407:HOH:O	2.14	0.46
1:L:158:LYS:HB3	4:L:307:HOH:O	2.18	0.43
1:L:19:ARG:HD3	4:L:410:HOH:O	2.19	0.43
1:L:155:SER:HA	1:L:156:PRO:HD3	1.89	0.42

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:382:HOH:O	4:A:414:HOH:O[10_545]	1.86	0.34
4:A:401:HOH:O	4:A:401:HOH:O[10_545]	1.87	0.33

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	208/214 (97%)	203 (98%)	5 (2%)	0	100	100
2	H	214/235 (91%)	207 (97%)	6 (3%)	1 (0%)	31	20
3	A	142/152 (93%)	141 (99%)	1 (1%)	0	100	100
All	All	564/601 (94%)	551 (98%)	12 (2%)	1 (0%)	49	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	112	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	175/179 (98%)	174 (99%)	1 (1%)	87	88
2	H	180/197 (91%)	179 (99%)	1 (1%)	87	88
3	A	126/132 (96%)	125 (99%)	1 (1%)	83	84
All	All	481/508 (95%)	478 (99%)	3 (1%)	87	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	158	LYS
2	H	170	THR
3	A	217	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	210/214 (98%)	-0.31	4 (1%) 66 71	24, 37, 65, 88	0
2	H	218/235 (92%)	-0.30	3 (1%) 75 78	24, 35, 62, 86	0
3	A	144/152 (94%)	-0.26	4 (2%) 53 59	18, 27, 58, 80	0
All	All	572/601 (95%)	-0.29	11 (1%) 66 71	18, 34, 62, 88	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	220	PHE	5.2
1	L	158	LYS	4.5
3	A	142	VAL	3.7
1	L	211	THR	3.4
1	L	2	SER	3.0
2	H	200	GLY	2.7
3	A	218	HIS	2.6
2	H	75	SER	2.5
2	H	25	SER	2.1
1	L	8	PRO	2.1
3	A	285	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.