



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 13, 2018 – 03:19 pm GMT

PDB ID : 2YF2  
Title : Crystal structure of the oligomerisation domain of C4b-binding protein from Gallus gallus  
Authors : Caesar, J.J.E.; Hill, F.; Lea, S.M.  
Deposited on : 2011-04-01  
Resolution : 2.24 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2027 (2.26-2.22)
Clashscore	122126	2170 (2.26-2.22)
Ramachandran outliers	120053	2129 (2.26-2.22)
Sidechain outliers	120020	2130 (2.26-2.22)
RSRZ outliers	108989	1991 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	65	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>5%</div> <div>12%</div> </div> </div>
1	B	65	<div> <div>3%</div> <div> <div></div> <div>65%</div> <div>6%</div> <div>29%</div> </div> </div>
1	C	65	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>11%</div> </div> </div>
1	D	65	<div> <div>3%</div> <div> <div></div> <div>66%</div> <div>5%</div> <div>29%</div> </div> </div>
1	E	65	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>•</div> <div>15%</div> </div> </div>
1	F	65	<div> <div>2%</div> <div> <div></div> <div>65%</div> <div>5%</div> <div>31%</div> </div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	65	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	1459	-	-	-	X
2	EDO	B	1452	-	-	X	-
2	EDO	D	1448	-	-	-	X
2	EDO	D	1449	-	-	-	X
2	EDO	F	1448	-	-	X	X
3	ACT	A	1460	-	-	X	-
3	ACT	B	1453	-	-	X	-
3	ACT	B	1454	-	-	X	-
3	ACT	D	1450	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C4B BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	57	Total	C	N	O	S	0	0	0
			451	291	72	86	2			
1	B	46	Total	C	N	O	S	0	0	0
			364	232	58	72	2			
1	C	58	Total	C	N	O	S	0	0	0
			460	296	75	87	2			
1	D	46	Total	C	N	O	S	0	0	0
			360	230	58	70	2			
1	E	55	Total	C	N	O	S	0	0	0
			437	283	70	82	2			
1	F	45	Total	C	N	O	S	0	0	0
			355	227	57	69	2			
1	G	57	Total	C	N	O	S	0	0	0
			455	293	74	86	2			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	396	SER	CYS	engineered mutation	UNP E1C2T5
B	396	SER	CYS	engineered mutation	UNP E1C2T5
C	396	SER	CYS	engineered mutation	UNP E1C2T5
D	396	SER	CYS	engineered mutation	UNP E1C2T5
E	396	SER	CYS	engineered mutation	UNP E1C2T5
F	396	SER	CYS	engineered mutation	UNP E1C2T5
G	396	SER	CYS	engineered mutation	UNP E1C2T5
A	459	GLY	-	expression tag	UNP E1C2T5
A	460	SER	-	expression tag	UNP E1C2T5
B	459	GLY	-	expression tag	UNP E1C2T5
B	460	SER	-	expression tag	UNP E1C2T5
C	459	GLY	-	expression tag	UNP E1C2T5
C	460	SER	-	expression tag	UNP E1C2T5
D	459	GLY	-	expression tag	UNP E1C2T5
D	460	SER	-	expression tag	UNP E1C2T5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	459	GLY	-	expression tag	UNP E1C2T5
E	460	SER	-	expression tag	UNP E1C2T5
F	459	GLY	-	expression tag	UNP E1C2T5
F	460	SER	-	expression tag	UNP E1C2T5
G	459	GLY	-	expression tag	UNP E1C2T5
G	460	SER	-	expression tag	UNP E1C2T5

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	F	1	Total C O 4 2 2	0	0
2	F	1	Total C O 4 2 2	0	0
2	F	1	Total C O 4 2 2	0	0
2	G	1	Total C O 4 2 2	0	0
2	G	1	Total C O 4 2 2	0	0
2	G	1	Total C O 4 2 2	0	0
2	G	1	Total C O 4 2 2	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $\text{C}_2\text{H}_3\text{O}_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	E	1	Total C O 4 2 2	0	0
3	E	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0
4	B	12	Total O 12 12	0	0
4	C	13	Total O 13 13	0	0
4	D	11	Total O 11 11	0	0
4	E	10	Total O 10 10	0	0
4	F	12	Total O 12 12	0	0
4	G	9	Total O 9 9	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: C4B BINDING PROTEIN



#### • Molecule 1: C4B BINDING PROTEIN





#### • Molecule 1: C4B BINDING PROTEIN



#### • Molecule 1: C4B BINDING PROTEIN




#### • Molecule 1: C4B BINDING PROTEIN





#### • Molecule 1: C4B BINDING PROTEIN







● Molecule 1: C4B BINDING PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.89Å 140.24Å 85.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.06 – 2.24 55.06 – 2.24	Depositor EDS
% Data completeness (in resolution range)	95.6 (55.06-2.24) 95.6 (55.06-2.24)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 2.25Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, $R_{free}$	0.220 , 0.236 0.228 , 0.244	Depositor DCC
$R_{free}$ test set	1200 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.8	Xtriage
Anisotropy	0.845	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.001 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.000 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3073	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/456	0.53	0/615
1	B	0.44	0/367	0.55	0/496
1	C	0.46	0/466	0.57	0/628
1	D	0.44	0/363	0.54	0/490
1	E	0.44	0/442	0.53	0/595
1	F	0.43	0/358	0.54	0/483
1	G	0.43	0/461	0.55	0/621
All	All	0.44	0/2913	0.55	0/3928

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	451	0	464	3	0
1	B	364	0	373	9	0
1	C	460	0	472	5	0
1	D	360	0	372	3	0
1	E	437	0	454	2	0
1	F	355	0	367	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	455	0	467	1	0
2	A	4	0	6	0	0
2	B	24	0	36	10	0
2	C	12	0	18	5	0
2	D	8	0	12	3	0
2	F	12	0	18	5	0
2	G	16	0	24	1	0
3	A	4	0	3	2	0
3	B	12	0	9	9	0
3	C	4	0	3	0	0
3	D	8	0	6	2	0
3	E	8	0	6	1	0
3	F	4	0	3	1	0
3	G	4	0	3	0	0
4	A	4	0	0	0	0
4	B	12	0	0	0	0
4	C	13	0	0	0	0
4	D	11	0	0	1	0
4	E	10	0	0	0	0
4	F	12	0	0	0	0
4	G	9	0	0	0	0
All	All	3073	0	3116	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:422:THR:HB	2:B:1452:EDO:C2	2.06	0.84
1:C:433:LYS:HE2	2:D:1448:EDO:H21	1.59	0.84
1:B:422:THR:HB	2:B:1452:EDO:H21	1.59	0.82
1:B:422:THR:OG1	3:B:1454:ACT:H2	1.80	0.79
1:G:441:LEU:HG	2:G:1462:EDO:H12	1.65	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	55/65 (85%)	55 (100%)	0	0	100	100
1	B	44/65 (68%)	44 (100%)	0	0	100	100
1	C	56/65 (86%)	55 (98%)	1 (2%)	0	100	100
1	D	44/65 (68%)	43 (98%)	1 (2%)	0	100	100
1	E	53/65 (82%)	53 (100%)	0	0	100	100
1	F	43/65 (66%)	43 (100%)	0	0	100	100
1	G	55/65 (85%)	54 (98%)	1 (2%)	0	100	100
All	All	350/455 (77%)	347 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	52/59 (88%)	52 (100%)	0	100	100
1	B	43/59 (73%)	43 (100%)	0	100	100
1	C	53/59 (90%)	53 (100%)	0	100	100
1	D	42/59 (71%)	42 (100%)	0	100	100
1	E	51/59 (86%)	51 (100%)	0	100	100
1	F	42/59 (71%)	42 (100%)	0	100	100
1	G	53/59 (90%)	53 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	336/413 (81%)	336 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	412	GLN
1	C	412	GLN
1	E	412	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	1459	-	3,3,3	0.49	0	2,2,2	0.18	0
3	ACT	A	1460	-	1,3,3	1.40	0	0,3,3	0.00	-
2	EDO	B	1447	-	3,3,3	0.36	0	2,2,2	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EDO	B	1448	-	3,3,3	0.51	0	2,2,2	0.23	0
2	EDO	B	1449	-	3,3,3	0.48	0	2,2,2	0.25	0
2	EDO	B	1450	-	3,3,3	0.49	0	2,2,2	0.34	0
2	EDO	B	1451	-	3,3,3	0.38	0	2,2,2	0.46	0
2	EDO	B	1452	-	3,3,3	0.30	0	2,2,2	0.09	0
3	ACT	B	1453	-	1,3,3	0.73	0	0,3,3	0.00	-
3	ACT	B	1454	-	1,3,3	4.42	1 (100%)	0,3,3	0.00	-
3	ACT	B	1455	-	1,3,3	0.86	0	0,3,3	0.00	-
2	EDO	C	1460	-	3,3,3	0.42	0	2,2,2	0.34	0
2	EDO	C	1461	-	3,3,3	0.35	0	2,2,2	0.51	0
2	EDO	C	1462	-	3,3,3	0.37	0	2,2,2	0.39	0
3	ACT	C	1463	-	1,3,3	0.94	0	0,3,3	0.00	-
2	EDO	D	1448	-	3,3,3	0.50	0	2,2,2	0.16	0
2	EDO	D	1449	-	3,3,3	0.50	0	2,2,2	0.31	0
3	ACT	D	1450	-	1,3,3	0.21	0	0,3,3	0.00	-
3	ACT	D	1451	-	1,3,3	1.91	0	0,3,3	0.00	-
3	ACT	E	1459	-	1,3,3	2.92	1 (100%)	0,3,3	0.00	-
3	ACT	E	1460	-	1,3,3	2.10	1 (100%)	0,3,3	0.00	-
2	EDO	F	1448	-	3,3,3	0.46	0	2,2,2	0.25	0
2	EDO	F	1449	-	3,3,3	0.45	0	2,2,2	0.34	0
2	EDO	F	1450	-	3,3,3	0.51	0	2,2,2	0.30	0
3	ACT	F	1451	-	1,3,3	1.34	0	0,3,3	0.00	-
2	EDO	G	1460	-	3,3,3	0.52	0	2,2,2	0.27	0
2	EDO	G	1461	-	3,3,3	0.52	0	2,2,2	0.28	0
2	EDO	G	1462	-	3,3,3	0.42	0	2,2,2	0.40	0
2	EDO	G	1463	-	3,3,3	0.48	0	2,2,2	0.30	0
3	ACT	G	1464	-	1,3,3	0.69	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1459	-	-	0/1/1/1	0/0/0/0
3	ACT	A	1460	-	-	0/0/0/0	0/0/0/0
2	EDO	B	1447	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1448	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1449	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1450	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1451	-	-	0/1/1/1	0/0/0/0
2	EDO	B	1452	-	-	0/1/1/1	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	B	1453	-	-	0/0/0/0	0/0/0/0
3	ACT	B	1454	-	-	0/0/0/0	0/0/0/0
3	ACT	B	1455	-	-	0/0/0/0	0/0/0/0
2	EDO	C	1460	-	-	0/1/1/1	0/0/0/0
2	EDO	C	1461	-	-	0/1/1/1	0/0/0/0
2	EDO	C	1462	-	-	0/1/1/1	0/0/0/0
3	ACT	C	1463	-	-	0/0/0/0	0/0/0/0
2	EDO	D	1448	-	-	0/1/1/1	0/0/0/0
2	EDO	D	1449	-	-	0/1/1/1	0/0/0/0
3	ACT	D	1450	-	-	0/0/0/0	0/0/0/0
3	ACT	D	1451	-	-	0/0/0/0	0/0/0/0
3	ACT	E	1459	-	-	0/0/0/0	0/0/0/0
3	ACT	E	1460	-	-	0/0/0/0	0/0/0/0
2	EDO	F	1448	-	-	0/1/1/1	0/0/0/0
2	EDO	F	1449	-	-	0/1/1/1	0/0/0/0
2	EDO	F	1450	-	-	0/1/1/1	0/0/0/0
3	ACT	F	1451	-	-	0/0/0/0	0/0/0/0
2	EDO	G	1460	-	-	0/1/1/1	0/0/0/0
2	EDO	G	1461	-	-	0/1/1/1	0/0/0/0
2	EDO	G	1462	-	-	0/1/1/1	0/0/0/0
2	EDO	G	1463	-	-	0/1/1/1	0/0/0/0
3	ACT	G	1464	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1454	ACT	CH3-C	-4.42	1.43	1.48
3	E	1460	ACT	CH3-C	2.10	1.51	1.48
3	E	1459	ACT	CH3-C	2.92	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

16 monomers are involved in 33 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1460	ACT	2	0
2	B	1447	EDO	1	0
2	B	1449	EDO	3	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1452	EDO	6	0
3	B	1453	ACT	4	0
3	B	1454	ACT	5	0
2	C	1460	EDO	2	0
2	C	1461	EDO	2	0
2	C	1462	EDO	1	0
2	D	1448	EDO	2	0
2	D	1449	EDO	2	0
3	D	1450	ACT	2	0
3	E	1460	ACT	1	0
2	F	1448	EDO	5	0
3	F	1451	ACT	1	0
2	G	1462	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	57/65 (87%)	0.12	1 (1%) 68 70	34, 48, 72, 88	0
1	B	46/65 (70%)	0.46	2 (4%) 35 35	31, 46, 83, 108	0
1	C	58/65 (89%)	0.29	1 (1%) 70 71	31, 49, 73, 100	0
1	D	46/65 (70%)	0.23	2 (4%) 35 35	29, 46, 78, 90	0
1	E	55/65 (84%)	0.10	2 (3%) 42 42	31, 55, 81, 102	0
1	F	45/65 (69%)	0.09	1 (2%) 62 64	34, 52, 76, 95	0
1	G	57/65 (87%)	0.28	2 (3%) 44 44	35, 55, 90, 122	0
All	All	364/455 (80%)	0.22	11 (3%) 50 51	29, 50, 83, 122	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	404	VAL	6.1
1	G	456	ILE	5.0
1	B	401	ASP	3.9
1	E	456	ILE	3.3
1	F	446	GLN	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	F	1448	4/4	0.39	0.44	86,88,89,90	0
2	EDO	D	1448	4/4	0.61	0.46	78,79,79,83	0
2	EDO	F	1450	4/4	0.64	0.31	84,85,86,90	0
3	ACT	B	1455	4/4	0.64	0.26	98,100,100,100	0
2	EDO	G	1461	4/4	0.64	0.39	79,81,83,83	0
2	EDO	A	1459	4/4	0.65	0.53	74,76,78,79	0
2	EDO	B	1448	4/4	0.74	0.30	73,75,75,76	0
2	EDO	C	1460	4/4	0.74	0.21	60,60,60,68	0
2	EDO	B	1449	4/4	0.75	0.18	73,75,77,77	0
2	EDO	C	1461	4/4	0.77	0.36	66,67,67,68	0
2	EDO	G	1463	4/4	0.78	0.30	51,53,55,62	0
3	ACT	D	1450	4/4	0.78	0.23	83,83,84,84	0
2	EDO	D	1449	4/4	0.80	0.44	68,68,69,75	0
2	EDO	C	1462	4/4	0.80	0.31	52,53,54,56	0
2	EDO	B	1450	4/4	0.83	0.26	55,56,57,58	0
2	EDO	B	1447	4/4	0.84	0.26	54,57,58,60	0
2	EDO	F	1449	4/4	0.84	0.35	59,61,62,64	0
3	ACT	C	1463	4/4	0.85	0.21	59,59,60,60	0
2	EDO	G	1462	4/4	0.85	0.55	70,72,73,73	0
3	ACT	D	1451	4/4	0.86	0.29	64,64,64,64	0
2	EDO	G	1460	4/4	0.87	0.16	61,64,67,70	0
3	ACT	B	1453	4/4	0.87	0.19	62,63,63,63	0
2	EDO	B	1451	4/4	0.88	0.28	55,58,62,62	0
2	EDO	B	1452	4/4	0.89	0.40	47,47,47,49	0
3	ACT	E	1460	4/4	0.89	0.19	65,66,66,66	0
3	ACT	G	1464	4/4	0.91	0.13	50,50,51,51	0
3	ACT	B	1454	4/4	0.92	0.29	58,59,60,60	0
3	ACT	E	1459	4/4	0.93	0.09	51,51,51,52	0
3	ACT	A	1460	4/4	0.94	0.26	80,80,80,80	0
3	ACT	F	1451	4/4	0.94	0.32	64,64,65,65	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.