



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2018 – 11:26 am GMT

PDB ID : 4ZCC
Title : Renalase in complex with NADH
Authors : Silvaggi, N.R.; Moran, G.R.; Roman, J.V.
Deposited on : 2015-04-15
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	7193 (2.00-2.00)
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)
RSRZ outliers	108989	7011 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>6%</div> <div>5%</div> </div> </div>
1	B	336	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>7%</div> <div>5%</div> </div> </div>
1	C	336	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>7%</div> <div>• •</div> </div> </div>
1	D	336	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>11%</div> <div>•</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20810 atoms, of which 9759 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Renalase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	320	Total	C	H	N	O	S	0	0	0
			4828	1552	2365	449	450	12			
1	B	319	Total	C	H	N	O	S	0	0	0
			4811	1547	2357	447	448	12			
1	C	321	Total	C	H	N	O	S	0	2	0
			4861	1562	2383	451	453	12			
1	D	328	Total	C	H	N	O	S	0	0	0
			4938	1584	2418	460	464	12			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	145	SER	GLY	engineered mutation	UNP Q48MT7
A	329	LEU	-	expression tag	UNP Q48MT7
A	330	GLU	-	expression tag	UNP Q48MT7
A	331	HIS	-	expression tag	UNP Q48MT7
A	332	HIS	-	expression tag	UNP Q48MT7
A	333	HIS	-	expression tag	UNP Q48MT7
A	334	HIS	-	expression tag	UNP Q48MT7
A	335	HIS	-	expression tag	UNP Q48MT7
A	336	HIS	-	expression tag	UNP Q48MT7
B	145	SER	GLY	engineered mutation	UNP Q48MT7
B	329	LEU	-	expression tag	UNP Q48MT7
B	330	GLU	-	expression tag	UNP Q48MT7
B	331	HIS	-	expression tag	UNP Q48MT7
B	332	HIS	-	expression tag	UNP Q48MT7
B	333	HIS	-	expression tag	UNP Q48MT7
B	334	HIS	-	expression tag	UNP Q48MT7
B	335	HIS	-	expression tag	UNP Q48MT7
B	336	HIS	-	expression tag	UNP Q48MT7
C	145	SER	GLY	engineered mutation	UNP Q48MT7
C	329	LEU	-	expression tag	UNP Q48MT7
C	330	GLU	-	expression tag	UNP Q48MT7

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	331	HIS	-	expression tag	UNP Q48MT7
C	332	HIS	-	expression tag	UNP Q48MT7
C	333	HIS	-	expression tag	UNP Q48MT7
C	334	HIS	-	expression tag	UNP Q48MT7
C	335	HIS	-	expression tag	UNP Q48MT7
C	336	HIS	-	expression tag	UNP Q48MT7
D	145	SER	GLY	engineered mutation	UNP Q48MT7
D	329	LEU	-	expression tag	UNP Q48MT7
D	330	GLU	-	expression tag	UNP Q48MT7
D	331	HIS	-	expression tag	UNP Q48MT7
D	332	HIS	-	expression tag	UNP Q48MT7
D	333	HIS	-	expression tag	UNP Q48MT7
D	334	HIS	-	expression tag	UNP Q48MT7
D	335	HIS	-	expression tag	UNP Q48MT7
D	336	HIS	-	expression tag	UNP Q48MT7

- # FAD

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 84	C 27	H 31	N 9	O 15	P 2	0	0
2	B	1	Total 84	C 27	H 31	N 9	O 15	P 2	0	0
2	C	1	Total 84	C 27	H 31	N 9	O 15	P 2	0	0



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	D	1	Total	C	H	N	O	P	0	0
			84	27	31	9	15	2		

- # NAI

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			4	1	1	2		
4	B	1	Total	C	H	O	0	0
			4	1	1	2		
4	C	1	Total	C	H	O	0	0
			4	1	1	2		
4	D	1	Total	C	H	O	0	0
			4	1	1	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	198	Total	O	0	0
			198	198		
5	B	191	Total	O	0	0
			191	191		
5	C	167	Total	O	0	0
			167	167		
5	D	180	Total	O	0	0
			180	180		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Renalase




• Molecule 1: Renalase



• Molecule 1: Renalase



• Molecule 1: Renalase



V269	F270	S271	L272	H276	A279	E310	E330	HIS	HIS	HIS	HIS	HIS	HIS
------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.50Å 71.50Å 143.80Å 90.00° 97.50° 90.00°	Depositor
Resolution (Å)	46.22 – 2.00 46.22 – 2.00	Depositor EDS
% Data completeness (in resolution range)	92.8 (46.22-2.00) 88.7 (46.22-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.00Å)	Xtriage
Refinement program	PHENIX dev_1965	Depositor
R, R_{free}	0.170 , 0.215 0.176 , 0.220	Depositor DCC
R_{free} test set	3000 reflections (3.59%)	wwPDB-VP
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	20810	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 79.25 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8182e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, NAI, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2531	0.76	1/3446 (0.0%)
1	B	0.69	0/2522	0.74	1/3434 (0.0%)
1	C	0.64	0/2556	0.77	1/3477 (0.0%)
1	D	0.66	1/2589 (0.0%)	0.75	0/3525
All	All	0.67	1/10198 (0.0%)	0.75	3/13882 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	276	TRP	CB-CG	6.21	1.61	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	302	ASP	CB-CG-OD1	6.59	124.23	118.30
1	B	302	ASP	CB-CG-OD1	5.72	123.45	118.30
1	A	40	ARG	NE-CZ-NH2	-5.20	117.70	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2463	2365	2378	19	0
1	B	2454	2357	2370	18	0
1	C	2478	2383	2388	22	1
1	D	2520	2418	2431	28	1
2	A	53	31	31	3	0
2	B	53	31	31	3	0
2	C	53	31	31	5	0
2	D	53	31	31	3	0
3	A	44	27	27	2	0
3	B	44	27	27	3	0
3	C	44	27	27	2	0
3	D	44	27	27	0	0
4	A	3	1	1	0	0
4	B	3	1	1	0	0
4	C	3	1	1	0	0
4	D	3	1	1	1	0
5	A	198	0	0	5	0
5	B	191	0	0	4	0
5	C	167	0	0	4	0
5	D	180	0	0	11	0
All	All	11051	9759	9803	85	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:VAL:N	5:B:501:HOH:O	1.85	1.06
1:D:147:ASN:ND2	5:D:501:HOH:O	1.92	1.01
1:B:239:ARG:NH1	5:B:502:HOH:O	2.01	0.91
1:B:280:ARG:HB3	3:B:402:NAI:H42N	1.63	0.80
1:D:23:SER:OG	5:D:503:HOH:O	2.00	0.79

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:318:GLU:OE2	1:D:225:THR:OG1[2_553]	1.97	0.23

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/336 (94%)	310 (98%)	5 (2%)	1 (0%)	43	39
1	B	315/336 (94%)	310 (98%)	5 (2%)	0	100	100
1	C	317/336 (94%)	314 (99%)	3 (1%)	0	100	100
1	D	326/336 (97%)	323 (99%)	3 (1%)	0	100	100
All	All	1274/1344 (95%)	1257 (99%)	16 (1%)	1 (0%)	53	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	327	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/266 (95%)	252 (100%)	0	100	100
1	B	251/266 (94%)	250 (100%)	1 (0%)	92	94
1	C	255/266 (96%)	251 (98%)	4 (2%)	65	70
1	D	258/266 (97%)	257 (100%)	1 (0%)	92	94
All	All	1016/1064 (96%)	1010 (99%)	6 (1%)	87	91

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	46	SER
1	D	173	LEU
1	C	147	ASN
1	C	44	LYS
1	C	217	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	153	HIS
1	A	326	HIS
1	B	153	HIS
1	C	153	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	401	-	50,58,58	1.33	7 (14%)	56,89,89	1.87	7 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAI	A	402	-	41,48,48	1.01	2 (4%)	45,73,73	1.79	7 (15%)
4	FMT	A	403	-	0,2,2	0.00	-	0,1,1	0.00	-
2	FAD	B	401	-	50,58,58	1.43	8 (16%)	56,89,89	1.79	7 (12%)
3	NAI	B	402	-	41,48,48	0.93	1 (2%)	45,73,73	1.60	7 (15%)
4	FMT	B	403	-	0,2,2	0.00	-	0,1,1	0.00	-
2	FAD	C	401	-	50,58,58	1.46	9 (18%)	56,89,89	2.08	8 (14%)
3	NAI	C	402	-	41,48,48	1.08	3 (7%)	45,73,73	1.71	7 (15%)
4	FMT	C	403	-	0,2,2	0.00	-	0,1,1	0.00	-
2	FAD	D	401	-	50,58,58	1.54	11 (22%)	56,89,89	2.30	7 (12%)
3	NAI	D	402	-	41,48,48	1.08	3 (7%)	45,73,73	2.17	10 (22%)
4	FMT	D	403	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	401	-	-	0/28/50/50	0/6/6/6
3	NAI	A	402	-	-	0/25/72/72	0/5/5/5
4	FMT	A	403	-	-	0/0/0/0	0/0/0/0
2	FAD	B	401	-	-	0/28/50/50	0/6/6/6
3	NAI	B	402	-	-	0/25/72/72	0/5/5/5
4	FMT	B	403	-	-	0/0/0/0	0/0/0/0
2	FAD	C	401	-	-	0/28/50/50	0/6/6/6
3	NAI	C	402	-	-	0/25/72/72	0/5/5/5
4	FMT	C	403	-	-	0/0/0/0	0/0/0/0
2	FAD	D	401	-	-	0/28/50/50	0/6/6/6
3	NAI	D	402	-	-	0/25/72/72	0/5/5/5
4	FMT	D	403	-	-	0/0/0/0	0/0/0/0

The worst 5 of 44 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FAD	C8A-N9A	-2.37	1.34	1.36
2	D	401	FAD	C4'-C3'	-2.19	1.49	1.53
2	D	401	FAD	C1'-N10	-2.11	1.46	1.48
2	C	401	FAD	C1'-N10	-2.03	1.46	1.48
2	B	401	FAD	C10-N1	2.02	1.35	1.33

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	FAD	N3A-C2A-N1A	-7.91	122.09	128.86
2	D	401	FAD	N3A-C2A-N1A	-7.54	122.41	128.86
3	A	402	NAI	N3A-C2A-N1A	-7.41	122.52	128.86
3	C	402	NAI	N3A-C2A-N1A	-6.68	123.15	128.86
3	D	402	NAI	N3A-C2A-N1A	-6.66	123.16	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	FAD	3	0
3	A	402	NAI	2	0
2	B	401	FAD	3	0
3	B	402	NAI	3	0
2	C	401	FAD	5	0
3	C	402	NAI	2	0
2	D	401	FAD	3	0
4	D	403	FMT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/336 (95%)	-0.08	8 (2%) 57 56	16, 26, 59, 111	0
1	B	319/336 (94%)	-0.12	4 (1%) 77 76	16, 25, 58, 104	0
1	C	321/336 (95%)	0.04	15 (4%) 31 31	16, 30, 70, 114	0
1	D	328/336 (97%)	-0.07	4 (1%) 79 78	17, 29, 65, 88	0
All	All	1288/1344 (95%)	-0.06	31 (2%) 59 58	16, 27, 67, 114	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	328	GLN	5.8
1	A	89	HIS	4.7
1	C	192	PHE	3.8
1	C	45	ARG	3.7
1	C	262	ASP	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAI	A	402	44/44	0.92	0.14	22,49,71,93	0
3	NAI	D	402	44/44	0.93	0.14	26,45,85,112	0
3	NAI	B	402	44/44	0.94	0.14	18,50,77,102	0
3	NAI	C	402	44/44	0.95	0.14	24,44,104,118	0
4	FMT	D	403	3/3	0.96	0.09	32,37,39,47	0
4	FMT	C	403	3/3	0.97	0.07	28,34,38,45	0
2	FAD	B	401	53/53	0.97	0.11	10,19,26,31	0
2	FAD	C	401	53/53	0.97	0.09	14,22,35,37	0
4	FMT	A	403	3/3	0.97	0.07	23,23,25,28	0
2	FAD	D	401	53/53	0.98	0.09	13,20,31,41	0
2	FAD	A	401	53/53	0.98	0.10	9,19,31,40	0
4	FMT	B	403	3/3	0.99	0.06	20,21,28,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.