



wwPDB EM Validation Summary Report ⓘ

Dec 18, 2022 – 03:56 am GMT

PDB ID : 6ZY2
EMDB ID : EMD-11547
Title : Cryo-EM structure of apo MlaFEDB
Authors : Dong, C.J.; Dong, H.H.
Deposited on : 2020-07-30
Resolution : 3.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	<div> <div>34%</div> <div>79%</div> <div>19%</div> </div>
1	D	183	<div> <div>27%</div> <div>74%</div> <div>5%</div> <div>21%</div> </div>
1	I	183	<div> <div>24%</div> <div>72%</div> <div>26%</div> </div>
1	J	183	<div> <div>38%</div> <div>74%</div> <div>5%</div> <div>21%</div> </div>
1	K	183	<div> <div>42%</div> <div>72%</div> <div>8%</div> <div>20%</div> </div>
1	L	183	<div> <div>33%</div> <div>70%</div> <div>6%</div> <div>23%</div> </div>
2	B	105	<div> <div>30%</div> <div>81%</div> <div>10%</div> <div>10%</div> </div>
2	C	105	<div> <div>31%</div> <div>84%</div> <div>8%</div> <div>9%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	269	<div><div></div><div>9%</div><div>87%</div><div>10%</div><div></div></div>
3	G	269	<div><div></div><div>7%</div><div>90%</div><div>7%</div><div></div></div>
4	E	260	<div><div></div><div>10%</div><div>90%</div><div>10%</div><div></div></div>
4	H	260	<div><div></div><div>7%</div><div>91%</div><div>9%</div><div></div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YrbD protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	149	Total	C	N	O	S	0	0
			1059	684	173	199	3		
1	D	144	Total	C	N	O	S	0	0
			1068	692	174	199	3		
1	I	136	Total	C	N	O	S	0	0
			1027	665	169	191	2		
1	J	144	Total	C	N	O	S	0	0
			1065	686	176	200	3		
1	K	146	Total	C	N	O	S	0	0
			1094	702	180	210	2		
1	L	140	Total	C	N	O	S	0	0
			1058	686	174	195	3		

- Molecule 2 is a protein called ABC transporter maintaining OM lipid asymmetry, cytoplasmic STAS component.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	95	Total	C	N	O	S	0	0
			734	465	126	140	3		
2	C	96	Total	C	N	O	S	0	0
			743	470	127	143	3		

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	98	HIS	-	expression tag	UNP W8T4U6
B	99	HIS	-	expression tag	UNP W8T4U6
B	100	HIS	-	expression tag	UNP W8T4U6
B	101	HIS	-	expression tag	UNP W8T4U6
B	102	HIS	-	expression tag	UNP W8T4U6
B	103	HIS	-	expression tag	UNP W8T4U6
B	104	HIS	-	expression tag	UNP W8T4U6
B	105	HIS	-	expression tag	UNP W8T4U6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	98	HIS	-	expression tag	UNP W8T4U6
C	99	HIS	-	expression tag	UNP W8T4U6
C	100	HIS	-	expression tag	UNP W8T4U6
C	101	HIS	-	expression tag	UNP W8T4U6
C	102	HIS	-	expression tag	UNP W8T4U6
C	103	HIS	-	expression tag	UNP W8T4U6
C	104	HIS	-	expression tag	UNP W8T4U6
C	105	HIS	-	expression tag	UNP W8T4U6

- Molecule 3 is a protein called Toluene tolerance protein Ttg2A.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	F	261	Total	C	N	O	S	0	0
			1984	1260	350	360	14		
3	G	262	Total	C	N	O	S	0	0
			1989	1263	351	361	14		


- Molecule 4 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	259	Total	C	N	O	S	0	0
			1948	1276	326	333	13		
4	H	259	Total	C	N	O	S	0	0
			1949	1278	325	333	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: YrbD protein




• Molecule 1: YrbD protein




• Molecule 1: YrbD protein




- Molecule 1: YrbD protein




- Molecule 1: YrbD protein



- Molecule 1: YrbD protein






- Molecule 2: ABC transporter maintaining OM lipid asymmetry, cytoplasmic STAS component










- Molecule 2: ABC transporter maintaining OM lipid asymmetry, cytoplasmic STAS component




- Molecule 3: Toluene tolerance protein Ttg2A




- Molecule 3: Toluene tolerance protein Ttg2A



- Molecule 4: Uncharacterized protein



- Molecule 4: Uncharacterized protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	76541	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.125	Depositor
Minimum map value	-0.067	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.02	Depositor
Map size (\AA)	259.584, 259.584, 259.584	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.014, 1.014, 1.014	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1075	0.46	0/1470
1	D	0.25	0/1083	0.51	0/1475
1	I	0.24	0/1041	0.49	0/1412
1	J	0.25	0/1081	0.47	0/1473
1	K	0.25	0/1109	0.50	0/1508
1	L	0.25	0/1074	0.46	0/1458
2	B	0.24	0/744	0.48	0/1011
2	C	0.24	0/753	0.49	0/1023
3	F	0.26	0/2024	0.51	0/2747
3	G	0.25	0/2029	0.49	0/2754
4	E	0.25	0/1986	0.45	0/2698
4	H	0.25	0/1987	0.44	0/2699
All	All	0.25	0/15986	0.48	0/21728

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1059	0	1025	3	0
1	D	1068	0	1089	7	0
1	I	1027	0	1054	4	0
1	J	1065	0	1055	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	K	1094	0	1102	7	0
1	L	1058	0	1083	8	0
2	B	734	0	756	6	0
2	C	743	0	762	4	0
3	F	1984	0	2025	17	0
3	G	1989	0	2030	11	0
4	E	1948	0	2056	16	0
4	H	1949	0	2059	16	0
All	All	15718	0	16096	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:41:CYS:HB2	2:B:75:GLN:HE22	1.65	0.62
3:F:75:MET:HB2	3:F:79:ARG:HB3	1.84	0.60
3:F:171:PRO:HD2	3:F:202:SER:HB3	1.84	0.60
4:E:240:VAL:HG13	4:H:58:ILE:HD12	1.88	0.56
4:E:95:LEU:HA	4:E:99:LEU:HB2	1.87	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/183 (79%)	141 (97%)	4 (3%)	0	100	100
1	D	140/183 (76%)	138 (99%)	2 (1%)	0	100	100
1	I	128/183 (70%)	127 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	J	140/183 (76%)	133 (95%)	7 (5%)	0	100	100
1	K	140/183 (76%)	139 (99%)	1 (1%)	0	100	100
1	L	134/183 (73%)	131 (98%)	3 (2%)	0	100	100
2	B	93/105 (89%)	92 (99%)	1 (1%)	0	100	100
2	C	94/105 (90%)	84 (89%)	10 (11%)	0	100	100
3	F	259/269 (96%)	244 (94%)	15 (6%)	0	100	100
3	G	260/269 (97%)	240 (92%)	20 (8%)	0	100	100
4	E	257/260 (99%)	248 (96%)	9 (4%)	0	100	100
4	H	257/260 (99%)	250 (97%)	7 (3%)	0	100	100
All	All	2047/2366 (86%)	1967 (96%)	80 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/150 (68%)	102 (100%)	0	100	100
1	D	111/150 (74%)	111 (100%)	0	100	100
1	I	108/150 (72%)	108 (100%)	0	100	100
1	J	107/150 (71%)	106 (99%)	1 (1%)	78	90
1	K	115/150 (77%)	114 (99%)	1 (1%)	78	90
1	L	111/150 (74%)	110 (99%)	1 (1%)	78	90
2	B	82/92 (89%)	82 (100%)	0	100	100
2	C	83/92 (90%)	83 (100%)	0	100	100
3	F	214/220 (97%)	214 (100%)	0	100	100
3	G	214/220 (97%)	214 (100%)	0	100	100
4	E	207/209 (99%)	206 (100%)	1 (0%)	88	95
4	H	207/209 (99%)	207 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1661/1942 (86%)	1657 (100%)	4 (0%)	93 98

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	138	LYS
1	K	77	THR
1	L	55	ARG
4	E	111	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	105	ASN
3	G	191	ASN
3	G	233	GLN
3	G	230	GLN
2	B	75	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	I	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	I	39:THR	C	40:TYR	N	3.64


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-11547. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128



Y Index: 128



Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 128



Y Index: 112




Z Index: 146


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 88 nm^3 ; this corresponds to an approximate mass of 80 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-11547 and PDB model 6ZY2. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

9.4 Atom inclusion [i](#)



At the recommended contour level, 69% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.5852	<div></div> 0.3790
A	<div></div> 0.4718	<div></div> 0.3130
B	<div></div> 0.5457	<div></div> 0.3710
C	<div></div> 0.5185	<div></div> 0.3470
D	<div></div> 0.5062	<div></div> 0.3460
E	<div></div> 0.6799	<div></div> 0.4280
F	<div></div> 0.7018	<div></div> 0.4350
G	<div></div> 0.7103	<div></div> 0.4430
H	<div></div> 0.7198	<div></div> 0.4480
I	<div></div> 0.4901	<div></div> 0.3220
J	<div></div> 0.4381	<div></div> 0.2940
K	<div></div> 0.3816	<div></div> 0.2940
L	<div></div> 0.4313	<div></div> 0.2880

1.0

0.0

<0.0