



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 11:38 pm BST

PDB ID : 469D  
Title : CRYSTAL STRUCTURE AND IMPROVED ANTISENSE PROPERTIES  
OF 2'-O-(2-METHOXYETHYL)-RNA  
Authors : Teplova, M.; Minasov, G.; Tereshko, V.; Inamati, G.; Cook, P.D.; Egli, M.  
Deposited on : 1999-04-29  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
RNA backbone	3102	1007 (2.38-1.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	12	
1	B	12	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	C43	B	21[B]	-	-	X	-
1	G48	B	22	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 711 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R>(\*C43)P\*(G48)P\*(C43)P\*(G48)P\*(A44)P\*(A44)P\*(U36)P\*(U36)P\*(C43)P\*(G48)P\*(C43)P\*(G48))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	12	Total	C	N	O	P	0	0	0
			301	150	46	94	11			
1	B	12	Total	C	N	O	P	0	1	0
			304	152	46	95	11			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	47	Total	O	0	0
			47	47		
3	B	58	Total	O	0	0
			58	58		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (5'-R(\*(C43)P\*(G48)P\*(C43)P\*(G48)P\*(A44)P\*(A44)P\*(U36)P\*(U36)P\*(C43)P\*(G48)P\*(C43)P\*(G48))-3')

Chain A:  8% 92%



- Molecule 1: RNA (5'-R(\*(C43)P\*(G48)P\*(C43)P\*(G48)P\*(A44)P\*(A44)P\*(U36)P\*(U36)P\*(C43)P\*(G48)P\*(C43)P\*(G48))-3')

Chain B:  33% 67%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	41.20Å 34.44Å 46.62Å 90.00° 92.40° 90.00°	Depositor
Resolution (Å)	8.00 – 1.70 7.95 – 1.70	Depositor EDS
% Data completeness (in resolution range)	88.7 (8.00-1.70) 88.7 (7.95-1.70)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.43 (at 1.70Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.206 , 0.258 0.184 , 0.240	Depositor DCC
$R_{free}$ test set	638 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.2	Xtriage
Anisotropy	0.592	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.17 , 83.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.049 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	711	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: G48, U36, MG, A44, C43

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	301	0	203	16	0
1	B	304	0	200	21	0
2	A	1	0	0	0	0
3	A	47	0	0	0	0
3	B	58	0	0	6	0
All	All	711	0	403	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:G48:H22	1:B:24:G48:HCD3	1.14	1.08

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:G48:N2	1:B:24:G48:HCD3	1.97	0.79
1:A:9:C43:HD'3	1:A:10:G48:H5'	1.67	0.76
1:B:21[B]:C43:HD'3	1:B:22:G48:C4'	2.24	0.68
1:B:13:C43:HA'1	1:B:14:G48:H5'	1.80	0.64
1:A:12:G48:CA'	3:B:121:HOH:O	2.46	0.63
1:A:9:C43:HD'3	1:A:10:G48:C5'	2.29	0.62
1:B:21[B]:C43:HD'3	1:B:22:G48:H4'	1.81	0.62
1:B:18:A44:HCD3	3:B:104:HOH:O	1.99	0.62
1:A:11:C43:HB'1	3:B:202:HOH:O	1.99	0.61
1:B:21[B]:C43:CD'	1:B:22:G48:HCA1	2.34	0.57
1:A:6:A44:HCA2	1:A:7:U36:O4'	2.07	0.54
1:A:5:A44:HCA1	1:B:21[A]:C43:CD'	2.38	0.54
1:B:21[B]:C43:HD'3	1:B:22:G48:O4'	2.07	0.54
1:A:5:A44:HCA1	1:B:21[A]:C43:HD'3	1.91	0.52
1:B:21[B]:C43:CD'	1:B:22:G48:H4'	2.40	0.52
1:A:12:G48:HCB2	3:B:121:HOH:O	2.12	0.49
1:B:21[B]:C43:HD'1	1:B:22:G48:HCA1	1.95	0.48
1:A:2:G48:H2'	1:A:3:C43:H6	1.96	0.48
1:B:21[B]:C43:HD'3	1:B:22:G48:HCA1	1.96	0.47
1:B:17:A44:O2'	1:B:18:A44:H5'	2.15	0.47
1:A:11:C43:HD'2	1:A:12:G48:C4'	2.47	0.45
1:B:21[B]:C43:HD'2	3:B:171:HOH:O	2.16	0.45
1:A:4:G48:H1'	1:A:4:G48:HCA1	1.62	0.44
1:B:21[B]:C43:HB'2	1:B:22:G48:H5'	1.98	0.44
1:A:11:C43:H1'	1:A:11:C43:HA'2	1.71	0.43
1:B:14:G48:HCA1	1:B:14:G48:H1'	1.71	0.43
1:A:12:G48:HCA1	3:B:121:HOH:O	2.16	0.42
1:A:1:C43:O5'	1:A:1:C43:H6	2.18	0.42
1:B:16:G48:O2'	1:B:17:A44:H5'	2.19	0.42
1:B:13:C43:O5'	1:B:13:C43:H6	2.21	0.41
1:B:17:A44:HCA2	1:B:18:A44:O4'	2.20	0.40
1:B:21[B]:C43:HD'3	1:B:22:G48:C1'	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains ⓘ

There are no protein molecules in this entry.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	0/12	-	-
1	B	0/12	-	-
All	All	0/24	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

25 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	G48	A	2	1	21,29,30	1.08	2 (9%)	24,41,44	2.52	6 (25%)
1	C43	A	11	1	18,25,26	0.86	0	21,34,37	0.95	1 (4%)
1	A44	B	17	1	21,28,29	0.71	0	22,39,42	0.84	1 (4%)
1	C43	B	23	1	18,25,26	0.96	0	21,34,37	1.09	1 (4%)
1	C43	A	9	1	18,25,26	0.89	0	21,34,37	0.94	1 (4%)
1	A44	A	5	1	21,28,29	0.72	0	22,39,42	0.97	1 (4%)
1	A44	A	6	1	21,28,29	0.71	0	22,39,42	0.91	1 (4%)
1	C43	B	21[B]	-	18,25,26	1.33	4 (22%)	21,34,37	1.82	2 (9%)
1	G48	B	24	1	21,29,30	1.17	2 (9%)	24,41,44	2.52	7 (29%)
1	C43	B	21[A]	-	18,25,26	1.37	4 (22%)	21,34,37	2.09	4 (19%)
1	U36	A	7	1	17,25,26	1.09	2 (11%)	18,34,37	1.14	1 (5%)
1	C43	B	15	1	18,25,26	0.89	0	21,34,37	1.00	1 (4%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	U36	A	8	1	17,25,26	1.17	2 (11%)	18,34,37	1.03	1 (5%)
1	G48	B	22	1	21,29,30	1.21	2 (9%)	24,41,44	2.41	5 (20%)
1	A44	B	18	1	21,28,29	0.78	0	22,39,42	0.81	1 (4%)
1	G48	A	10	1	21,29,30	1.28	3 (14%)	24,41,44	2.44	5 (20%)
1	G48	B	16	1	21,29,30	1.24	2 (9%)	24,41,44	2.43	5 (20%)
1	C43	A	3	1	18,25,26	0.77	0	21,34,37	0.98	1 (4%)
1	U36	B	20	1	17,25,26	1.35	2 (11%)	18,34,37	1.46	2 (11%)
1	G48	A	12	1	21,29,30	1.20	2 (9%)	24,41,44	2.47	6 (25%)
1	U36	B	19	1	17,25,26	1.14	2 (11%)	18,34,37	1.04	1 (5%)
1	G48	A	4	1	21,29,30	1.21	2 (9%)	24,41,44	2.44	5 (20%)
1	G48	B	14	1	21,29,30	1.23	1 (4%)	24,41,44	2.40	5 (20%)
1	C43	B	13	1	18,22,26	0.79	0	21,30,37	0.95	1 (4%)
1	C43	A	1	1	18,22,26	0.73	0	21,30,37	0.96	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	G48	A	2	1	-	1/8/30/31	0/3/3/3
1	C43	A	11	1	-	2/10/30/31	0/2/2/2
1	A44	B	17	1	-	2/8/30/31	0/3/3/3
1	C43	B	23	1	-	1/10/30/31	0/2/2/2
1	C43	A	9	1	-	1/10/30/31	0/2/2/2
1	A44	A	5	1	-	0/8/30/31	0/3/3/3
1	A44	A	6	1	-	1/8/30/31	0/3/3/3
1	C43	B	21[B]	-	-	3/10/30/31	0/2/2/2
1	G48	B	24	1	-	2/8/30/31	0/3/3/3
1	C43	B	21[A]	-	-	3/10/30/31	0/2/2/2
1	U36	A	7	1	-	2/10/30/31	0/2/2/2
1	C43	B	15	1	-	0/10/30/31	0/2/2/2
1	U36	A	8	1	-	1/10/30/31	0/2/2/2
1	G48	B	22	1	-	2/8/30/31	0/3/3/3
1	A44	B	18	1	-	1/8/30/31	0/3/3/3
1	G48	A	10	1	-	0/8/30/31	0/3/3/3
1	G48	B	16	1	-	0/8/30/31	0/3/3/3
1	C43	A	3	1	-	2/10/30/31	0/2/2/2

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	U36	B	20	1	-	2/10/30/31	0/2/2/2
1	G48	A	12	1	-	3/8/30/31	0/3/3/3
1	U36	B	19	1	-	2/10/30/31	0/2/2/2
1	G48	A	4	1	-	1/8/30/31	0/3/3/3
1	G48	B	14	1	-	1/8/30/31	0/3/3/3
1	C43	B	13	1	-	3/9/27/31	0/2/2/2
1	C43	A	1	1	-	1/9/27/31	0/2/2/2

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	14	G48	C6-N1	4.28	1.40	1.33
1	A	12	G48	C6-N1	4.15	1.40	1.33
1	B	16	G48	C6-N1	4.03	1.40	1.33
1	B	22	G48	C6-N1	3.99	1.40	1.33
1	B	24	G48	C6-N1	3.88	1.39	1.33
1	A	4	G48	C6-N1	3.65	1.39	1.33
1	A	10	G48	C6-N1	3.63	1.39	1.33
1	A	2	G48	C6-N1	3.52	1.39	1.33
1	B	20	U36	C4-N3	2.99	1.38	1.33
1	A	8	U36	C4-N3	2.81	1.37	1.33
1	A	7	U36	C4-N3	2.80	1.37	1.33
1	B	21[B]	C43	O2'-C2'	2.73	1.47	1.43
1	B	21[A]	C43	O2'-C2'	2.73	1.47	1.43
1	B	21[A]	C43	CB'-CA'	-2.71	1.35	1.49
1	B	19	U36	C4-N3	2.67	1.37	1.33
1	B	20	U36	C6-N1	2.59	1.39	1.35
1	A	8	U36	C6-N1	2.50	1.38	1.35
1	B	19	U36	C6-N1	2.49	1.38	1.35
1	A	7	U36	C6-N1	2.40	1.38	1.35
1	A	4	G48	C6-C5	2.34	1.45	1.41
1	B	22	G48	C8-N7	-2.32	1.30	1.34
1	B	16	G48	C8-N7	-2.26	1.30	1.34
1	B	21[B]	C43	CB'-CA'	2.20	1.60	1.49
1	A	10	G48	C6-C5	2.20	1.45	1.41
1	B	21[B]	C43	C6-C5	-2.17	1.33	1.38
1	B	21[A]	C43	C6-C5	-2.17	1.33	1.38
1	B	21[B]	C43	O2'-CA'	2.15	1.48	1.43
1	B	21[A]	C43	O2'-CA'	2.15	1.48	1.43
1	B	24	G48	C8-N7	-2.12	1.30	1.34
1	A	10	G48	C8-N7	-2.05	1.31	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	12	G48	C8-N7	-2.04	1.31	1.34
1	A	2	G48	C8-N7	-2.03	1.31	1.34

All (66) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	G48	C5-C6-N1	-8.96	111.18	123.43
1	A	12	G48	C5-C6-N1	-8.86	111.32	123.43
1	A	4	G48	C5-C6-N1	-8.85	111.32	123.43
1	B	16	G48	C5-C6-N1	-8.69	111.54	123.43
1	B	22	G48	C5-C6-N1	-8.61	111.66	123.43
1	B	24	G48	C5-C6-N1	-8.59	111.68	123.43
1	B	14	G48	C5-C6-N1	-8.54	111.75	123.43
1	A	2	G48	C5-C6-N1	-8.49	111.81	123.43
1	B	21[B]	C43	CA'-O2'-C2'	6.41	129.60	114.32
1	B	21[A]	C43	CA'-O2'-C2'	6.41	129.60	114.32
1	A	2	G48	C6-N1-C2	6.02	125.50	115.93
1	A	4	G48	C6-N1-C2	5.84	125.21	115.93
1	A	10	G48	C6-N1-C2	5.80	125.14	115.93
1	B	16	G48	C6-N1-C2	5.78	125.11	115.93
1	B	14	G48	C6-N1-C2	5.76	125.08	115.93
1	A	12	G48	C6-N1-C2	5.76	125.08	115.93
1	B	24	G48	C6-N1-C2	5.74	125.05	115.93
1	B	22	G48	C6-N1-C2	5.68	124.96	115.93
1	B	21[A]	C43	O2'-CA'-CB'	4.20	124.28	109.72
1	B	21[B]	C43	C2-N3-C4	3.75	120.15	116.34
1	B	21[A]	C43	C2-N3-C4	3.75	120.15	116.34
1	B	23	C43	C2-N3-C4	3.74	120.13	116.34
1	A	7	U36	C5-C4-N3	-3.69	115.18	123.31
1	B	20	U36	C5-C4-N3	-3.69	115.20	123.31
1	B	13	C43	C2-N3-C4	3.68	120.08	116.34
1	B	19	U36	C5-C4-N3	-3.63	115.32	123.31
1	A	11	C43	C2-N3-C4	3.57	119.96	116.34
1	A	3	C43	C2-N3-C4	3.53	119.92	116.34
1	A	8	U36	C5-C4-N3	-3.51	115.58	123.31
1	A	1	C43	C2-N3-C4	3.51	119.90	116.34
1	A	9	C43	C2-N3-C4	3.48	119.87	116.34
1	A	2	G48	CA'-O2'-C2'	3.42	122.49	114.32
1	B	20	U36	CA'-O2'-C2'	3.42	122.48	114.32
1	B	15	C43	C2-N3-C4	3.42	119.80	116.34
1	B	14	G48	N3-C2-N1	-2.93	123.31	127.22
1	B	16	G48	N3-C2-N1	-2.90	123.36	127.22

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	21[A]	C43	OC'-CB'-CA'	2.87	132.97	111.07
1	A	2	G48	C2-N3-C4	-2.83	112.12	115.36
1	B	24	G48	C2-N3-C4	-2.80	112.16	115.36
1	B	22	G48	C2-N3-C4	-2.80	112.16	115.36
1	A	2	G48	N3-C2-N1	-2.78	123.51	127.22
1	A	10	G48	C2-N3-C4	-2.77	112.20	115.36
1	A	6	A44	C5-C6-N6	2.72	124.49	120.35
1	A	12	G48	C2-N3-C4	-2.70	112.27	115.36
1	B	16	G48	C2-N3-C4	-2.68	112.29	115.36
1	A	4	G48	C6-C5-C4	-2.66	118.26	120.80
1	A	5	A44	C5-C6-N6	2.65	124.39	120.35
1	A	12	G48	C6-C5-C4	-2.64	118.28	120.80
1	A	2	G48	C6-C5-C4	-2.61	118.31	120.80
1	B	18	A44	C5-C6-N6	2.60	124.31	120.35
1	B	24	G48	O3'-C3'-C2'	2.57	118.47	111.17
1	B	22	G48	N3-C2-N1	-2.57	123.80	127.22
1	B	14	G48	C6-C5-C4	-2.56	118.35	120.80
1	B	22	G48	C6-C5-C4	-2.56	118.35	120.80
1	B	24	G48	N3-C2-N1	-2.54	123.83	127.22
1	A	4	G48	C2-N3-C4	-2.51	112.49	115.36
1	A	4	G48	N3-C2-N1	-2.49	123.90	127.22
1	B	14	G48	C2-N3-C4	-2.43	112.59	115.36
1	B	24	G48	C6-C5-C4	-2.39	118.51	120.80
1	A	12	G48	N3-C2-N1	-2.39	124.04	127.22
1	A	10	G48	C6-C5-C4	-2.39	118.52	120.80
1	B	17	A44	C5-C6-N6	2.30	123.85	120.35
1	A	10	G48	N3-C2-N1	-2.29	124.16	127.22
1	B	24	G48	O3'-C3'-C4'	2.27	117.61	111.05
1	A	12	G48	O3'-C3'-C2'	2.22	117.48	111.17
1	B	16	G48	C6-C5-C4	-2.22	118.68	120.80

There are no chirality outliers.

All (37) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	22	G48	C1'-C2'-O2'-CA'
1	A	12	G48	C1'-C2'-O2'-CA'
1	A	4	G48	C1'-C2'-O2'-CA'
1	B	14	G48	C1'-C2'-O2'-CA'
1	B	20	U36	O2'-CA'-CB'-OC'
1	B	21[B]	C43	O2'-CA'-CB'-OC'
1	A	9	C43	O2'-CA'-CB'-OC'

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
1	B	21[A]	C43	CB'-CA'-O2'-C2'
1	A	11	C43	CA'-CB'-OC'-CD'
1	A	3	C43	CA'-CB'-OC'-CD'
1	A	11	C43	C1'-C2'-O2'-CA'
1	B	23	C43	CA'-CB'-OC'-CD'
1	B	13	C43	CA'-CB'-OC'-CD'
1	B	19	U36	CA'-CB'-OC'-CD'
1	B	18	A44	CA'-CB'-OC'-CD'
1	B	21[B]	C43	CA'-CB'-OC'-CD'
1	A	8	U36	CA'-CB'-OC'-CD'
1	B	21[B]	C43	CB'-CA'-O2'-C2'
1	B	20	U36	CB'-CA'-O2'-C2'
1	A	7	U36	CA'-CB'-OC'-CD'
1	A	6	A44	CA'-CB'-OC'-CD'
1	B	17	A44	O2'-CA'-CB'-OC'
1	B	17	A44	CA'-CB'-OC'-CD'
1	B	24	G48	O2'-CA'-CB'-OC'
1	B	22	G48	O2'-CA'-CB'-OC'
1	A	7	U36	CB'-CA'-O2'-C2'
1	B	24	G48	CB'-CA'-O2'-C2'
1	A	1	C43	CA'-CB'-OC'-CD'
1	A	12	G48	CB'-CA'-O2'-C2'
1	B	13	C43	CB'-CA'-O2'-C2'
1	B	19	U36	O2'-CA'-CB'-OC'
1	A	12	G48	O2'-CA'-CB'-OC'
1	B	21[A]	C43	CA'-CB'-OC'-CD'
1	A	2	G48	C1'-C2'-O2'-CA'
1	B	13	C43	C1'-C2'-O2'-CA'
1	A	3	C43	O2'-CA'-CB'-OC'
1	B	21[A]	C43	O2'-CA'-CB'-OC'

There are no ring outliers.

20 monomers are involved in 33 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	2	G48	3	0
1	A	11	C43	3	0
1	B	17	A44	3	0
1	A	9	C43	2	0
1	A	5	A44	2	0
1	A	6	A44	1	0
1	B	21[B]	C43	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	24	G48	2	0
1	B	21[A]	C43	2	0
1	A	7	U36	1	0
1	B	22	G48	9	0
1	B	18	A44	3	0
1	A	10	G48	2	0
1	B	16	G48	1	0
1	A	3	C43	1	0
1	A	12	G48	4	0
1	A	4	G48	1	0
1	B	14	G48	2	0
1	B	13	C43	2	0
1	A	1	C43	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	0/12	-	-	-	-
1	B	0/12	-	-	-	-
All	All	0/24	-	-	-	-

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	A44	B	17	26/27	0.93	0.10	9,24,40,40	0
1	U36	B	20	24/25	0.93	0.10	10,20,40,40	0
1	A44	A	6	26/27	0.94	0.10	11,19,32,40	0
1	G48	B	24	27/28	0.94	0.10	17,30,40,40	0
1	A44	B	18	26/27	0.94	0.10	11,26,40,40	0
1	C43	A	11	24/25	0.94	0.10	7,20,40,40	0
1	G48	A	12	27/28	0.94	0.10	9,22,40,40	0
1	C43	B	23	24/25	0.95	0.10	15,21,34,40	0
1	C43	B	21[A]	24/25	0.95	0.10	10,22,28,34	3
1	U36	A	7	24/25	0.95	0.09	12,21,31,31	0
1	U36	A	8	24/25	0.95	0.09	8,17,25,30	0
1	C43	A	9	24/25	0.95	0.09	8,15,29,36	0
1	G48	A	2	27/28	0.95	0.09	17,22,33,40	0
1	C43	B	21[B]	24/25	0.95	0.10	10,21,32,34	3
1	U36	B	19	24/25	0.95	0.09	12,21,33,40	0
1	G48	A	4	27/28	0.95	0.10	11,21,34,40	0
1	C43	A	1	21/25	0.95	0.10	18,25,40,40	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	C43	A	3	24/25	0.96	0.09	14,21,31,40	0
1	G48	B	22	27/28	0.96	0.09	12,23,40,40	0
1	C43	B	13	21/25	0.96	0.10	13,21,32,39	0
1	G48	B	16	27/28	0.96	0.08	13,18,26,30	0
1	G48	B	14	27/28	0.97	0.08	12,18,33,40	0
1	C43	B	15	24/25	0.97	0.09	13,18,31,40	0
1	A44	A	5	26/27	0.97	0.09	9,22,37,40	0
1	G48	A	10	27/28	0.98	0.07	8,16,27,40	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	100	1/1	0.92	0.06	34,34,34,34	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.