



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 04:58 am BST

PDB ID : 6A06
Title : Structure of pSTING complex
Authors : Yuan, Z.L.; Shang, G.J.; Cong, X.Y.; Gu, L.C.
Deposited on : 2018-06-05
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	201	<div> <div>7%</div> <div>83%</div> <div>10%</div> <div>6%</div> </div>
1	B	201	<div> <div>7%</div> <div>79%</div> <div>10%</div> <div>10%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3253 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Stimulator of interferon genes protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	188	Total	C	N	O	S	0	2	0
			1506	947	277	276	6			
1	B	181	Total	C	N	O	S	0	1	0
			1449	914	262	267	6			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	150	ALA	-	expression tag	UNP B8XX90
A	151	ALA	-	expression tag	UNP B8XX90
A	260	GLU	GLY	conflict	UNP B8XX90
A	343	LEU	-	expression tag	UNP B8XX90
A	344	GLU	-	expression tag	UNP B8XX90
A	345	HIS	-	expression tag	UNP B8XX90
A	346	HIS	-	expression tag	UNP B8XX90
A	347	HIS	-	expression tag	UNP B8XX90
A	348	HIS	-	expression tag	UNP B8XX90
A	349	HIS	-	expression tag	UNP B8XX90
A	350	HIS	-	expression tag	UNP B8XX90
B	150	ALA	-	expression tag	UNP B8XX90
B	151	ALA	-	expression tag	UNP B8XX90
B	260	GLU	GLY	conflict	UNP B8XX90
B	343	LEU	-	expression tag	UNP B8XX90
B	344	GLU	-	expression tag	UNP B8XX90
B	345	HIS	-	expression tag	UNP B8XX90
B	346	HIS	-	expression tag	UNP B8XX90
B	347	HIS	-	expression tag	UNP B8XX90
B	348	HIS	-	expression tag	UNP B8XX90
B	349	HIS	-	expression tag	UNP B8XX90
B	350	HIS	-	expression tag	UNP B8XX90

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is cGAMP (three-letter code: 1SY) (formula: C₂₀H₂₄N₁₀O₁₃P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total 45	C 20	N 10	O 13	P 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	110	Total 110	O 110	0	0
4	B	123	Total 123	O 123	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.63Å 63.31Å 101.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.90 – 1.79 30.90 – 1.79	Depositor EDS
% Data completeness (in resolution range)	98.9 (30.90-1.79) 95.2 (30.90-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.54 (at 1.79Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.175 , 0.220 0.176 , 0.221	Depositor DCC
R_{free} test set	2000 reflections (6.60%)	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3253	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.16 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3414e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1SY, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1543	0.52	0/2089
1	B	0.39	0/1482	0.56	0/2008
All	All	0.37	0/3025	0.54	0/4097

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1506	0	1486	15	0
1	B	1449	0	1420	15	0
2	A	10	0	0	0	0
2	B	10	0	0	1	0
3	B	45	0	22	2	0
4	A	110	0	0	2	0
4	B	123	0	0	3	0
All	All	3253	0	2928	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:ARG:NH1	1:A:246:GLU:OE2	2.26	0.69
1:B:197:ARG:NH1	4:B:503:HOH:O	2.27	0.68
1:A:204:LEU:HD22	1:A:289:LYS:HD3	1.80	0.62
1:A:285:LEU:O	1:A:289:LYS:HG2	2.02	0.59
1:B:202:PHE:CE1	1:B:265:LEU:HD21	2.38	0.58
1:A:202:PHE:CE1	1:A:311:LEU:HG	2.42	0.55
1:B:281:ARG:H	1:B:281:ARG:HD3	1.74	0.52
1:A:335:GLN:HG2	1:A:338:ARG:HH12	1.75	0.52
1:B:185:ARG:NH2	4:B:507:HOH:O	2.44	0.51
1:A:227:GLN:HG2	1:A:241:THR:HG22	1.92	0.50
1:A:304:GLU:H	1:A:304:GLU:CD	2.14	0.49
1:B:209:PRO:HD3	1:B:318:THR:HG21	1.94	0.49
1:A:209:PRO:HD2	1:A:261:TYR:CZ	2.47	0.49
1:B:304:GLU:O	1:B:308:ASN:ND2	2.45	0.49
1:A:220:ARG:HD2	1:A:246:GLU:OE2	2.13	0.48
1:B:197:ARG:NH1	1:B:336:GLU:HB3	2.29	0.48
1:A:269:PHE:HA	1:A:284:ARG:HD2	1.96	0.46
3:B:403:1SY:H21	3:B:403:1SY:O26	2.16	0.45
1:B:227:GLN:HG3	4:B:588:HOH:O	2.16	0.44
1:B:269:PHE:CE2	1:B:273:GLN:HG3	2.53	0.43
1:A:163:TYR:HA	3:B:403:1SY:H16	2.01	0.43
1:A:204:LEU:HB3	1:A:289:LYS:HE3	2.01	0.42
1:B:273:GLN:O	1:B:275:GLY:N	2.53	0.42
1:B:284:ARG:NH1	2:B:402:SO4:O4	2.44	0.42
1:B:169:ARG:HH21	1:B:170:LEU:HD21	1.85	0.41
1:B:335:GLN:HA	1:B:338:ARG:NH2	2.36	0.41
1:A:210:ASP:N	1:A:210:ASP:OD1	2.53	0.41
1:B:249:GLU:OE2	1:B:327:GLN:NE2	2.51	0.41
1:A:213:SER:HB2	4:A:549:HOH:O	2.22	0.40
1:B:226:PRO:O	1:B:227:GLN:HB2	2.20	0.40
1:A:334[B]:ARG:HB2	4:A:540:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	188/201 (94%)	184 (98%)	4 (2%)	0	100	100
1	B	178/201 (89%)	170 (96%)	8 (4%)	0	100	100
All	All	366/402 (91%)	354 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/169 (94%)	158 (99%)	1 (1%)	86	84
1	B	154/169 (91%)	152 (99%)	2 (1%)	69	62
All	All	313/338 (93%)	310 (99%)	3 (1%)	76	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	308	ASN
1	B	227	GLN
1	B	281	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	402	-	4,4,4	0.15	0	6,6,6	0.08	0
3	1SY	B	403	-	43,51,51	2.71	19 (44%)	52,80,80	1.94	8 (15%)
2	SO4	A	402	-	4,4,4	0.17	0	6,6,6	0.31	0
2	SO4	A	401	-	4,4,4	0.17	0	6,6,6	0.17	0
2	SO4	B	401	-	4,4,4	0.14	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1SY	B	403	-	-	5/22/62/62	0/6/7/7

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	1SY	C36-N42	8.33	1.48	1.35
3	B	403	1SY	C38-C37	6.79	1.53	1.41
3	B	403	1SY	C40-N41	5.45	1.44	1.33
3	B	403	1SY	C38-N39	5.21	1.42	1.33
3	B	403	1SY	C40-N39	4.40	1.43	1.35
3	B	403	1SY	P27-O28	3.23	1.69	1.60
3	B	403	1SY	C6-N01	3.08	1.45	1.34
3	B	403	1SY	C22-C21	-3.05	1.46	1.52
3	B	403	1SY	O31-C32	3.04	1.45	1.41
3	B	403	1SY	P18-O17	3.03	1.71	1.59
3	B	403	1SY	C3'-C4'	-2.96	1.44	1.52
3	B	403	1SY	C2-N3	2.64	1.36	1.32
3	B	403	1SY	C5-C4	-2.63	1.34	1.40
3	B	403	1SY	O4'-C1'	2.39	1.44	1.41
3	B	403	1SY	C40-N42	2.24	1.45	1.34
3	B	403	1SY	P18-O20	2.17	1.66	1.60
3	B	403	1SY	P27-O26	2.13	1.67	1.59
3	B	403	1SY	C2'-C1'	-2.08	1.50	1.53
3	B	403	1SY	C2'-C3'	-2.05	1.48	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	403	1SY	N3-C2-N1	-5.73	119.72	128.68
3	B	403	1SY	C5-C6-N01	5.48	128.68	120.35
3	B	403	1SY	N42-C40-N39	-4.86	120.74	127.22
3	B	403	1SY	C37-C38-N39	-4.24	117.64	123.43
3	B	403	1SY	C38-N39-C40	3.99	122.27	115.93
3	B	403	1SY	N01-C6-N1	-3.71	110.88	118.57
3	B	403	1SY	C32-N33-C36	2.87	131.68	126.64
3	B	403	1SY	O19-P18-O20	2.09	115.05	106.78

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	403	1SY	C21-O20-P18-O17
3	B	403	1SY	C25-O26-P27-O28
3	B	403	1SY	C21-O20-P18-O44
3	B	403	1SY	C25-O26-P27-O29
3	B	403	1SY	C25-O26-P27-O30

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	402	SO4	1	0
3	B	403	1SY	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	188/201 (93%)	0.33	15 (7%) 12 9	18, 28, 66, 93	0
1	B	181/201 (90%)	0.35	14 (7%) 13 10	19, 30, 63, 80	0
All	All	369/402 (91%)	0.34	29 (7%) 12 9	18, 29, 66, 93	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	277	ALA	7.6
1	A	151	ALA	5.0
1	B	303	PRO	4.9
1	B	153	PHE	4.5
1	B	151	ALA	4.2
1	A	277	ALA	3.5
1	B	274	ASP	3.5
1	A	302	ALA	3.5
1	A	320	GLY	3.4
1	B	276	ARG	3.3
1	A	306	GLN	3.2
1	A	278	GLY	3.0
1	B	278	GLY	2.7
1	B	306	GLN	2.5
1	A	338	ARG	2.4
1	B	191	GLY	2.4
1	A	318	THR	2.3
1	B	152	ASN	2.2
1	B	282	GLU	2.2
1	A	193	ILE	2.2
1	B	305	ALA	2.2
1	B	301	ASP	2.2
1	A	185	ARG	2.2
1	A	303	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	305	ALA	2.1
1	A	321	GLY	2.1
1	B	307	ASN	2.1
1	A	311	LEU	2.0
1	A	209	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	401	5/5	0.86	0.27	111,113,114,115	0
2	SO4	B	402	5/5	0.88	0.16	102,105,105,108	0
2	SO4	A	401	5/5	0.94	0.11	54,55,63,65	0
2	SO4	A	402	5/5	0.96	0.08	34,40,46,50	0
3	1SY	B	403	45/45	0.97	0.10	16,22,26,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 1SY B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.