



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 11:15 pm BST

PDB ID : 1A3E  
Title : COMPLEX OF HUMAN ALPHA-THROMBIN WITH THE BIFUNCTIONAL BORONATE INHIBITOR BOROLOG2  
Authors : Skordalakes, E.; Elgendy, S.; Goodwin, C.A.; Green, D.; Scully, M.F.; Kakkar, V.V.; Freyssinet, J.M.; Dodson, G.; Deadman, J.  
Deposited on : 1998-01-21  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	36	
2	H	259	
3	I	18	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2566 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-THROMBIN (SMALL SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	27	Total	C	N	O	S	0	0	0
			222	140	36	45	1			


- Molecule 2 is a protein called ALPHA-THROMBIN (LARGE SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	251	Total	C	N	O	S	0	0	0
			2017	1288	355	360	14			

- Molecule 3 is a protein called Hirudin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	I	11	Total	C	N	O	0	0	0
			86	55	11	20			

- Molecule 4 is BOROLOG2 (three-letter code: T16) (formula:  $C_{23}H_{32}Br_2N_3O_6$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	H	1	Total	B	Br	C	N	O	0	0
			35	1	2	23	3	6		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	24	Total	O	0	0
			24	24		
5	H	178	Total	O	0	0
			178	178		
5	I	4	Total	O	0	0
			4	4		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: ALPHA-THROMBIN (SMALL SUBUNIT)

Chain L: 




- Molecule 2: ALPHA-THROMBIN (LARGE SUBUNIT)

Chain H: 



- Molecule 3: Hirudin

Chain I: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.20 Å   72.04 Å   73.01 Å 90.00°   100.93°   90.00°	Depositor
Resolution (Å)	20.00 – 1.85	Depositor
% Data completeness (in resolution range)	98.4 (20.00-1.85)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.05	Depositor
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.170 , 0.220	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2566	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: T16

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	1.02	0/224	1.24	2/298 (0.7%)
2	H	0.92	0/2069	1.20	25/2796 (0.9%)
3	I	1.21	0/88	1.13	0/118
All	All	0.94	0/2381	1.20	27/3212 (0.8%)

There are no bond length outliers.

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	175	ARG	NE-CZ-NH2	7.38	123.99	120.30
1	L	14(D)	ARG	NE-CZ-NH2	7.38	123.99	120.30
2	H	101	ARG	NE-CZ-NH2	7.34	123.97	120.30
2	H	97	ARG	NE-CZ-NH2	7.29	123.94	120.30
2	H	137	ARG	NE-CZ-NH2	7.28	123.94	120.30
2	H	50	ARG	NE-CZ-NH2	7.20	123.90	120.30
2	H	173	ARG	NE-CZ-NH2	7.11	123.85	120.30
2	H	221	ARG	NE-CZ-NH2	7.00	123.80	120.30
2	H	67	ARG	NE-CZ-NH2	7.00	123.80	120.30
2	H	73	ARG	NE-CZ-NH2	6.91	123.75	120.30
1	L	4	ARG	NE-CZ-NH2	6.88	123.74	120.30
2	H	206	ARG	NE-CZ-NH2	6.81	123.70	120.30
2	H	187	ARG	NE-CZ-NH2	6.80	123.70	120.30
2	H	77(A)	ARG	NE-CZ-NH2	6.50	123.55	120.30
2	H	126	ARG	NE-CZ-NH2	6.37	123.49	120.30
2	H	233	ARG	NE-CZ-NH2	6.34	123.47	120.30
2	H	180	MET	CG-SD-CE	6.20	110.12	100.20
2	H	26	MET	CG-SD-CE	6.17	110.07	100.20
2	H	210	MET	CG-SD-CE	6.10	109.96	100.20
2	H	32	MET	CG-SD-CE	6.09	109.95	100.20
2	H	106	MET	CG-SD-CE	6.09	109.94	100.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	93	ARG	NE-CZ-NH2	6.04	123.32	120.30
2	H	84	MET	CG-SD-CE	6.02	109.83	100.20
2	H	35	ARG	NE-CZ-NH2	5.76	123.18	120.30
2	H	201	MET	CG-SD-CE	5.72	109.35	100.20
2	H	165	ARG	NE-CZ-NH2	5.70	123.15	120.30
2	H	146	GLU	C-N-CA	5.04	134.31	121.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	222	0	225	2	0
2	H	2017	0	1978	26	0
3	I	86	0	63	3	0
4	H	35	0	29	1	0
5	H	178	0	0	5	1
5	I	4	0	0	0	0
5	L	24	0	0	1	0
All	All	2566	0	2295	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:1:T16:BR2	5:H:352:HOH:O	2.72	0.58
2:H:37:SER:HA	2:H:37(A):PRO:C	2.23	0.58
2:H:147:THR:CA	5:H:380:HOH:O	2.54	0.56
2:H:165:ARG:HD3	2:H:169:LYS:HZ2	1.71	0.56
2:H:60(G):ASN:ND2	5:H:340:HOH:O	2.40	0.55
2:H:139:THR:HG22	2:H:157:VAL:HG22	1.94	0.50
2:H:165:ARG:HB3	2:H:166:PRO:HD3	1.93	0.49

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:81:LYS:HE2	5:H:386:HOH:O	2.12	0.48
2:H:40:LEU:HD22	2:H:41:LEU:N	2.30	0.46
2:H:82:ILE:HD12	3:I:14:ILE:HG23	1.97	0.46
1:L:10:LYS:HD2	5:L:30:HOH:O	2.16	0.45
2:H:82:ILE:CD1	3:I:14:ILE:HG23	2.46	0.45
2:H:77(A):ARG:O	2:H:78:ASN:HB2	2.18	0.44
2:H:50:ARG:HG2	2:H:108:LEU:HB2	1.98	0.44
2:H:127:GLU:O	2:H:129(B):SER:OG	2.34	0.44
2:H:22:ALA:HB2	2:H:157:VAL:HG23	1.99	0.44
2:H:33:LEU:HD21	2:H:106:MET:HE3	2.01	0.43
2:H:33:LEU:HD11	2:H:106:MET:HE1	2.01	0.43
2:H:73:ARG:NH1	5:H:367:HOH:O	2.52	0.42
2:H:211:GLY:HA2	2:H:229:THR:O	2.18	0.42
2:H:129:ALA:O	2:H:130:LEU:HB2	2.20	0.42
2:H:17:VAL:O	2:H:188:GLY:HA2	2.20	0.42
3:I:15:PRO:HB2	3:I:17:GLU:OE1	2.20	0.42
2:H:50:ARG:HG3	2:H:111:PRO:HA	2.02	0.42
2:H:130:LEU:HA	2:H:130:LEU:HD23	1.92	0.41
2:H:204(B):ASN:C	2:H:204(B):ASN:HD22	2.23	0.41
1:L:1:CYS:C	2:H:122:CYS:SG	2.99	0.41
2:H:163:VAL:HB	2:H:182:CYS:SG	2.61	0.41
2:H:136:GLY:HA3	2:H:199:PHE:CZ	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:H:280:HOH:O	5:H:280:HOH:O[2_555]	1.88	0.32

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	25/36 (69%)	24 (96%)	1 (4%)	0	100	100
2	H	247/259 (95%)	239 (97%)	8 (3%)	0	100	100
3	I	9/18 (50%)	8 (89%)	0	1 (11%)	0	0
All	All	281/313 (90%)	271 (96%)	9 (3%)	1 (0%)	34	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	I	10	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	25/31 (81%)	24 (96%)	1 (4%)	31	14
2	H	216/225 (96%)	209 (97%)	7 (3%)	39	22
3	I	8/15 (53%)	7 (88%)	1 (12%)	4	0
All	All	249/271 (92%)	240 (96%)	9 (4%)	35	18

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	14(D)	ARG
2	H	40	LEU
2	H	50	ARG
2	H	64	LEU
2	H	127	GLU
2	H	129(B)	SER
2	H	182	CYS
2	H	204(B)	ASN
3	I	10	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	38	GLN
2	H	78	ASN
2	H	204(B)	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	T16	H	1	2	32,36,36	3.18	6 (18%)	40,47,47	4.44	19 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	T16	H	1	2	-	9/32/48/48	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	1	T16	BR2-C1	-12.14	1.71	1.93
4	H	1	T16	BR1-C1	-6.87	1.80	1.93
4	H	1	T16	C6-N5	6.86	1.49	1.34
4	H	1	T16	C9-N8	4.59	1.45	1.34
4	H	1	T16	C13-N12	4.39	1.43	1.34
4	H	1	T16	O9-C9	2.76	1.27	1.22

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1	T16	BR2-C1-BR1	-13.56	90.84	109.37
4	H	1	T16	C10-N12-C13	12.74	154.43	121.65
4	H	1	T16	C10-C9-N8	10.41	137.71	118.62
4	H	1	T16	O9-C9-C10	-7.97	105.27	119.66
4	H	1	T16	BR2-C1-C2	-7.64	96.13	109.11
4	H	1	T16	C11-C10-C9	6.97	124.65	109.93
4	H	1	T16	O13-C13-N12	-4.94	114.61	122.95
4	H	1	T16	C14-C13-N12	4.09	122.92	115.83
4	H	1	T16	C11-C10-N12	4.07	119.38	110.79
4	H	1	T16	BR1-C1-C2	-3.89	102.50	109.11
4	H	1	T16	C7C-N8-C7	3.82	118.06	112.00
4	H	1	T16	O9-C9-N8	-3.33	115.44	121.38
4	H	1	T16	C14-C15-C16	2.90	118.41	113.23
4	H	1	T16	C4-N5-C6	-2.75	115.66	122.77
4	H	1	T16	C3-C2-C1	-2.75	105.94	113.55
4	H	1	T16	C9-C10-N12	-2.55	102.86	108.81
4	H	1	T16	C6-C7-N8	-2.41	105.93	112.56
4	H	1	T16	C21-C11-C10	2.26	119.62	113.39
4	H	1	T16	C7A-C7B-C7C	2.13	111.03	104.98

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	1	T16	C11-C10-N12-C13
4	H	1	T16	C9-C10-C11-C21
4	H	1	T16	N12-C10-C11-C21
4	H	1	T16	O13-C13-N12-C10
4	H	1	T16	C14-C13-N12-C10
4	H	1	T16	C11-C10-C9-O9
4	H	1	T16	C14-C15-C16-C17
4	H	1	T16	O9-C9-N8-C7
4	H	1	T16	BR1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1	T16	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.