



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 06:25 am BST

PDB ID : 5A4X  
Title : The crystal structure of Arabidopsis thaliana CAR4 in complex with two calcium ions and Zn  
Authors : Diaz, M.; Albert, A.  
Deposited on : 2015-06-15  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	<div> <div>5%</div> <div> <div></div> <div>60%</div> <div>24%</div> <div>5%</div> <div>11%</div> </div> </div>
2	B	177	<div> <div>7%</div> <div> <div></div> <div>69%</div> <div>19%</div> <div>•</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2665 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AT3G17980.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	158	Total	C	N	O	S	0	0	0
			1258	792	219	241	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	120	SER	LYS	conflict	UNP Q9LVH4

- Molecule 2 is a protein called AT3G17980.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	163	Total	C	N	O	S	0	0	0
			1282	806	224	246	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	65	SER	LYS	conflict	UNP Q9LVH4
B	131	GLY	ASP	conflict	UNP Q9LVH4
B	192	SER	LYS	conflict	UNP Q9LVH4

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Ca	0	0
			2	2		
3	A	2	Total	Ca	0	0
			2	2		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Zn 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	63	Total 63	O 63	0	0
5	B	57	Total 57	O 57	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: AT3G17980



#### • Molecule 2: AT3G17980



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.12Å 88.64Å 110.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.76 – 2.20 46.76 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.76-2.20) 100.0 (46.76-2.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 2.00Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.220 , 0.266 0.214 , 0.258	Depositor DCC
$R_{free}$ test set	1230 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.7	Xtriage
Anisotropy	0.550	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2665	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.8859e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.41	0/1276	0.58	0/1729
2	B	0.42	0/1300	0.59	0/1760
All	All	0.41	0/2576	0.58	0/3489

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1258	0	1288	29	0
2	B	1282	0	1314	29	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	B	1	0	0	0	0
5	A	63	0	0	5	0
5	B	57	0	0	8	0
All	All	2665	0	2602	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:ASP:OD2	1:A:172:ARG:NH1	1.92	1.03
2:B:114:ASP:OD1	2:B:172:ARG:NH1	2.05	0.88
2:B:91:ASP:HB2	2:B:94:LEU:HG	1.65	0.78
2:B:92:SER:HA	2:B:123:ILE:HD13	1.66	0.77
2:B:51:ASP:HB2	2:B:54:SER:O	1.87	0.75
1:A:116:GLU:HG2	1:A:172:ARG:HH21	1.56	0.71
1:A:157:THR:OG1	5:A:2052:HOH:O	2.09	0.70
2:B:118:GLU:OE1	5:B:2024:HOH:O	2.11	0.69
2:B:152:GLU:OE2	5:B:2046:HOH:O	2.11	0.68
1:A:40:ARG:NH1	1:A:84:ASP:OD1	2.28	0.67
1:A:166:ASP:OD1	5:A:2005:HOH:O	2.13	0.66
1:A:187:ASP:OD1	5:A:2002:HOH:O	2.13	0.66
1:A:40:ARG:HH21	1:A:184:GLN:HE22	1.44	0.63
2:B:151:ALA:HA	2:B:170:ARG:NH2	2.13	0.62
2:B:65:SER:OG	5:B:2015:HOH:O	2.01	0.62
2:B:170:ARG:NH2	5:B:2043:HOH:O	2.31	0.61
1:A:133:LEU:HD22	1:A:137:THR:HG21	1.83	0.59
2:B:133:LEU:HD22	2:B:137:THR:HG21	1.86	0.58
2:B:51:ASP:O	2:B:52:ILE:HB	2.05	0.56
2:B:52:ILE:HG22	2:B:53:SER:N	2.19	0.56
1:A:102:ASP:HB2	1:A:112:MET:SD	2.45	0.56
1:A:38:ARG:NH2	1:A:184:GLN:OE1	2.38	0.56
2:B:99:THR:HG22	2:B:114:ASP:HB3	1.88	0.56
2:B:47:LEU:O	5:B:2005:HOH:O	2.18	0.55
1:A:91:ASP:HB3	1:A:94:LEU:HG	1.90	0.54
1:A:95:THR:OG1	1:A:118:GLU:OE2	2.23	0.54
2:B:54:SER:HB3	2:B:75:LYS:HD2	1.90	0.54
2:B:99:THR:HG22	2:B:114:ASP:CB	2.38	0.53
2:B:91:ASP:CB	2:B:94:LEU:HG	2.36	0.53
2:B:107:SER:O	5:B:2025:HOH:O	2.18	0.53
1:A:144:PRO:HD3	1:A:153:GLU:HA	1.92	0.52
2:B:95:THR:OG1	2:B:118:GLU:OE2	2.24	0.52
1:A:35:GLY:HA3	1:A:185:TRP:CZ2	2.45	0.51
1:A:130:LEU:H	1:A:130:LEU:HD12	1.76	0.51
2:B:159:VAL:HG23	5:B:2049:HOH:O	2.09	0.51
2:B:36:LEU:HD23	2:B:88:SER:HA	1.94	0.50
2:B:172:ARG:HH11	2:B:172:ARG:HB2	1.78	0.48
2:B:63:MET:HB3	2:B:87:LEU:HD13	1.95	0.47
2:B:74:ASN:O	5:B:2018:HOH:O	2.20	0.47
1:A:119:ILE:O	1:A:123:ILE:HG12	2.15	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:GLU:OE2	5:A:2063:HOH:O	2.20	0.46
1:A:57:PRO:HA	1:A:101:TYR:O	2.15	0.46
2:B:57:PRO:HA	2:B:101:TYR:O	2.16	0.46
1:A:38:ARG:HB3	1:A:184:GLN:HB2	2.00	0.44
1:A:123:ILE:O	1:A:127:ARG:HG3	2.16	0.44
1:A:188:LEU:HA	1:A:189:PRO:HD3	1.82	0.43
1:A:56:ASP:OD1	1:A:75:LYS:HE3	2.18	0.43
2:B:140:THR:HG22	2:B:141:THR:H	1.83	0.43
2:B:102:ASP:HB2	2:B:112:MET:SD	2.59	0.43
1:A:38:ARG:CZ	5:A:2003:HOH:O	2.66	0.43
2:B:125:ALA:O	2:B:128:MET:HB2	2.19	0.42
1:A:68:LEU:HD11	1:A:86:THR:O	2.19	0.42
1:A:134:PRO:O	1:A:137:THR:OG1	2.23	0.41
1:A:43:ARG:HD3	1:A:80:GLU:OE2	2.19	0.41
1:A:43:ARG:NH1	1:A:80:GLU:OE2	2.37	0.41
1:A:36:LEU:HB2	1:A:186:ILE:HB	2.02	0.41
2:B:94:LEU:HA	2:B:94:LEU:HD23	1.67	0.40
1:A:120:SER:HB3	1:A:121:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/177 (88%)	151 (97%)	5 (3%)	0	100	100
2	B	161/177 (91%)	154 (96%)	6 (4%)	1 (1%)	25	26
All	All	317/354 (90%)	305 (96%)	11 (4%)	1 (0%)	41	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	52	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/160 (91%)	128 (88%)	17 (12%)	5	4
2	B	147/159 (92%)	136 (92%)	11 (8%)	13	14
All	All	292/319 (92%)	264 (90%)	28 (10%)	8	8

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ARG
1	A	43	ARG
1	A	53	SER
1	A	75	LYS
1	A	92	SER
1	A	108	LYS
1	A	127	ARG
1	A	128	MET
1	A	129	GLN
1	A	130	LEU
1	A	133	LEU
1	A	135	SER
1	A	159	VAL
1	A	162	LYS
1	A	164	VAL
1	A	172	ARG
1	A	188	LEU
2	B	33	LEU
2	B	43	ARG
2	B	53	SER
2	B	92	SER
2	B	93	ASN
2	B	102	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	128	MET
2	B	140	THR
2	B	164	VAL
2	B	172	ARG
2	B	188	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	158/177 (89%)	0.13	9 (5%) 23 22	21, 35, 68, 79	0
2	B	163/177 (92%)	0.20	12 (7%) 14 13	20, 36, 62, 68	0
All	All	321/354 (90%)	0.16	21 (6%) 18 17	20, 36, 63, 79	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	132	GLY	7.8
2	B	92	SER	7.6
2	B	189	PRO	6.4
1	A	93	ASN	5.3
2	B	190	GLY	5.0
2	B	93	ASN	4.9
1	A	132	GLY	4.4
2	B	191	SER	4.0
1	A	136	GLY	3.4
1	A	130	LEU	3.2
1	A	133	LEU	2.7
1	A	188	LEU	2.6
2	B	131	GLY	2.6
2	B	192	SER	2.3
2	B	133	LEU	2.2
1	A	137	THR	2.2
2	B	194	LEU	2.1
2	B	134	PRO	2.1
1	A	147	ARG	2.1
2	B	193	GLY	2.1
1	A	134	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	A	1191	1/1	0.97	0.14	32,32,32,32	0
3	CA	B	1195	1/1	0.98	0.11	25,25,25,25	0
3	CA	A	1190	1/1	0.99	0.13	26,26,26,26	0
3	CA	B	1196	1/1	0.99	0.10	26,26,26,26	0
4	ZN	B	1197	1/1	0.99	0.05	43,43,43,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.