



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 09:27 AM EDT

PDB ID : 1A7P  
Title : FV FRAGMENT OF MOUSE MONOCLONAL ANTIBODY D1.3 (BALB/C, IGG1, K) ENGINEERED MUTANT PRO95L->SER ON VARIANT CHAIN L GLU81->ASP AND CHAIN H LEU312->VAL  
Authors : Marks, C.; Henrick, K.; Winter, G.  
Deposited on : 1998-03-16  
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	107	<div> <div>2%</div> <div>64%</div> <div>25%</div> <div>8%</div> <div>.</div> </div>
2	H	116	<div> <div>%</div> <div>71%</div> <div>20%</div> <div>8%</div> <div>.</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG1-KAPPA D1.3 FV (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	107	Total	C	N	O	S	0	0	0
			815	516	135	162	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	3	VAL	GLU	conflict	UNP P01635
L	50	TYR	LYS	conflict	UNP P01635
L	51	THR	ALA	conflict	UNP P01635
L	52	THR	GLN	conflict	UNP P01635
L	81	ASP	GLU	variant	UNP P01635
L	?	-	PRO	deletion	UNP P01635
L	95	SER	PRO	engineered mutation	UNP P01635
L	96	ARG	TRP	conflict	UNP P01635

- Molecule 2 is a protein called IGG1-KAPPA D1.3 FV (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	116	Total	C	N	O	S	0	0	0
			895	560	154	177	4			

- Molecule 3 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	59	Total	O	0	0
			59	59		
4	H	82	Total	O	0	0
			82	82		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IGG1-KAPPA D1.3 FV (LIGHT CHAIN)



- Molecule 2: IGG1-KAPPA D1.3 FV (HEAVY CHAIN)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.30Å 90.30Å 56.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.01 16.77 – 2.01	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-2.01) 99.0 (16.77-2.01)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.12 (at 2.00Å)	Xtriage
Refinement program	CCP4	Depositor
R, $R_{free}$	0.174 , (Not available) 0.172 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.5	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 78.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1855	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	1.01	0/835	2.12	27/1134 (2.4%)
2	H	0.99	0/915	2.28	33/1243 (2.7%)
All	All	1.00	0/1750	2.20	60/2377 (2.5%)

There are no bond length outliers.

All (60) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	266	ARG	NE-CZ-NH2	-24.50	108.05	120.30
1	L	96	ARG	NE-CZ-NH1	15.16	127.88	120.30
2	H	292	ARG	NE-CZ-NH2	13.93	127.27	120.30
1	L	1	ASP	CB-CG-OD1	13.18	130.16	118.30
2	H	302	ARG	NE-CZ-NH2	-13.07	113.76	120.30
2	H	304	ASP	CB-CG-OD1	11.31	128.48	118.30
2	H	266	ARG	NE-CZ-NH1	11.19	125.90	120.30
2	H	305	TYR	CB-CG-CD2	-10.62	114.63	121.00
1	L	96	ARG	NE-CZ-NH2	-10.22	115.19	120.30
2	H	258	ASP	CB-CG-OD2	10.12	127.41	118.30
2	H	282	MET	CA-CB-CG	9.96	130.23	113.30
2	H	201	GLN	CA-C-O	9.84	140.76	120.10
1	L	32	TYR	CB-CG-CD1	-9.15	115.51	121.00
2	H	288	ASP	CA-CB-CG	-9.08	93.42	113.40
2	H	288	ASP	CB-CG-OD2	-9.02	110.19	118.30
2	H	297	ARG	NE-CZ-NH1	8.76	124.68	120.30
1	L	103	LYS	CB-CA-C	-8.75	92.90	110.40
2	H	218	LEU	CA-CB-CG	8.59	135.06	115.30
1	L	70	GLN	CA-CB-CG	8.37	131.80	113.40
1	L	94	THR	N-CA-CB	-8.04	95.02	110.30
1	L	86	TYR	CB-CG-CD2	-7.92	116.25	121.00
1	L	87	TYR	CB-CG-CD1	-7.85	116.29	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	289	ASP	CB-CG-OD2	7.72	125.25	118.30
1	L	86	TYR	CB-CG-CD1	7.69	125.61	121.00
2	H	292	ARG	NH1-CZ-NH2	-7.32	111.35	119.40
1	L	61	ARG	NE-CZ-NH2	7.26	123.93	120.30
1	L	81	ASP	CB-CG-OD1	7.22	124.80	118.30
1	L	10	SER	N-CA-CB	7.12	121.17	110.50
2	H	297	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	L	21	ILE	CA-CB-CG2	6.70	124.30	110.90
1	L	11	LEU	CA-CB-CG	6.68	130.67	115.30
2	H	308	GLN	CB-CG-CD	6.68	128.96	111.60
1	L	21	ILE	CB-CA-C	6.60	124.80	111.60
2	H	288	ASP	OD1-CG-OD2	6.49	135.62	123.30
1	L	107	LYS	CA-C-O	-6.43	106.60	120.10
1	L	87	TYR	CB-CG-CD2	6.35	124.81	121.00
2	H	238	ARG	NE-CZ-NH2	-6.14	117.23	120.30
2	H	265	SER	CA-CB-OG	-6.13	94.66	111.20
2	H	266	ARG	NH1-CZ-NH2	6.05	126.06	119.40
2	H	201	GLN	CA-C-N	-5.93	104.16	117.20
2	H	304	ASP	CB-CG-OD2	-5.89	113.00	118.30
1	L	34	ALA	CB-CA-C	5.83	118.84	110.10
2	H	202	VAL	N-CA-C	-5.76	95.46	111.00
2	H	201	GLN	CA-CB-CG	5.67	125.87	113.40
1	L	1	ASP	CA-CB-CG	5.65	125.84	113.40
2	H	274	SER	N-CA-CB	-5.65	102.03	110.50
1	L	63	SER	N-CA-CB	5.63	118.94	110.50
1	L	78	LEU	O-C-N	5.58	131.63	122.70
2	H	294	TYR	CG-CD1-CE1	5.56	125.75	121.30
2	H	213	ALA	N-CA-CB	-5.56	102.32	110.10
1	L	14	SER	O-C-N	5.49	131.48	122.70
2	H	266	ARG	CG-CD-NE	-5.48	100.29	111.80
1	L	83	PHE	CB-CA-C	5.43	121.27	110.40
2	H	259	TYR	CB-CG-CD1	-5.32	117.81	121.00
1	L	1	ASP	OD1-CG-OD2	-5.28	113.27	123.30
2	H	288	ASP	CB-CG-OD1	-5.20	113.62	118.30
2	H	305	TYR	CD1-CE1-CZ	-5.16	115.16	119.80
1	L	72	SER	N-CA-CB	-5.09	102.86	110.50
1	L	39	LYS	CA-CB-CG	-5.08	102.22	113.40
2	H	294	TYR	CA-CB-CG	5.08	123.06	113.40

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	815	0	781	15	0
2	H	895	0	853	14	0
3	H	4	0	3	0	0
4	H	82	0	0	0	0
4	L	59	0	0	1	0
All	All	1855	0	1637	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:292:ARG:HH11	2:H:311:THR:HG22	1.36	0.91
1:L:103:LYS:HB2	1:L:103:LYS:NZ	1.95	0.80
2:H:292:ARG:NH1	2:H:311:THR:HG22	2.06	0.70
2:H:211:LEU:HD11	2:H:315:SER:HB2	1.77	0.65
1:L:76:ASN:ND2	4:L:521:HOH:O	2.28	0.65
2:H:272:ASP:OD1	2:H:274:SER:HB3	1.97	0.64
1:L:103:LYS:HB2	1:L:103:LYS:HZ2	1.62	0.62
2:H:218:LEU:HB2	2:H:285:LEU:HD11	1.82	0.61
1:L:6:GLN:HE21	1:L:21:ILE:HD11	1.66	0.60
1:L:48:VAL:HG22	1:L:54:LEU:HD23	1.85	0.58
1:L:103:LYS:HB2	1:L:103:LYS:HZ3	1.71	0.53
2:H:211:LEU:HD13	2:H:313:THR:HB	1.93	0.49
2:H:218:LEU:HD12	2:H:282:MET:HG3	1.97	0.47
2:H:211:LEU:HD12	2:H:313:THR:O	2.15	0.46
2:H:211:LEU:HD11	2:H:315:SER:CB	2.43	0.45
2:H:282:MET:HE2	2:H:282:MET:HB3	1.87	0.44
2:H:297:ARG:O	2:H:303:LEU:HA	2.19	0.43
1:L:36:TYR:OH	1:L:89:GLN:NE2	2.52	0.42
1:L:74:LYS:HB2	1:L:74:LYS:HE2	1.92	0.42
2:H:201:GLN:OE1	2:H:201:GLN:CA	2.67	0.42
1:L:2:ILE:HD12	1:L:29:ILE:HG22	2.00	0.42
1:L:12:SER:HA	1:L:105:GLU:O	2.20	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:14:SER:HB2	1:L:17:GLU:HG3	2.02	0.41
1:L:24:ARG:HA	1:L:69:THR:O	2.21	0.41
1:L:96:ARG:HB2	2:H:247:TRP:CD1	2.56	0.41
2:H:275:LYS:O	2:H:276:SER:HB2	2.20	0.41
1:L:21:ILE:HD12	1:L:35:TRP:CZ3	2.56	0.40
1:L:83:PHE:CD1	1:L:104:LEU:HD22	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	105/107 (98%)	101 (96%)	4 (4%)	0	100	100
2	H	114/116 (98%)	109 (96%)	4 (4%)	1 (1%)	17	11
All	All	219/223 (98%)	210 (96%)	8 (4%)	1 (0%)	29	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	284	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	89/91 (98%)	80 (90%)	9 (10%)	7	4
2	H	98/100 (98%)	88 (90%)	10 (10%)	7	4
All	All	187/191 (98%)	168 (90%)	19 (10%)	7	4

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	1	ASP
1	L	11	LEU
1	L	21	ILE
1	L	72	SER
1	L	81	ASP
1	L	94	THR
1	L	97	THR
1	L	103	LYS
1	L	104	LEU
2	H	201	GLN
2	H	221	THR
2	H	225	SER
2	H	276	SER
2	H	282	MET
2	H	283	ASN
2	H	284	SER
2	H	302	ARG
2	H	311	THR
2	H	315	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	89	GLN
2	H	256	ASN
2	H	283	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACY	H	400	-	1,3,3	3.74	1 (100%)	0,3,3	-	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	400	ACY	CH3-C	3.74	1.53	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	107/107 (100%)	-0.33	2 (1%) 66 65	10, 20, 30, 49	6 (5%)
2	H	116/116 (100%)	-0.43	1 (0%) 84 83	8, 17, 30, 42	3 (2%)
All	All	223/223 (100%)	-0.38	3 (1%) 77 76	8, 18, 30, 49	9 (4%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	1	ASP	2.9
1	L	107	LYS	2.4
2	H	283	ASN	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACY	H	400	4/4	0.78	0.23	46,46,47,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.