



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 09:12 am BST

PDB ID : 6A83  
Title : Crystal structure of the C-terminal periplasmic domain of EcEptC from Escherichia coli complex with Zn  
Authors : Zhao, Y.Q.; Gu, Y.J.; Cheng, W.  
Deposited on : 2018-07-06  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>8%</div> <div>9%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoethanolamine transferase EptC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	357	Total	C	N	O	S	0	0	0
			2915	1845	498	564	8			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	184	MET	-	initiating methionine	UNP P0CB39
A	185	GLY	-	expression tag	UNP P0CB39
A	186	SER	-	expression tag	UNP P0CB39
A	187	SER	-	expression tag	UNP P0CB39
A	188	HIS	-	expression tag	UNP P0CB39
A	189	HIS	-	expression tag	UNP P0CB39
A	190	HIS	-	expression tag	UNP P0CB39
A	191	HIS	-	expression tag	UNP P0CB39
A	192	HIS	-	expression tag	UNP P0CB39
A	193	HIS	-	expression tag	UNP P0CB39
A	194	SER	-	expression tag	UNP P0CB39
A	195	SER	-	expression tag	UNP P0CB39
A	196	GLY	-	expression tag	UNP P0CB39
A	197	LEU	-	expression tag	UNP P0CB39
A	198	VAL	-	expression tag	UNP P0CB39
A	199	PRO	-	expression tag	UNP P0CB39
A	200	ARG	-	expression tag	UNP P0CB39
A	201	GLY	-	expression tag	UNP P0CB39
A	202	SER	-	expression tag	UNP P0CB39
A	203	HIS	-	expression tag	UNP P0CB39
A	204	MET	-	expression tag	UNP P0CB39

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	43	Total	O	0	0
			43	43		



- Molecule 1: Phosphoethanolamine transferase EptC



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.57Å 103.57Å 122.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.69 – 2.60 47.69 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.69-2.60) 99.9 (47.69-2.60)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.71 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.212 , 0.237 0.220 , 0.244	Depositor DCC
$R_{free}$ test set	1161 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	65.1	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 33.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.030 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2960	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/2995	0.50	1/4082 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	355	GLN	N-CA-C	-7.12	91.77	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2915	0	2790	32	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	43	0	0	2	0
All	All	2960	0	2790	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:358:THR:HG22	1:A:359:GLN:H	1.15	1.07
1:A:357:ARG:HD3	1:A:357:ARG:H	1.19	1.02
1:A:358:THR:CG2	1:A:359:GLN:H	1.81	0.94
1:A:358:THR:HG22	1:A:359:GLN:N	1.73	0.94
1:A:357:ARG:O	1:A:358:THR:OG1	1.97	0.80
1:A:334:THR:HB	1:A:345:ARG:NH1	2.06	0.71
1:A:357:ARG:N	1:A:357:ARG:HD3	2.00	0.69
1:A:334:THR:HB	1:A:345:ARG:HH12	1.61	0.65
1:A:251:ARG:NH1	1:A:468:ASP:OD1	2.29	0.64
1:A:359:GLN:O	1:A:360:SER:HB3	1.99	0.62
1:A:317:MET:HG3	1:A:384:LEU:HD22	1.83	0.59
1:A:356:GLN:HG2	1:A:361:ALA:HB1	1.83	0.59
1:A:341:THR:O	1:A:345:ARG:HG2	2.08	0.54
1:A:358:THR:O	1:A:359:GLN:C	2.48	0.52
1:A:388:HIS:HE1	4:A:714:HOH:O	1.93	0.49
1:A:399:ARG:NH2	4:A:709:HOH:O	2.46	0.49
1:A:358:THR:O	1:A:359:GLN:O	2.30	0.48
1:A:356:GLN:HG2	1:A:361:ALA:CB	2.43	0.48
1:A:358:THR:CG2	1:A:359:GLN:N	2.43	0.47
1:A:268:LEU:HD21	1:A:444:ILE:HG21	1.97	0.46
1:A:249:ARG:HH22	1:A:577:GLN:C	2.19	0.45
1:A:257:TYR:CE2	1:A:259:ARG:HB2	2.51	0.45
1:A:355:GLN:O	1:A:356:GLN:HB2	2.17	0.44
1:A:287:TYR:O	1:A:291:ILE:HG22	2.17	0.44
1:A:357:ARG:H	1:A:357:ARG:CD	2.05	0.43
1:A:286:PRO:O	1:A:477:ARG:HD2	2.18	0.43
1:A:340:LEU:HD12	1:A:340:LEU:HA	1.73	0.43
1:A:505:ARG:HH22	1:A:526:ASP:CG	2.22	0.43
1:A:261:THR:O	1:A:434:LEU:HA	2.20	0.41
1:A:278:PHE:HB2	1:A:491:PHE:HB3	2.02	0.41
1:A:356:GLN:HG3	1:A:357:ARG:N	2.36	0.41
1:A:357:ARG:C	1:A:358:THR:OG1	2.59	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/394 (90%)	342 (96%)	11 (3%)	2 (1%)	25	47

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	359	GLN
1	A	357	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/356 (90%)	316 (98%)	6 (2%)	57	79

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	237	ARG
1	A	260	GLU
1	A	357	ARG
1	A	388	HIS
1	A	437	ASP
1	A	475	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	357/394 (90%)	-0.10	11 (3%) 49 42	41, 62, 96, 132	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	334	THR	5.6
1	A	335	ALA	3.8
1	A	357	ARG	3.4
1	A	221	ALA	3.2
1	A	333	MET	3.2
1	A	572	ASP	2.9
1	A	332	THR	2.5
1	A	337	ASN	2.2
1	A	577	GLN	2.1
1	A	360	SER	2.1
1	A	421	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	A	602	1/1	0.61	0.17	30,30,30,30	0
2	ZN	A	601	1/1	0.99	0.12	59,59,59,59	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.