



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 20, 2021 – 08:03 AM GMT

PDB ID : 7A9Y
Title : Structural comparison of cellular retinoic acid binding protein I and II in the presence and absence of natural and synthetic ligands
Authors : Tomlinson, C.W.E.; Cornish, K.A.S.; Pohl, E.
Deposited on : 2020-09-02
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.17.1.dev1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.17.1.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	137	<div> <div>3%</div> <div>83%</div> <div>16%</div> <div>.</div> </div>
1	BBB	137	<div> <div>4%</div> <div>90%</div> <div>9%</div> <div>..</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4628 atoms, of which 2225 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cellular retinoic acid-binding protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	BBB	136	Total	C	H	N	O	S	29	3	0
			2183	687	1078	195	216	7			
1	AAA	137	Total	C	H	N	O	S	29	2	0
			2194	689	1087	197	215	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
BBB	29	CYS	LEU	engineered mutation	UNP P29762
AAA	29	CYS	LEU	engineered mutation	UNP P29762

- Molecule 2 is MYRISTIC ACID (three-letter code: MYR) (formula: C₁₄H₂₈O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	BBB	1	Total	C	H	O	0	0
			43	14	27	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	BBB	1	Total	C	H	O	2	0
			14	3	8	3		

- Molecule 4 is N-TRIDECANOIC ACID (three-letter code: TDA) (formula: $C_{13}H_{26}O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	AAA	1	Total	C	H	O	0	0
			40	13	25	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	BBB	67	Total 67	O 67	0	0
5	AAA	87	Total 87	O 87	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cellular retinoic acid-binding protein 1



- Molecule 1: Cellular retinoic acid-binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	36.89Å 109.98Å 170.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.05 – 1.64 36.05 – 1.64	Depositor EDS
% Data completeness (in resolution range)	99.9 (36.05-1.64) 99.6 (36.05-1.64)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 1.64Å)	Xtriage
Refinement program	REFMAC 5.8.0266, REFMAC 5.8.0266	Depositor
R, R_{free}	0.191 , 0.218 0.199 , 0.224	Depositor DCC
R_{free} test set	2046 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 48.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4628	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TDA, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.75	0/1129	1.00	6/1523 (0.4%)
1	BBB	0.74	0/1127	0.95	2/1520 (0.1%)
All	All	0.74	0/2256	0.97	8/3043 (0.3%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	132	ARG	NE-CZ-NH1	9.06	124.83	120.30
1	AAA	132	ARG	NE-CZ-NH2	-7.83	116.39	120.30
1	BBB	132	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	BBB	132	ARG	NE-CZ-NH2	-6.11	117.24	120.30
1	AAA	11[A]	ARG	NE-CZ-NH1	-5.45	117.58	120.30
1	AAA	11[B]	ARG	NE-CZ-NH1	-5.45	117.58	120.30
1	AAA	136	ARG	NE-CZ-NH2	-5.37	117.61	120.30
1	AAA	112	ARG	NE-CZ-NH2	-5.29	117.66	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1107	1087	1084	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	BBB	1105	1078	1077	8	0
2	BBB	16	27	27	1	0
3	BBB	6	8	8	0	0
4	AAA	15	25	25	0	0
5	AAA	87	0	0	0	1
5	BBB	67	0	0	1	0
All	All	2403	2225	2221	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:11[B]:ARG:NH2	1:AAA:118:GLU:OE2	2.18	0.77
1:AAA:33:ALA:O	1:AAA:37:ALA:HB3	2.01	0.60
1:BBB:39:LYS:HD2	1:AAA:11[A]:ARG:HG3	1.83	0.59
1:BBB:100:LEU:HD21	5:BBB:305:HOH:O	2.04	0.58
1:BBB:3[A]:ASN:OD1	1:BBB:3[A]:ASN:N	2.30	0.57
1:AAA:2:PRO:HG3	1:AAA:91:GLU:O	2.09	0.53
1:BBB:2:PRO:HG3	1:BBB:91:GLU:O	2.12	0.49
1:AAA:10:MET:HE2	1:AAA:38:SER:O	2.13	0.48
1:AAA:68:VAL:HG11	1:AAA:88:TRP:CD1	2.51	0.45
1:AAA:39:LYS:N	1:AAA:40:PRO:HD3	2.33	0.44
1:AAA:75:GLU:HA	1:AAA:80[B]:ARG:O	2.19	0.43
1:AAA:122:THR:CG2	1:AAA:129:VAL:HG13	2.48	0.43
1:BBB:37:ALA:HB2	2:BBB:201:MYR:H62	1.99	0.43
1:AAA:10:MET:CE	1:AAA:38:SER:O	2.67	0.42
1:BBB:68:VAL:HG11	1:BBB:88:TRP:CD1	2.54	0.42
1:AAA:109:TYR:CE2	1:AAA:124:GLY:HA3	2.54	0.42
1:AAA:23:LEU:HD22	1:AAA:80[B]:ARG:HH12	1.86	0.41
1:BBB:39:LYS:N	1:BBB:40:PRO:HD3	2.36	0.41
1:BBB:80[A]:ARG:HE	1:BBB:80[A]:ARG:HB2	1.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:AAA:301:HOH:O	5:AAA:355:HOH:O[8_444]	2.00	0.20

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	137/137 (100%)	133 (97%)	4 (3%)	0	100	100
1	BBB	137/137 (100%)	135 (98%)	2 (2%)	0	100	100
All	All	274/274 (100%)	268 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	119/118 (101%)	118 (99%)	1 (1%)	81	68
1	BBB	120/118 (102%)	116 (97%)	4 (3%)	38	11
All	All	239/236 (101%)	234 (98%)	5 (2%)	60	26

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	BBB	3[A]	ASN
1	BBB	3[B]	ASN
1	BBB	15	ASN
1	BBB	137	GLU
1	AAA	104	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TDA	AAA	201	-	11,14,14	0.20	0	10,14,14	0.21	0
3	GOL	BBB	202	-	5,5,5	0.11	0	5,5,5	0.45	0
2	MYR	BBB	201	-	12,15,15	0.20	0	11,15,15	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TDA	AAA	201	-	-	3/10/12/12	-
3	GOL	BBB	202	-	-	2/4/4/4	-
2	MYR	BBB	201	-	-	3/11/13/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	BBB	202	GOL	C1-C2-C3-O3
3	BBB	202	GOL	O2-C2-C3-O3
4	AAA	201	TDA	C24-C25-C26-C27
2	BBB	201	MYR	C11-C10-C9-C8
4	AAA	201	TDA	C29-C30-C31-C32
4	AAA	201	TDA	C28-C29-C30-C31
2	BBB	201	MYR	C5-C6-C7-C8
2	BBB	201	MYR	C6-C7-C8-C9

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	BBB	201	MYR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	137/137 (100%)	0.14	4 (2%) 51 50	22, 31, 58, 74	0
1	BBB	136/137 (99%)	0.20	5 (3%) 41 39	25, 37, 57, 75	0
All	All	273/274 (99%)	0.17	9 (3%) 46 44	22, 35, 57, 75	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	101	LEU	3.3
1	BBB	2	PRO	2.9
1	BBB	47	ASP	2.8
1	BBB	91	GLU	2.5
1	BBB	104	ASP	2.4
1	AAA	37	ALA	2.3
1	AAA	38	SER	2.2
1	BBB	117	ASP	2.0
1	AAA	104	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	BBB	202	6/6	0.56	0.21	62,71,75,77	2
2	MYR	BBB	201	16/16	0.92	0.09	31,46,51,52	0
4	TDA	AAA	201	15/15	0.97	0.12	25,59,62,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.