



wwPDB X-ray Structure Validation Summary Report ⓘ

May 30, 2020 – 02:17 pm BST

PDB ID : 1ADO
Title : FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE FROM RABBIT MUSCLE
Authors : Blom, N.S.; Sygusch, J.
Deposited on : 1996-12-02
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	363	
1	B	363	
1	C	363	
1	D	363	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16926 atoms, of which 2570 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ALDOLASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	363	3398	1731	641	489	526	11	0	0	0
1	B	363	3398	1731	641	489	526	11	0	0	0
1	C	363	3398	1731	641	489	526	11	0	0	0
1	D	363	3398	1731	641	489	526	11	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	344	SER	PRO	CONFLICT	UNP P00883
B	344	SER	PRO	CONFLICT	UNP P00883
C	344	SER	PRO	CONFLICT	UNP P00883
D	344	SER	PRO	CONFLICT	UNP P00883

- Molecule 2 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula: C₃H₇O₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	O	P		
2	A	1	24	6	4	12	2	0	1
2	B	1	12	3	2	6	1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	C	1	5	4	1	0	0
3	D	1	5	4	1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	762	Total 762	O 762	0	0
4	B	682	Total 682	O 682	0	0
4	C	953	Total 953	O 953	0	0
4	D	891	Total 891	O 891	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: ALDOLASE



- Molecule 1: ALDOLASE




- Molecule 1: ALDOLASE



- Molecule 1: ALDOLASE

Chain D:  81% 18%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	163.88Å 57.47Å 85.03Å 90.00° 102.70° 90.00°	Depositor
Resolution (Å)	12.00 – 1.90	Depositor
% Data completeness (in resolution range)	85.2 (12.00-1.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.162 , 0.203	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	16926	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 13P, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2810	0.61	1/3806 (0.0%)
1	B	0.36	0/2810	0.63	1/3806 (0.0%)
1	C	0.35	0/2810	0.62	1/3806 (0.0%)
1	D	0.35	0/2810	0.60	0/3806
All	All	0.35	0/11240	0.61	3/15224 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	353	SER	N-CA-C	-5.79	95.35	111.00
1	A	300	SER	N-CA-C	-5.35	96.56	111.00
1	B	356	LEU	CA-CB-CG	5.21	127.27	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	213	TYR	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2757	641	2774	42	0
1	B	2757	641	2774	42	0
1	C	2757	641	2774	57	0
1	D	2757	641	2774	56	0
2	A	20	4	10	2	0
2	B	10	2	5	1	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	762	0	0	12	0
4	B	682	0	0	13	1
4	C	953	0	0	27	1
4	D	891	0	0	30	0
All	All	14356	2570	11111	191	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 191 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:170:LEU:HB3	4:D:1994:HOH:O	1.75	0.86
1:A:293:LYS:HG2	1:A:297:LEU:HD11	1.64	0.80
1:A:264:VAL:HG23	4:A:1314:HOH:O	1.83	0.78
1:D:208:VAL:HG13	4:D:2108:HOH:O	1.83	0.77
1:D:353:SER:HA	4:D:2066:HOH:O	1.89	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1319:HOH:O	4:C:1454:HOH:O[2_645]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/363 (99%)	341 (94%)	16 (4%)	4 (1%)	14	5
1	B	361/363 (99%)	343 (95%)	12 (3%)	6 (2%)	9	2
1	C	361/363 (99%)	337 (93%)	19 (5%)	5 (1%)	11	3
1	D	361/363 (99%)	345 (96%)	14 (4%)	2 (1%)	25	15
All	All	1444/1452 (99%)	1366 (95%)	61 (4%)	17 (1%)	13	4

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	352	ALA
1	B	353	SER
1	B	358	ILE
1	C	348	ALA
1	C	354	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/291 (100%)	283 (97%)	8 (3%)	44	38
1	B	291/291 (100%)	277 (95%)	14 (5%)	25	16
1	C	291/291 (100%)	284 (98%)	7 (2%)	49	43
1	D	291/291 (100%)	283 (97%)	8 (3%)	44	38
All	All	1164/1164 (100%)	1127 (97%)	37 (3%)	39	30

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	318	GLU
1	B	361	HIS
1	D	310	LEU
1	B	347	GLN
1	B	357	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	179	GLN
1	B	220	HIS
1	C	361	HIS
1	B	119	ASN
1	D	85	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	13P	B	1053	-	9,9,9	1.13	1 (11%)	10,12,12	1.11	1 (10%)
3	SO4	C	1326	-	4,4,4	0.75	0	6,6,6	0.87	0
2	13P	A	1104[B]	-	9,9,9	1.26	1 (11%)	10,12,12	1.41	2 (20%)
2	13P	A	1104[A]	-	9,9,9	1.90	2 (22%)	10,12,12	1.84	2 (20%)
3	SO4	D	1261	-	4,4,4	0.78	0	6,6,6	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	13P	B	1053	-	-	4/7/8/8	-
2	13P	A	1104[B]	-	-	5/7/8/8	-
2	13P	A	1104[A]	-	-	6/7/8/8	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1104[A]	13P	O1-C1	3.83	1.45	1.43
2	A	1104[A]	13P	P-O1P	3.77	1.62	1.50
2	A	1104[B]	13P	P-O1P	3.16	1.60	1.50
2	B	1053	13P	P-O1P	2.96	1.60	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1104[A]	13P	O1-P-O1P	4.41	118.86	106.47
2	A	1104[A]	13P	O2P-P-O1P	-3.09	98.57	110.68
2	A	1104[B]	13P	O1-P-O1P	2.48	113.42	106.47
2	A	1104[B]	13P	O2P-P-O1	2.44	113.23	106.73
2	B	1053	13P	O2P-P-O1	2.29	112.82	106.73

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1053	13P	C1-O1-P-O2P
2	B	1053	13P	O1-C1-C2-O2
2	A	1104[B]	13P	C1-O1-P-O1P

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1104[B]	13P	C1-O1-P-O2P
2	A	1104[B]	13P	C1-O1-P-O3P

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1053	13P	1	0
2	A	1104[A]	13P	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.