



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 11:28 pm BST

PDB ID : 2ADF  
Title : Crystal Structure and Paratope Determination of 82D6A3, an Antithrombotic Antibody Directed Against the von Willebrand factor A3-Domain  
Authors : Staelens, S.; Hadders, M.A.; Vauterin, S.; Platteau, C.; Vanhoorelbeke, K.; Huizinga, E.G.; Deckmyn, H.  
Deposited on : 2005-07-20  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	196	
2	H	218	
3	L	209	

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5165 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Von Willebrand factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	189	Total	C	N	O	S	0	0	0
			1414	894	242	272	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	916	GLY	-	EXPRESSION TAG	UNP P04275
A	917	SER	-	EXPRESSION TAG	UNP P04275
A	918	HIS	-	EXPRESSION TAG	UNP P04275
A	919	MET	-	EXPRESSION TAG	UNP P04275

- Molecule 2 is a protein called 82D6A3 IgG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	218	Total	C	N	O	S	0	0	0
			1665	1056	272	330	7			

- Molecule 3 is a protein called 82D6A3 IgG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	209	Total	C	N	O	S	0	0	0
			1624	1015	274	329	6			

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	O	S	0	0
			5	4	1		
5	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	H	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	136	Total	O	0	0
			136	136		
7	H	155	Total	O	0	0
			155	155		
7	L	151	Total	O	0	0
			151	151		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Von Willebrand factor



- Molecule 2: 82D6A3 IgG



- Molecule 3: 82D6A3 IgG



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.18 Å 89.08 Å 123.51 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.35 – 1.90 27.34 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (27.35-1.90) 99.8 (27.34-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.25 (at 1.91 Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.191 , 0.220 0.246 , 0.270	Depositor DCC
$R_{free}$ test set	3243 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.9	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5165	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACY, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/1439	0.86	10/1957 (0.5%)
2	H	0.46	0/1709	0.74	2/2333 (0.1%)
3	L	0.60	2/1662 (0.1%)	0.81	3/2253 (0.1%)
All	All	0.52	2/4810 (0.0%)	0.80	15/6543 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	209	ASN	C-OXT	13.29	1.48	1.23
3	L	122	GLU	CD-OE1	5.28	1.31	1.25

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	979	ASP	CB-CG-OD2	7.34	124.91	118.30
3	L	159	LEU	CA-CB-CG	6.80	130.94	115.30
2	H	99	ASP	CB-CG-OD2	5.67	123.41	118.30
1	A	1009	ASP	CB-CG-OD2	5.66	123.40	118.30
1	A	1082	ASP	CB-CG-OD2	5.63	123.36	118.30
1	A	1050	ASP	CB-CG-OD2	5.54	123.28	118.30
1	A	1090	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	A	922	ASP	CB-CG-OD2	5.28	123.06	118.30
3	L	70	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	1046	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	996	ASP	CB-CG-OD2	5.20	122.98	118.30
3	L	164	ASP	CB-CG-OD2	5.08	122.88	118.30
1	A	1069	ASP	CB-CG-OD2	5.04	122.84	118.30
1	A	945	ASP	CB-CG-OD2	5.03	122.83	118.30
2	H	178	ASP	CB-CG-OD2	5.01	122.81	118.30



There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1414	0	1425	10	0
2	H	1665	0	1621	13	0
3	L	1624	0	1567	14	0
4	A	4	0	3	0	0
5	H	5	0	0	0	0
5	L	5	0	0	0	0
6	H	6	0	8	1	0
7	A	136	0	0	2	0
7	H	155	0	0	4	0
7	L	151	0	0	2	0
All	All	5165	0	4624	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:200:SER:HB3	3:L:202:SER:O	1.63	0.99
2:H:161:SER:H	2:H:201:ASN:HD21	1.24	0.84
1:A:1021:GLU:HG2	7:A:1227:HOH:O	1.78	0.83
1:A:984:VAL:O	1:A:986:PRO:HD3	1.87	0.74
2:H:6:GLN:HE21	2:H:109:GLY:HA3	1.53	0.73
3:L:200:SER:CB	3:L:202:SER:O	2.37	0.72
3:L:22:THR:HG22	7:L:1049:HOH:O	1.93	0.66
2:H:161:SER:H	2:H:201:ASN:ND2	1.95	0.63
6:H:1004:GOL:H32	7:H:1066:HOH:O	1.99	0.62
2:H:157:LEU:HD11	2:H:200:CYS:HB2	1.82	0.61
3:L:16:GLY:HA2	3:L:77:ASN:HD22	1.65	0.61
1:A:952:LYS:NZ	1:A:996:ASP:OD1	2.34	0.58
2:H:157:LEU:HD13	2:H:202:VAL:HG22	1.84	0.58

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:200:SER:HB2	7:L:1024:HOH:O	2.04	0.57
2:H:113:THR:HG22	7:H:1014:HOH:O	2.02	0.57
3:L:199:THR:O	3:L:200:SER:HB2	2.06	0.55
3:L:79:GLU:OE1	3:L:81:GLU:OE2	2.27	0.53
1:A:966:GLN:HE21	1:A:983:ASN:H	1.57	0.52
3:L:202:SER:HB3	3:L:203:PRO:HD2	1.92	0.52
3:L:90:GLN:HE22	3:L:93:ASN:H	1.58	0.51
2:H:143:LEU:HD12	2:H:198:ILE:HG21	1.97	0.47
1:A:1007:ILE:HD12	1:A:1041:SER:HB2	1.97	0.46
3:L:35:TRP:CE2	3:L:73:PHE:HB2	2.51	0.45
2:H:6:GLN:HE22	2:H:95:PHE:HA	1.81	0.45
1:A:1094:LEU:HB3	1:A:1095:PRO:HD3	1.99	0.45
1:A:952:LYS:CE	1:A:996:ASP:OD1	2.64	0.45
3:L:194:GLU:HG2	3:L:205:VAL:HG22	2.00	0.44
3:L:111:ALA:HA	3:L:199:THR:HB	1.99	0.44
2:H:6:GLN:HE21	2:H:109:GLY:CA	2.28	0.42
3:L:196:THR:CG2	3:L:203:PRO:HB3	2.49	0.42
1:A:1063:GLY:HA3	1:A:1073:LEU:HD11	2.00	0.42
2:H:11:LEU:HD22	2:H:152:PRO:HG3	2.01	0.41
2:H:171:PHE:HA	2:H:172:PRO:HD3	1.96	0.41
3:L:3:GLN:HG2	3:L:26:SER:HB3	2.02	0.41
1:A:1074:ARG:NH2	7:A:1242:HOH:O	2.50	0.40
2:H:82:GLN:NE2	7:H:1158:HOH:O	2.55	0.40
1:A:1025:ALA:HB1	1:A:1031:LYS:HZ2	1.87	0.40
2:H:113:THR:HG21	7:H:1105:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	187/196 (95%)	180 (96%)	7 (4%)	0	100 100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	216/218 (99%)	210 (97%)	4 (2%)	2 (1%)	17	7
3	L	207/209 (99%)	200 (97%)	6 (3%)	1 (0%)	29	18
All	All	610/623 (98%)	590 (97%)	17 (3%)	3 (0%)	29	18

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	197	SER
3	L	197	HIS
2	H	198	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/160 (98%)	154 (99%)	2 (1%)	69	68
2	H	187/187 (100%)	187 (100%)	0	100	100
3	L	185/185 (100%)	178 (96%)	7 (4%)	33	24
All	All	528/532 (99%)	519 (98%)	9 (2%)	60	57

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1022	MET
1	A	1046	ASP
3	L	1	ASP
3	L	3	GLN
3	L	22	THR
3	L	90	GLN
3	L	104	GLU
3	L	159	LEU
3	L	201	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	959	ASN
1	A	966	GLN
2	H	6	GLN
2	H	169	HIS
2	H	201	ASN
3	L	77	ASN
3	L	90	GLN
3	L	137	ASN
3	L	160	ASN
3	L	189	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ACY	A	1112	-	1,3,3	1.43	0	0,3,3	0.00	-
5	SO4	L	1003	-	4,4,4	0.17	0	6,6,6	0.26	0
5	SO4	H	1002	-	4,4,4	0.16	0	6,6,6	0.19	0
6	GOL	H	1004	-	5,5,5	0.38	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	H	1004	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	H	1004	GOL	O1-C1-C2-C3
6	H	1004	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	1004	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.