



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 06:05 am BST

PDB ID : 6AFX
Title : Proton pyrophosphatase - E225A
Authors : Tsai, J.-Y.; Tang, K.-Z.; Sun, Y.-J.
Deposited on : 2018-08-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	766	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>••</div> </div> </div>
1	B	766	<div> <div>4%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div>•</div> </div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyrophosphate-energized vacuolar membrane proton pump.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	740	Total	C	N	O	S	0	0	0
			5427	3538	862	998	29			
1	B	740	Total	C	N	O	S	0	0	0
			5427	3538	862	998	29			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	225	ALA	GLU	engineered mutation	UNP P21616
B	225	ALA	GLU	engineered mutation	UNP P21616

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	5	Total	Mg	0	0
			5	5		
3	A	5	Total	Mg	0	0
			5	5		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	K	0	0
			1	1		
4	A	1	Total	K	0	0
			1	1		

- Molecule 5 is 2-(2-{2-[2-(2-METHOXY-ETHOXY)-ETHOXY]-ETHOXY}-ETHOXY)-ETHANOL (three-letter code: 1PG) (formula: C₁₁H₂₄O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			17	11	6		
5	A	1	Total	C	O	0	0
			17	11	6		
5	A	1	Total	C	O	0	0
			17	11	6		
5	A	1	Total	C	O	0	0
			17	11	6		
5	A	1	Total	C	O	0	0
			13	8	5		
5	B	1	Total	C	O	0	0
			17	11	6		
5	B	1	Total	C	O	0	0
			17	11	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	258	Total	O	0	0
			258	258		
6	B	230	Total	O	0	0
			230	230		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyrophosphate-energized vacuolar membrane proton pump



- Molecule 1: Pyrophosphate-energized vacuolar membrane proton pump



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	217.20Å 88.32Å 158.84Å 90.00° 125.14° 90.00°	Depositor
Resolution (Å)	29.80 – 2.30 29.80 – 2.30	Depositor EDS
% Data completeness (in resolution range)	66.2 (29.80-2.30) 82.7 (29.80-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.34 (at 2.31Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.165 , 0.199 0.165 , 0.200	Depositor DCC
R_{free} test set	2000 reflections (2.13%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11489	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, PO4, MG, 1PG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/5533	0.52	0/7519
1	B	0.39	0/5533	0.50	0/7519
All	All	0.40	0/11066	0.51	0/15038

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5427	0	5575	41	0
1	B	5427	0	5575	29	0
2	A	10	0	0	1	0
2	B	10	0	0	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	81	0	113	10	0
5	B	34	0	48	1	0
6	A	258	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	230	0	0	0	0
All	All	11489	0	11311	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:421:VAL:HG12	5:A:1012:1PG:H101	1.66	0.77
1:A:441:ARG:NH2	1:B:606:GLU:OE2	2.28	0.67
1:B:438:ASP:OD2	1:B:441:ARG:NH1	2.29	0.65
1:B:408:VAL:HG22	1:B:484:VAL:HG13	1.79	0.64
1:B:273:ASN:HB3	1:B:276:VAL:HG23	1.78	0.64
1:A:273:ASN:HB3	1:A:276:VAL:HG23	1.81	0.63
1:A:570:VAL:HG22	1:B:570:VAL:HG22	1.86	0.58
1:B:179:VAL:HG11	1:B:354:VAL:HG12	1.86	0.56
1:B:264:ARG:NH2	1:B:613:ASN:OD1	2.39	0.55
1:A:159:LEU:HD13	1:A:196:LEU:HD13	1.88	0.54
1:B:122:GLN:NE2	1:B:133:LYS:O	2.39	0.54
1:A:434:GLN:NE2	5:A:1013:1PG:H52	2.23	0.53
1:B:617:GLY:HA3	1:B:622:THR:HB	1.91	0.52
1:A:404:LEU:HD21	1:A:480:TYR:HE2	1.76	0.51
5:A:1012:1PG:H62	6:A:1196:HOH:O	2.10	0.50
1:A:706:ARG:NH2	6:A:1106:HOH:O	2.42	0.49
1:A:663:GLU:HG2	6:A:1108:HOH:O	2.12	0.49
1:A:408:VAL:HG22	1:A:484:VAL:HG13	1.94	0.48
1:A:428:ASN:OD1	5:A:1013:1PG:H82	2.14	0.48
1:B:6:LEU:HD12	1:B:7:PRO:HD2	1.95	0.47
1:A:100:MET:HB2	1:A:237:MET:SD	2.55	0.47
1:A:363:LYS:NZ	6:A:1105:HOH:O	2.40	0.47
1:A:30:TRP:HD1	1:A:163:ILE:HD13	1.79	0.47
1:A:696:TYR:OH	5:A:1013:1PG:H41	2.14	0.47
1:A:701:ALA:HB3	5:A:1013:1PG:H102	1.96	0.47
1:B:126:TYR:CE1	1:B:221:GLY:HA3	2.50	0.47
1:A:363:LYS:HA	1:A:363:LYS:HD3	1.47	0.46
1:B:404:LEU:HD21	1:B:480:TYR:HE2	1.80	0.46
1:B:643:ILE:HB	1:B:644:PRO:HD3	1.98	0.46
1:A:30:TRP:CD1	1:A:163:ILE:HD13	2.51	0.45
1:A:273:ASN:HB3	1:A:276:VAL:CG2	2.45	0.45
1:A:438:ASP:OD2	1:A:441:ARG:NH1	2.50	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:460:ILE:HD11	1:B:588:TRP:CD1	2.52	0.44
1:B:629:THR:O	1:B:633:ILE:HG13	2.17	0.44
1:A:352:LYS:HB2	1:A:352:LYS:HE3	1.80	0.44
1:A:6:LEU:HA	1:A:7:PRO:HD3	1.83	0.44
1:B:67:ASP:HB3	1:B:70:VAL:HG12	2.00	0.44
1:A:701:ALA:CB	5:A:1013:1PG:H102	2.47	0.43
5:A:1011:1PG:H72	5:A:1011:1PG:H52	1.71	0.43
1:A:687:GLY:HA3	1:A:730:LYS:HB3	1.99	0.43
1:B:212:PHE:HB3	1:B:223:LEU:HD13	2.00	0.43
1:A:606:GLU:OE1	1:B:441:ARG:NH2	2.50	0.43
1:A:423:GLU:HG3	1:A:535:THR:HG21	2.01	0.43
1:B:196:LEU:HA	1:B:196:LEU:HD23	1.89	0.43
1:A:404:LEU:HA	1:A:404:LEU:HD12	1.91	0.43
1:A:168:ASN:HB2	1:A:508:ASN:HB3	2.00	0.43
1:A:430:TYR:CE1	5:A:1012:1PG:H51	2.54	0.42
1:A:699:ALA:CB	5:A:1013:1PG:H101	2.49	0.42
1:A:479:MET:SD	1:A:557:GLY:HA3	2.59	0.42
1:B:100:MET:HB2	1:B:237:MET:SD	2.59	0.42
1:B:626:ASP:O	1:B:629:THR:HG22	2.19	0.42
1:A:174:GLU:HG3	1:A:185:THR:OG1	2.19	0.42
1:A:624:LYS:HA	1:A:624:LYS:HD3	1.76	0.42
1:B:562:ARG:HA	1:B:562:ARG:HD2	1.87	0.42
1:B:38:LEU:HD23	1:B:173:LEU:HD12	2.02	0.42
1:A:570:VAL:CG2	1:B:570:VAL:HG22	2.48	0.41
1:A:663:GLU:O	1:A:666:SER:HB2	2.19	0.41
1:B:648:LEU:HA	5:B:1009:1PG:H91	2.01	0.41
1:B:470:ILE:O	1:B:474:PHE:HB2	2.20	0.41
1:B:240:PHE:HE2	1:B:650:MET:HE2	1.86	0.41
1:A:470:ILE:O	1:A:474:PHE:HB2	2.21	0.41
2:B:1001:PO4:P	2:B:1002:PO4:O1	2.78	0.41
1:A:643:ILE:HB	1:A:644:PRO:HD3	2.02	0.41
1:B:162:LYS:HA	1:B:162:LYS:HD3	1.90	0.41
1:B:250:LYS:HD3	1:B:250:LYS:HA	1.87	0.41
1:A:295:LEU:HD13	1:A:295:LEU:HA	1.90	0.41
1:A:181:LYS:HD3	1:A:181:LYS:HA	1.89	0.40
2:A:1001:PO4:P	2:A:1002:PO4:O1	2.79	0.40
1:A:560:VAL:HG13	1:A:565:ILE:HB	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	736/766 (96%)	723 (98%)	13 (2%)	0	100	100
1	B	736/766 (96%)	724 (98%)	12 (2%)	0	100	100
All	All	1472/1532 (96%)	1447 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	567/587 (97%)	552 (97%)	15 (3%)	46	63
1	B	567/587 (97%)	556 (98%)	11 (2%)	57	73
All	All	1134/1174 (97%)	1108 (98%)	26 (2%)	50	67

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	LEU
1	A	106	LEU
1	A	179	VAL
1	A	223	LEU
1	A	285	VAL
1	A	287	ASP
1	A	363	LYS
1	A	474	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	484	VAL
1	A	502	TYR
1	A	508	ASN
1	A	673	LEU
1	A	677	VAL
1	A	738	ASN
1	A	765	ILE
1	B	31	LEU
1	B	34	SER
1	B	179	VAL
1	B	223	LEU
1	B	287	ASP
1	B	308	VAL
1	B	474	PHE
1	B	502	TYR
1	B	508	ASN
1	B	677	VAL
1	B	738	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 23 ligands modelled in this entry, 12 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	1PG	A	1011	-	16,16,16	0.88	0	15,15,15	0.39	0
5	1PG	A	1012	-	16,16,16	0.81	0	15,15,15	0.48	0
2	PO4	B	1001	3,4	4,4,4	1.03	1 (25%)	6,6,6	0.57	0
5	1PG	A	1013	-	12,12,16	0.94	0	11,11,15	0.47	0
5	1PG	A	1009	-	16,16,16	0.85	0	15,15,15	0.59	0
5	1PG	B	1010	-	16,16,16	0.84	0	15,15,15	0.48	0
5	1PG	A	1010	-	16,16,16	0.86	0	15,15,15	0.50	0
2	PO4	A	1001	3,4	4,4,4	1.35	1 (25%)	6,6,6	0.77	0
5	1PG	B	1009	-	16,16,16	0.90	0	15,15,15	0.41	0
2	PO4	A	1002	3	4,4,4	0.34	0	6,6,6	0.97	0
2	PO4	B	1002	3	4,4,4	0.47	0	6,6,6	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	1PG	B	1009	-	-	7/14/14/14	-
5	1PG	A	1012	-	-	7/14/14/14	-
5	1PG	A	1013	-	-	8/10/10/14	-
5	1PG	A	1009	-	-	7/14/14/14	-
5	1PG	B	1010	-	-	8/14/14/14	-
5	1PG	A	1010	-	-	6/14/14/14	-
5	1PG	A	1011	-	-	10/14/14/14	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	PO4	P-O1	2.63	1.57	1.50
2	B	1001	PO4	P-O1	2.02	1.55	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (53) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1011	1PG	C7-C6-O3-C5
5	A	1012	1PG	C4-C5-O3-C6
5	A	1011	1PG	O1-C2-C3-O2
5	B	1010	1PG	O4-C8-C9-O5
5	A	1011	1PG	O4-C8-C9-O5
5	A	1010	1PG	O4-C8-C9-O5
5	A	1012	1PG	O1-C2-C3-O2
5	A	1009	1PG	C2-C3-O2-C4
5	A	1013	1PG	O3-C6-C7-O4
5	B	1009	1PG	O1-C2-C3-O2
5	A	1009	1PG	O1-C2-C3-O2
5	A	1011	1PG	C6-C7-O4-C8
5	A	1009	1PG	O5-C10-C11-O6
5	A	1010	1PG	O2-C4-C5-O3
5	A	1011	1PG	O3-C6-C7-O4
5	A	1010	1PG	O1-C2-C3-O2
5	A	1009	1PG	O2-C4-C5-O3
5	B	1009	1PG	O4-C8-C9-O5
5	A	1010	1PG	O5-C10-C11-O6
5	A	1012	1PG	O2-C4-C5-O3
5	B	1010	1PG	O1-C2-C3-O2
5	B	1010	1PG	O2-C4-C5-O3
5	B	1009	1PG	C2-C3-O2-C4
5	A	1009	1PG	O3-C6-C7-O4
5	B	1009	1PG	C11-C10-O5-C9
5	A	1010	1PG	C4-C5-O3-C6
5	A	1012	1PG	C7-C6-O3-C5
5	B	1010	1PG	C6-C7-O4-C8
5	A	1011	1PG	C11-C10-O5-C9
5	A	1011	1PG	C9-C8-O4-C7
5	A	1013	1PG	C7-C6-O3-C5
5	B	1009	1PG	C8-C9-O5-C10
5	A	1012	1PG	C8-C9-O5-C10
5	A	1013	1PG	C8-C9-O5-C10
5	A	1013	1PG	C11-C10-O5-C9
5	A	1009	1PG	C7-C6-O3-C5
5	A	1012	1PG	O3-C6-C7-O4
5	A	1009	1PG	C11-C10-O5-C9
5	A	1012	1PG	C5-C4-O2-C3
5	B	1010	1PG	C7-C6-O3-C5
5	A	1011	1PG	O5-C10-C11-O6
5	A	1013	1PG	C9-C8-O4-C7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	1010	1PG	C4-C5-O3-C6
5	B	1010	1PG	C5-C4-O2-C3
5	B	1009	1PG	C3-C2-O1-C1
5	A	1010	1PG	O3-C6-C7-O4
5	A	1011	1PG	C8-C9-O5-C10
5	A	1013	1PG	C4-C5-O3-C6
5	B	1009	1PG	O3-C6-C7-O4
5	A	1013	1PG	O5-C10-C11-O6
5	A	1011	1PG	O2-C4-C5-O3
5	B	1010	1PG	C2-C3-O2-C4
5	A	1013	1PG	O4-C8-C9-O5


There are no ring outliers.

8 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1011	1PG	1	0
5	A	1012	1PG	3	0
2	B	1001	PO4	1	0
5	A	1013	1PG	6	0
2	A	1001	PO4	1	0
5	B	1009	1PG	1	0
2	A	1002	PO4	1	0
2	B	1002	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	740/766 (96%)	-0.45	15 (2%) 65 71	8, 20, 38, 68	0
1	B	740/766 (96%)	-0.21	30 (4%) 37 44	8, 25, 45, 75	0
All	All	1480/1532 (96%)	-0.33	45 (3%) 50 57	8, 22, 42, 75	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	41	VAL	7.2
1	B	41	VAL	6.7
1	B	766	PHE	5.4
1	A	40	ALA	5.1
1	B	2	GLY	5.1
1	B	40	ALA	4.9
1	B	68	HIS	4.7
1	A	67	ASP	4.6
1	A	348	PHE	4.2
1	B	69	ASN	3.9
1	A	2	GLY	3.8
1	A	622	THR	3.8
1	B	622	THR	3.8
1	A	68	HIS	3.7
1	B	39	SER	3.6
1	A	766	PHE	3.5
1	B	623	ALA	3.2
1	A	765	ILE	3.2
1	B	31	LEU	3.2
1	B	119	THR	3.1
1	B	67	ASP	3.0
1	A	39	SER	2.9
1	B	348	PHE	2.9
1	A	349	PHE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	214	ILE	2.8
1	B	90	PHE	2.8
1	A	3	ALA	2.8
1	B	620	GLU	2.7
1	B	614	THR	2.7
1	A	623	ALA	2.7
1	B	504	PRO	2.6
1	B	765	ILE	2.6
1	A	616	PRO	2.6
1	B	624	LYS	2.5
1	A	119	THR	2.5
1	B	616	PRO	2.5
1	B	120	SER	2.4
1	B	9	LEU	2.4
1	B	70	VAL	2.4
1	B	618	LEU	2.4
1	B	520	HIS	2.3
1	B	114	VAL	2.2
1	B	615	ILE	2.2
1	B	505	ILE	2.1
1	B	3	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	1PG	A	1009	17/17	0.79	0.27	19,39,55,66	0
5	1PG	A	1011	17/17	0.81	0.31	33,49,59,59	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	1PG	B	1009	17/17	0.82	0.26	29,37,56,58	0
5	1PG	B	1010	17/17	0.84	0.27	33,44,60,65	0
5	1PG	A	1010	17/17	0.84	0.18	27,41,56,67	0
5	1PG	A	1013	13/17	0.88	0.35	28,34,40,45	0
5	1PG	A	1012	17/17	0.89	0.24	28,40,56,60	0
2	PO4	B	1001	5/5	0.92	0.15	11,15,20,44	0
2	PO4	A	1001	5/5	0.93	0.15	5,8,15,36	0
3	MG	B	1005	1/1	0.96	0.20	15,15,15,15	0
3	MG	A	1006	1/1	0.96	0.20	15,15,15,15	0
3	MG	B	1004	1/1	0.96	0.16	22,22,22,22	0
3	MG	A	1003	1/1	0.97	0.14	13,13,13,13	0
3	MG	B	1007	1/1	0.97	0.14	18,18,18,18	0
3	MG	B	1006	1/1	0.98	0.20	18,18,18,18	0
4	K	B	1008	1/1	0.98	0.10	27,27,27,27	0
2	PO4	B	1002	5/5	0.99	0.16	7,13,16,16	0
2	PO4	A	1002	5/5	0.99	0.17	7,9,11,12	0
3	MG	A	1005	1/1	0.99	0.19	11,11,11,11	0
3	MG	A	1004	1/1	0.99	0.17	16,16,16,16	0
3	MG	B	1003	1/1	0.99	0.14	13,13,13,13	0
4	K	A	1008	1/1	0.99	0.10	21,21,21,21	0
3	MG	A	1007	1/1	0.99	0.15	12,12,12,12	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 1PG A 1009:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 1PG A 1011:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 1PG B 1009:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 1PG B 1010:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 1PG A 1010:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 1PG A 1013:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.