



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 02:48 pm BST

PDB ID : 2AGV  
Title : Crystal structure of the SR CA<sup>2+</sup>-ATPASE with BHQ and TG  
Authors : Toyoshima, C.; Obara, K.; Norimatsu, Y.  
Deposited on : 2005-07-27  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	994	
1	B	994	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 15830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sarcoplasmic/endoplasmic reticulum calcium ATPase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	994	7671	4876	1287	1451	57	0	0	0
1	B	994	7671	4876	1287	1451	57	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	994	GLY	-	SEE REMARK 999	UNP P04191
B	994	GLY	-	SEE REMARK 999	UNP P04191

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is OCTANOIC ACID [3S-[3ALPHA, 3ABETA, 4ALPHA, 6BETA, 6ABETA, 7BETA, 8ALPHA(Z), 9BALPHA]]-6-(ACETYLOXY)-2,3,-3A,4,5,6,6A,7,8,9B-DECAHYDRO-3,3A-DIHYDROXY-3,6,9-TRIMETHYL-8-[(2-METHYL-1-OXO-2-BUTENYL)OXY]-2-OXO-4-(1-OXOBUTOXY)-AZULENO[4,5-B]FURAN-7-YL ESTER (three-letter code: TG1) (formula: C<sub>34</sub>H<sub>50</sub>O<sub>12</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total	C	O	0	0
			46	34	12		
3	B	1	Total	C	O	0	0
			46	34	12		

- Molecule 4 is 2,5-DITERT-BUTYLBENZENE-1,4-DIOL (three-letter code: BHQ) (formula:  $C_{14}H_{22}O_2$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	A	1	Total	C	O	0	0
			16	14	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			16	14	2		

- Molecule 5 is PHOSPHATIDYLETHANOLAMINE (three-letter code: PTY) (formula:  $C_{40}H_{80}NO_8P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			19	9	1	8	1		
5	A	1	Total	C	N	O	P	0	0
			19	9	1	8	1		
5	A	1	Total	C	N	O	P	0	0
			19	9	1	8	1		
5	B	1	Total	C	N	O	P	0	0
			19	9	1	8	1		
5	B	1	Total	C	N	O	P	0	0
			19	9	1	8	1		
5	B	1	Total	C	N	O	P	0	0
			19	9	1	8	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	124	Total	O	0	0
			124	124		
6	B	124	Total	O	0	0
			124	124		





## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.39Å 71.39Å 591.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.98 – 2.40	Depositor
% Data completeness (in resolution range)	99.8 (14.98-2.40)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.239 , 0.263	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	15830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BHQ, NA, TG1, PTY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/7812	0.64	1/10592 (0.0%)
1	B	0.41	0/7812	0.64	2/10592 (0.0%)
All	All	0.42	0/15624	0.64	3/21184 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	505	ARG	NE-CZ-NH2	7.30	123.95	120.30
1	B	505	ARG	NE-CZ-NH2	7.30	123.95	120.30
1	B	512	MET	CG-SD-CE	5.91	109.65	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7671	0	7764	331	0
1	B	7671	0	7764	330	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	46	0	50	3	0
3	B	46	0	50	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	16	0	22	1	0
4	B	16	0	22	1	0
5	A	57	0	33	2	0
5	B	57	0	33	2	0
6	A	124	0	0	7	0
6	B	124	0	0	7	0
All	All	15830	0	15738	660	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 660 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:567:ARG:HD3	1:B:570:PRO:HA	1.42	1.01
1:A:865:VAL:HB	1:A:868:HIS:HB2	1.42	1.01
1:B:865:VAL:HB	1:B:868:HIS:HB2	1.41	0.99
1:A:567:ARG:HD3	1:A:570:PRO:HA	1.42	0.98
1:B:126:MET:HE2	1:B:139:ARG:HD3	1.49	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	992/994 (100%)	920 (93%)	61 (6%)	11 (1%)	14	20
1	B	992/994 (100%)	921 (93%)	60 (6%)	11 (1%)	14	20
All	All	1984/1988 (100%)	1841 (93%)	121 (6%)	22 (1%)	14	20

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	883	PHE
1	B	883	PHE
1	A	185	VAL
1	A	663	LEU
1	A	866	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	840/840 (100%)	785 (94%)	55 (6%)	17	27
1	B	840/840 (100%)	788 (94%)	52 (6%)	18	29
All	All	1680/1680 (100%)	1573 (94%)	107 (6%)	17	28

5 of 107 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	816	ILE
1	B	96	LEU
1	B	760	PHE
1	A	838	MET
1	A	968	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	919	ASN
1	B	108	GLN
1	B	875	GLN
1	A	880	HIS
1	A	914	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	BHQ	B	1104	-	16,16,16	1.89	6 (37%)	26,26,26	1.33	4 (15%)
5	PTY	B	1113	-	18,18,49	1.21	3 (16%)	21,23,54	1.29	3 (14%)
5	PTY	B	1111	-	18,18,49	1.12	2 (11%)	21,23,54	1.46	3 (14%)
3	TG1	B	1103	-	43,48,48	1.49	10 (23%)	44,72,72	1.66	9 (20%)
5	PTY	A	1011	-	18,18,49	1.09	1 (5%)	21,23,54	1.47	3 (14%)
5	PTY	A	1012	-	18,18,49	1.46	3 (16%)	21,23,54	1.54	3 (14%)
4	BHQ	A	1004	-	16,16,16	1.87	5 (31%)	26,26,26	1.32	4 (15%)
3	TG1	A	1003	-	43,48,48	1.45	8 (18%)	44,72,72	1.65	9 (20%)
5	PTY	A	1013	-	18,18,49	1.20	3 (16%)	21,23,54	1.29	3 (14%)
5	PTY	B	1112	-	18,18,49	1.46	3 (16%)	21,23,54	1.53	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BHQ	B	1104	-	-	0/12/12/12	0/1/1/1
5	PTY	B	1113	-	-	7/20/20/53	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PTY	B	1111	-	-	13/20/20/53	-
3	TG1	B	1103	-	-	8/33/99/99	0/3/3/3
5	PTY	A	1011	-	-	13/20/20/53	-
5	PTY	A	1012	-	-	10/20/20/53	-
4	BHQ	A	1004	-	-	0/12/12/12	0/1/1/1
3	TG1	A	1003	-	-	8/33/99/99	0/3/3/3
5	PTY	A	1013	-	-	7/20/20/53	-
5	PTY	B	1112	-	-	10/20/20/53	-

The worst 5 of 44 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1104	BHQ	C1-C2	4.17	1.45	1.40
4	A	1004	BHQ	C1-C2	4.10	1.45	1.40
3	A	1003	TG1	O4-C21	3.96	1.29	1.21
3	B	1103	TG1	O4-C21	3.47	1.28	1.21
4	B	1104	BHQ	C6-C5	3.34	1.44	1.39

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1103	TG1	C10-O9-C32	5.86	135.44	121.53
3	A	1003	TG1	C10-O9-C32	5.70	135.07	121.53
5	A	1011	PTY	O7-C8-C11	4.50	119.36	111.09
5	B	1111	PTY	O7-C8-C11	4.43	119.24	111.09
5	A	1012	PTY	O4-C1-C6	4.35	121.11	108.43

There are no chirality outliers.

5 of 76 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	1012	PTY	O4-C1-C6-O7
5	A	1012	PTY	N1-C2-C3-O11
5	B	1112	PTY	O4-C1-C6-O7
5	B	1112	PTY	N1-C2-C3-O11
5	B	1113	PTY	C11-C8-O7-C6



There are no ring outliers.


6 monomers are involved in 12 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1104	BHQ	1	0
3	B	1103	TG1	3	0
5	A	1012	PTY	2	0
4	A	1004	BHQ	1	0
3	A	1003	TG1	3	0
5	B	1112	PTY	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.