



wwPDB EM Validation Summary Report ⓘ

Nov 29, 2022 – 12:51 pm GMT

PDB ID : 8AGE
EMDB ID : EMD-15421
Title : Structure of yeast oligosaccharyltransferase complex with acceptor peptide bound
Authors : Ramirez, A.S.; de Capitani, M.; Pesciullesi, G.; Kowal, J.; Bloch, J.S.; Irobalieva, R.N.; Aebi, M.; Reymond, J.L.; Locher, K.P.
Deposited on : 2022-07-19
Resolution : 2.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	718	
2	B	65	
3	C	86	
4	D	130	
5	E	476	
6	F	283	
7	G	430	
8	H	350	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	P	7	 71% 29%
10	I	2	 100%
10	J	2	 50% 50%

2 Entry composition

There are 19 unique types of molecules in this entry. The entry contains 17534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dolichyl-diphosphooligosaccharide--protein glycotransferase.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	644	Total	C	N	O	S	0	0
			5191	3432	830	906	23		

- Molecule 2 is a protein called OST4 isoform 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	33	Total	C	N	O	S	0	0
			251	161	37	49	4		

There are 29 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	37	ARG	-	expression tag	UNP A0A8H8UM72
B	38	THR	-	expression tag	UNP A0A8H8UM72
B	39	LEU	-	expression tag	UNP A0A8H8UM72
B	40	GLN	-	expression tag	UNP A0A8H8UM72
B	41	VAL	-	expression tag	UNP A0A8H8UM72
B	42	ASP	-	expression tag	UNP A0A8H8UM72
B	43	GLY	-	expression tag	UNP A0A8H8UM72
B	44	GLY	-	expression tag	UNP A0A8H8UM72
B	45	SER	-	expression tag	UNP A0A8H8UM72
B	46	GLY	-	expression tag	UNP A0A8H8UM72
B	47	GLY	-	expression tag	UNP A0A8H8UM72
B	48	SER	-	expression tag	UNP A0A8H8UM72
B	49	LEU	-	expression tag	UNP A0A8H8UM72
B	50	GLU	-	expression tag	UNP A0A8H8UM72
B	51	VAL	-	expression tag	UNP A0A8H8UM72
B	52	LEU	-	expression tag	UNP A0A8H8UM72
B	53	PHE	-	expression tag	UNP A0A8H8UM72
B	54	GLN	-	expression tag	UNP A0A8H8UM72
B	55	GLY	-	expression tag	UNP A0A8H8UM72
B	56	PRO	-	expression tag	UNP A0A8H8UM72
B	57	THR	-	expression tag	UNP A0A8H8UM72

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	58	GLU	-	expression tag	UNP A0A8H8UM72
B	59	THR	-	expression tag	UNP A0A8H8UM72
B	60	SER	-	expression tag	UNP A0A8H8UM72
B	61	GLN	-	expression tag	UNP A0A8H8UM72
B	62	VAL	-	expression tag	UNP A0A8H8UM72
B	63	ALA	-	expression tag	UNP A0A8H8UM72
B	64	PRO	-	expression tag	UNP A0A8H8UM72
B	65	ALA	-	expression tag	UNP A0A8H8UM72

- Molecule 3 is a protein called Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit OST5.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	85	Total	C	N	O	S	0	0
			666	448	99	118	1		

- Molecule 4 is a protein called Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit OST2.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	109	Total	C	N	O	S	0	0
			883	595	141	141	6		

- Molecule 5 is a protein called Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	438	Total	C	N	O	S	0	0
			3537	2294	565	671	7		

- Molecule 6 is a protein called Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	258	Total	C	N	O	S	0	0
			1892	1242	313	334	3		

- Molecule 7 is a protein called Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit WBP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	394	Total	C	N	O	S	0	0
			3201	2053	528	616	4		


- Molecule 8 is a protein called OST3 isoform 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	133	Total	C	N	O	S	0	0
			1067	724	161	176	6		

- Molecule 9 is a protein called PEPTIDE.


Mol	Chain	Residues	Atoms				AltConf	Trace
9	P	7	Total	C	N	O	0	1
			44	26	8	10		

- Molecule 10 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
10	I	2	Total	C	N	O	0	0
			28	16	2	10		
10	J	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 11 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				AltConf
11	A	1	Total	C	N	O	0
			14	8	1	5	
11	G	1	Total	C	N	O	0
			42	24	3	15	
11	G	1	Total	C	N	O	0
			42	24	3	15	
11	G	1	Total	C	N	O	0
			42	24	3	15	

- Molecule 12 is 1-PALMITOYL-2-LINOLEOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: CPL) (formula: C₄₂H₈₀NO₈P).



Mol	Chain	Residues	Atoms					AltConf
12	A	1	Total	C	N	O	P	0
			52	42	1	8	1	
12	B	1	Total	C	N	O	P	0
			52	42	1	8	1	
12	E	1	Total	C	N	O	P	0
			52	42	1	8	1	
12	F	1	Total	C	N	O	P	0
			52	42	1	8	1	

- Molecule 13 is PHOSPHATIDYLETHANOLAMINE (three-letter code: PTY) (formula: $C_{40}H_{80}NO_8P$).




Mol	Chain	Residues	Atoms					AltConf
13	A	1	Total	C	N	O	P	0
			42	32	1	8	1	

- Molecule 14 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
14	A	1	Total	Mg	0
			1	1	

- Molecule 15 is (2 {S},3 {R},4 {R},5 {S},6 {S})-2-(hydroxymethyl)-6-[(1 {S},2 {R},3 {R},4 {R},5' {S},6 {S},7 {R},8 {S},9 {R},12 {R},13 {R},15 {S},16 {S},18 {R})-5',7,9,13-tetramethyl-3,15-bis(oxidanyl)spiro[5-oxapentacyclo[10.8.0.0^{2,9}.0^{4,8}.0^{13,18}]icosane-6,2'-oxane]-16-yl]oxy-oxane-3,4,5-triol (three-letter code: KZB) (formula: $C_{33}H_{54}O_{10}$).




Mol	Chain	Residues	Atoms			AltConf
15	A	1	Total	C	O	0
			129	99	30	
15	A	1	Total	C	O	0
			129	99	30	
15	A	1	Total	C	O	0
			129	99	30	
15	B	1	Total	C	O	0
			86	66	20	
15	B	1	Total	C	O	0
			86	66	20	
15	F	1	Total	C	O	0
			86	66	20	
15	F	1	Total	C	O	0
			86	66	20	
15	G	1	Total	C	O	0
			43	33	10	

- Molecule 16 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			AltConf
16	F	1	Total	C	O	0
			33	18	15	
16	F	1	Total	C	O	0
			33	18	15	
16	F	1	Total	C	O	0
			33	18	15	
16	G	1	Total	C	O	0
			11	6	5	

- Molecule 17 is beta-D-mannopyranose (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			AltConf
17	G	1	Total	C	O	0
			11	6	5	

- Molecule 18 is 2-[3,6-bis(dimethylamino)xanthen-9-yl]-5-methanoyl-benzoate (three-letter code: 323) (formula: C₂₅H₂₂N₂O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
18	P	1	Total	C	N	O	0
			31	25	2	4	


- Molecule 19 is water.

Mol	Chain	Residues	Atoms		AltConf
19	A	4	Total	O	0
			4	4	
19	C	1	Total	O	0
			1	1	
19	E	2	Total	O	0
			2	2	
19	G	2	Total	O	0
			2	2	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dolichyl-diphosphooligosaccharide--protein glycotransferase




- Molecule 2: OST4 isoform 1





- Molecule 3: Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit OST5





- Molecule 4: Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit OST2




- Molecule 5: Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 1




- Molecule 6: Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 2




- Molecule 7: Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit WBP1






• Molecule 8: OST3 isoform 1

Chain H: 33% 5% 62%



• Molecule 9: PEPTIDE

Chain P: 71% 29%



• Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 100%



• Molecule 10: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	72993	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48.4	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.865	Depositor
Minimum map value	-0.884	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.050	Depositor
Recommended contour level	0.312	Depositor
Map size (Å)	326.40002, 326.40002, 326.40002	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.8500001, 0.8500001, 0.8500001	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTY, CPL, KZB, MAN, 323, NAG, MG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/5343	0.45	0/7265
2	B	0.26	0/253	0.40	0/342
3	C	0.30	0/684	0.40	0/926
4	D	0.28	0/905	0.42	0/1220
5	E	0.27	0/3639	0.46	0/4954
6	F	0.28	0/1934	0.46	0/2637
7	G	0.26	0/3283	0.43	0/4459
8	H	0.28	0/1098	0.43	0/1494
9	P	0.31	0/44	0.50	0/60
All	All	0.28	0/17183	0.44	0/23357

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5191	0	5158	61	0
2	B	251	0	260	3	0
3	C	666	0	677	11	0
4	D	883	0	911	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	E	3537	0	3412	40	0
6	F	1892	0	1831	38	0
7	G	3201	0	3101	41	0
8	H	1067	0	1099	11	0
9	P	44	0	36	2	0
10	I	28	0	25	0	0
10	J	28	0	25	0	0
11	A	14	0	12	0	0
11	G	42	0	38	1	0
12	A	52	0	80	1	0
12	B	52	0	80	4	0
12	E	52	0	80	4	0
12	F	52	0	80	1	0
13	A	42	0	60	0	0
14	A	1	0	0	0	0
15	A	129	0	0	0	0
15	B	86	0	0	0	0
15	F	86	0	0	0	0
15	G	43	0	0	0	0
16	F	33	0	29	0	0
16	G	11	0	9	0	0
17	G	11	0	8	0	0
18	P	31	0	21	0	0
19	A	4	0	0	0	0
19	C	1	0	0	0	0
19	E	2	0	0	0	0
19	G	2	0	0	0	0
All	All	17534	0	17032	208	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 208 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:98:LYS:HB3	6:F:105:MET:HB3	1.55	0.88
7:G:140:ILE:HD11	7:G:179:ALA:HB2	1.68	0.76
1:A:32:ARG:NH1	1:A:160:SER:OG	2.20	0.74
1:A:656:THR:HG22	1:A:658:LEU:H	1.55	0.71
6:F:129:PRO:HB2	6:F:153:GLN:HB3	1.71	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	636/718 (89%)	609 (96%)	25 (4%)	2 (0%)	41	72
2	B	31/65 (48%)	31 (100%)	0	0	100	100
3	C	83/86 (96%)	82 (99%)	1 (1%)	0	100	100
4	D	107/130 (82%)	101 (94%)	6 (6%)	0	100	100
5	E	434/476 (91%)	414 (95%)	20 (5%)	0	100	100
6	F	256/283 (90%)	234 (91%)	21 (8%)	1 (0%)	34	66
7	G	392/430 (91%)	380 (97%)	12 (3%)	0	100	100
8	H	131/350 (37%)	126 (96%)	5 (4%)	0	100	100
9	P	5/7 (71%)	5 (100%)	0	0	100	100
All	All	2075/2545 (82%)	1982 (96%)	90 (4%)	3 (0%)	54	81

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	45	GLU
1	A	710	LYS
6	F	168	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	551/613 (90%)	542 (98%)	9 (2%)	62	88
2	B	30/55 (54%)	29 (97%)	1 (3%)	38	72
3	C	74/75 (99%)	73 (99%)	1 (1%)	67	90
4	D	96/115 (84%)	92 (96%)	4 (4%)	30	63
5	E	394/426 (92%)	381 (97%)	13 (3%)	38	72
6	F	181/246 (74%)	172 (95%)	9 (5%)	24	56
7	G	359/392 (92%)	348 (97%)	11 (3%)	40	74
8	H	115/316 (36%)	114 (99%)	1 (1%)	78	94
9	P	4/4 (100%)	4 (100%)	0	100	100
All	All	1804/2242 (80%)	1755 (97%)	49 (3%)	48	78

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	F	100	ASN
6	F	221	ASN
6	F	108	TYR
6	F	121	GLN
7	G	67	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
7	G	222	GLN
8	H	263	GLN
7	G	226	ASN
7	G	273	HIS
3	C	80	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	NAG	I	1	5,10	14,14,15	0.32	0	17,19,21	0.39	0
10	NAG	I	2	10	14,14,15	0.22	0	17,19,21	0.51	0
10	NAG	J	1	5,10	14,14,15	0.42	0	17,19,21	1.29	1 (5%)
10	NAG	J	2	10	14,14,15	0.23	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	I	1	5,10	-	0/6/23/26	0/1/1/1
10	NAG	I	2	10	-	0/6/23/26	0/1/1/1
10	NAG	J	1	5,10	-	4/6/23/26	0/1/1/1
10	NAG	J	2	10	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	J	1	NAG	C1-O5-C5	4.84	118.75	112.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	J	2	NAG	C4-C5-C6-O6
10	J	1	NAG	O5-C5-C6-O6
10	J	2	NAG	O5-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
10	J	1	NAG	C4-C5-C6-O6
10	J	1	NAG	C8-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 1 is monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	NAG	G	501	11,17	14,14,15	0.26	0	17,19,21	0.54	0
16	MAN	G	503	17,16	11,11,12	0.75	0	15,15,17	1.19	2 (13%)
11	NAG	A	801	11,1	14,14,15	0.41	0	17,19,21	0.49	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
15	KZB	A	807	-	49,49,49	0.62	0	74,80,80	1.57	10 (13%)
18	323	P	101	9	34,34,34	2.28	9 (26%)	45,50,50	1.30	6 (13%)
12	CPL	B	103	-	51,51,51	1.28	5 (9%)	57,59,59	1.06	2 (3%)
15	KZB	F	306	-	49,49,49	0.61	0	74,80,80	1.68	11 (14%)
15	KZB	A	805	-	49,49,49	0.69	0	74,80,80	1.92	12 (16%)
11	NAG	G	505	7	14,14,15	1.06	2 (14%)	17,19,21	1.13	1 (5%)
11	NAG	G	504	7	14,14,15	0.48	0	17,19,21	1.29	1 (5%)
17	BMA	G	502	11,16	11,11,12	0.62	0	15,15,17	0.88	0
15	KZB	B	102	-	49,49,49	0.65	0	74,80,80	1.80	13 (17%)
16	MAN	F	301	16	11,11,12	0.65	0	15,15,17	1.02	2 (13%)
15	KZB	A	806	-	49,49,49	0.66	0	74,80,80	2.42	19 (25%)
15	KZB	F	305	-	49,49,49	0.61	0	74,80,80	2.01	16 (21%)
16	MAN	F	302	16	11,11,12	0.63	0	15,15,17	0.96	2 (13%)
13	PTY	A	803	-	41,41,49	0.93	4 (9%)	44,46,54	1.08	2 (4%)
12	CPL	F	304	-	51,51,51	1.29	5 (9%)	57,59,59	1.06	2 (3%)
15	KZB	G	506	7	49,49,49	0.57	0	74,80,80	1.73	10 (13%)
15	KZB	B	101	-	49,49,49	0.69	0	74,80,80	1.80	13 (17%)
12	CPL	E	501	-	51,51,51	1.28	6 (11%)	57,59,59	1.01	2 (3%)
12	CPL	A	802	-	51,51,51	1.27	5 (9%)	57,59,59	0.94	1 (1%)
16	MAN	F	303	17	11,11,12	0.62	0	15,15,17	1.00	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	NAG	G	501	11,17	-	0/6/23/26	0/1/1/1
16	MAN	G	503	17,16	-	2/2/19/22	0/1/1/1
11	NAG	A	801	11,1	-	2/6/23/26	0/1/1/1
15	KZB	A	807	-	-	3/6/121/121	0/7/7/7
18	323	P	101	9	-	10/16/69/69	0/4/4/4
12	CPL	B	103	-	-	20/55/55/55	-
15	KZB	F	306	-	-	2/6/121/121	0/7/7/7
15	KZB	A	805	-	-	2/6/121/121	0/7/7/7
11	NAG	G	505	7	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	NAG	G	504	7	-	0/6/23/26	0/1/1/1
17	BMA	G	502	11,16	-	0/2/19/22	0/1/1/1
15	KZB	B	102	-	-	4/6/121/121	0/7/7/7
16	MAN	F	301	16	-	0/2/19/22	0/1/1/1
15	KZB	A	806	-	-	6/6/121/121	0/7/7/7
15	KZB	F	305	-	-	0/6/121/121	0/7/7/7
16	MAN	F	302	16	-	0/2/19/22	0/1/1/1
13	PTY	A	803	-	-	18/45/45/53	-
12	CPL	F	304	-	-	22/55/55/55	-
15	KZB	G	506	7	-	3/6/121/121	0/7/7/7
15	KZB	B	101	-	-	4/6/121/121	0/7/7/7
12	CPL	E	501	-	-	21/55/55/55	-
12	CPL	A	802	-	-	10/55/55/55	-
16	MAN	F	303	17	-	0/2/19/22	0/1/1/1

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
18	P	101	323	O1-C4	6.80	1.49	1.38
18	P	101	323	C7-N2	5.32	1.50	1.37
18	P	101	323	O1-C3	4.86	1.46	1.38
18	P	101	323	C18-C1	3.75	1.54	1.49
18	P	101	323	C23-C24	3.69	1.57	1.49

The worst 5 of 129 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	A	806	KZB	C84-C03-C04	-9.36	94.77	104.88
15	F	305	KZB	O05-C04-C13	8.92	128.53	110.17
15	B	101	KZB	O05-C04-C13	8.73	128.14	110.17
15	A	805	KZB	O05-C04-C13	8.72	128.12	110.17
15	A	807	KZB	O05-C04-C13	8.48	127.63	110.17

There are no chirality outliers.

5 of 131 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
12	E	501	CPL	O4P-C4-C5-N
12	F	304	CPL	O2-C2-C3-O3
12	F	304	CPL	O4P-C4-C5-N

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
13	A	803	PTY	O4-C1-C6-O7
13	A	803	PTY	C3-O11-P1-O13

There are no ring outliers.


5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	G	501	NAG	1	0
12	B	103	CPL	4	0
12	F	304	CPL	1	0
12	E	501	CPL	4	0
12	A	802	CPL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand 323 P 101




Bond lengths




Bond angles






Torsions






Rings

















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-15421. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 206




Y Index: 194




Z Index: 200


6.3.2 Raw map



X Index: 206



Y Index: 194




Z Index: 200

The images above show the largest variance slices of the map in three orthogonal directions.


6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.312. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 85 nm^3 ; this corresponds to an approximate mass of 77 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.357 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.357 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.80	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.81	6.94	3.92

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.81 differs from the reported value 2.8 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-15421 and PDB model 8AGE. Per-residue inclusion information can be found in section 3 on page 12.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.312 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.312).

9.4 Atom inclusion [i](#)



At the recommended contour level, 75% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.312) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7614	<div></div> 0.5710
A	<div></div> 0.8827	<div></div> 0.6100
B	<div></div> 0.7912	<div></div> 0.5650
C	<div></div> 0.7428	<div></div> 0.5580
D	<div></div> 0.8110	<div></div> 0.5630
E	<div></div> 0.7385	<div></div> 0.5700
F	<div></div> 0.5186	<div></div> 0.4830
G	<div></div> 0.7238	<div></div> 0.5680
H	<div></div> 0.8129	<div></div> 0.5770
I	<div></div> 0.9286	<div></div> 0.6130
J	<div></div> 0.1429	<div></div> 0.4020
P	<div></div> 0.7838	<div></div> 0.6040

1.0

0.0

<0.0