



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:52 pm BST

PDB ID : 3AHE  
Title : Phosphoketolase from Bifidobacterium Breve complexed with dihydroxyethyl thiamine diphosphate  
Authors : Suzuki, R.; Katayama, T.; Kim, B.-J.; Wakagi, T.; Shoun, H.; Ashida, H.; Yamamoto, K.; Fushinobu, S.  
Deposited on : 2010-04-22  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	845	 84% 10% • 5%

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	802	Total	C	N	O	S	0	0	0
			6379	4053	1085	1217	24			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP D6PAH1
A	-18	GLY	-	EXPRESSION TAG	UNP D6PAH1
A	-17	SER	-	EXPRESSION TAG	UNP D6PAH1
A	-16	SER	-	EXPRESSION TAG	UNP D6PAH1
A	-15	HIS	-	EXPRESSION TAG	UNP D6PAH1
A	-14	HIS	-	EXPRESSION TAG	UNP D6PAH1
A	-13	HIS	-	EXPRESSION TAG	UNP D6PAH1
A	-12	HIS	-	EXPRESSION TAG	UNP D6PAH1
A	-11	HIS	-	EXPRESSION TAG	UNP D6PAH1
A	-10	HIS	-	EXPRESSION TAG	UNP D6PAH1
A	-9	SER	-	EXPRESSION TAG	UNP D6PAH1
A	-8	SER	-	EXPRESSION TAG	UNP D6PAH1
A	-7	GLY	-	EXPRESSION TAG	UNP D6PAH1
A	-6	LEU	-	EXPRESSION TAG	UNP D6PAH1
A	-5	VAL	-	EXPRESSION TAG	UNP D6PAH1
A	-4	PRO	-	EXPRESSION TAG	UNP D6PAH1
A	-3	ARG	-	EXPRESSION TAG	UNP D6PAH1
A	-2	GLY	-	EXPRESSION TAG	UNP D6PAH1
A	-1	SER	-	EXPRESSION TAG	UNP D6PAH1
A	0	HIS	-	EXPRESSION TAG	UNP D6PAH1

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-[3-[(4-AMINO-2-METHYL-5-PYRIMIDINYL)METHYL]-2-(1,2-DIHYDROXYETHYL)-4-METHYL-1,3-THIAZOL-3-IUM-5-YL]ETHYL TRIHYDROGEN DIPHOSPHATE (three-letter code: THD) (formula: C<sub>14</sub>H<sub>22</sub>N<sub>4</sub>O<sub>9</sub>P<sub>2</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			30	14	4	9	2	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $\text{C}_2\text{H}_6\text{O}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is NONAETHYLENE GLYCOL (three-letter code: 2PE) (formula: C<sub>18</sub>H<sub>38</sub>O<sub>10</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			28	18	10		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	797	Total O 797 797	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	173.53Å 173.53Å 163.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.07 – 2.10 49.07 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.07-2.10) 99.7 (49.07-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.75 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.161 , 0.206 0.163 , 0.207	Depositor DCC
$R_{free}$ test set	3637 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.7	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7280	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, THD, 2PE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.32	20/6553 (0.3%)	1.02	28/8909 (0.3%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	134	SER	CB-OG	-7.96	1.31	1.42
1	A	195	GLN	CG-CD	6.51	1.66	1.51
1	A	617	ALA	CA-CB	6.44	1.66	1.52
1	A	185	ALA	CA-CB	6.39	1.65	1.52
1	A	5	VAL	N-CA	6.26	1.58	1.46
1	A	732	VAL	CB-CG2	6.12	1.65	1.52
1	A	19	GLU	CG-CD	6.12	1.61	1.51
1	A	543	VAL	CB-CG1	5.94	1.65	1.52
1	A	433	PHE	CG-CD1	5.80	1.47	1.38
1	A	237	PHE	CE2-CZ	5.59	1.48	1.37
1	A	701	ALA	CA-CB	5.53	1.64	1.52
1	A	447	TYR	CD2-CE2	5.50	1.47	1.39
1	A	165	TYR	CD1-CE1	5.45	1.47	1.39
1	A	597	LYS	CB-CG	5.40	1.67	1.52
1	A	705	VAL	CB-CG2	5.29	1.64	1.52
1	A	712	TYR	CB-CG	5.19	1.59	1.51
1	A	419	CYS	CB-SG	-5.11	1.73	1.81
1	A	331	GLU	CG-CD	5.05	1.59	1.51
1	A	654	GLU	CB-CG	5.04	1.61	1.52
1	A	217	TYR	CD1-CE1	5.04	1.47	1.39

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	766	ASP	CB-CG-OD2	-9.78	109.49	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	766	ASP	CB-CG-OD1	8.38	125.84	118.30
1	A	230	ASP	CB-CG-OD2	-7.68	111.39	118.30
1	A	715	ASP	CB-CG-OD2	-7.25	111.77	118.30
1	A	230	ASP	CB-CG-OD1	7.02	124.62	118.30
1	A	15	ARG	NE-CZ-NH1	6.93	123.77	120.30
1	A	43	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	A	60	ARG	NE-CZ-NH2	6.70	123.65	120.30
1	A	754	ARG	NE-CZ-NH2	-6.48	117.06	120.30
1	A	676	ASP	CB-CG-OD1	6.05	123.74	118.30
1	A	54	ARG	NE-CZ-NH2	-5.89	117.35	120.30
1	A	618	ARG	NE-CZ-NH1	5.89	123.24	120.30
1	A	385	ASP	CB-CG-OD1	5.83	123.55	118.30
1	A	665	MET	CA-CB-CG	5.67	122.95	113.30
1	A	392	ASP	CB-CG-OD1	5.61	123.34	118.30
1	A	353	ASP	CB-CG-OD1	5.55	123.29	118.30
1	A	43	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	A	687	ASP	CB-CG-OD1	5.52	123.27	118.30
1	A	385	ASP	CB-CG-OD2	-5.48	113.36	118.30
1	A	227	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	745	ASP	CB-CG-OD1	5.42	123.18	118.30
1	A	15	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	A	203	ARG	NE-CZ-NH1	-5.40	117.60	120.30
1	A	800	ASP	CB-CG-OD2	-5.39	113.45	118.30
1	A	27	LYS	CD-CE-NZ	-5.21	99.72	111.70
1	A	60	ARG	NE-CZ-NH1	-5.14	117.73	120.30
1	A	442	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	A	382	ILE	CB-CA-C	-5.11	101.39	111.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6379	0	6085	39	0
2	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	30	0	19	1	0
4	A	1	0	0	0	0
5	A	44	0	66	1	0
6	A	28	0	38	0	0
7	A	797	0	0	11	0
All	All	7280	0	6208	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:735:LYS:HD3	7:A:1498:HOH:O	1.92	0.70
1:A:43:ARG:NH2	1:A:135:TYR:OH	2.29	0.66
1:A:331:GLU:HB2	7:A:1508:HOH:O	1.99	0.63
1:A:170:ASN:HD21	1:A:431:ARG:HH11	1.47	0.61
1:A:27:LYS:HE2	7:A:950:HOH:O	2.05	0.54
1:A:435:PRO:HA	1:A:476:GLN:O	2.07	0.54
1:A:635:GLU:OE1	1:A:635:GLU:HA	2.08	0.54
1:A:637:ASN:HB3	5:A:829:EDO:H22	1.90	0.54
1:A:60:ARG:NH1	7:A:1336:HOH:O	2.37	0.54
1:A:368:GLU:OE1	7:A:1307:HOH:O	2.18	0.53
1:A:594:SER:HB2	1:A:597:LYS:HD2	1.92	0.51
1:A:273:GLU:OE1	7:A:1291:HOH:O	2.19	0.50
1:A:96:GLY:HA3	1:A:153:GLU:O	2.12	0.50
1:A:432:ILE:CD1	1:A:446:THR:HG21	2.42	0.50
1:A:78:ILE:O	1:A:82:ILE:HG13	2.13	0.49
1:A:756:ALA:HB2	1:A:781:ARG:CZ	2.43	0.49
1:A:775:ASP:HA	7:A:1136:HOH:O	2.12	0.48
1:A:144:ALA:HB1	1:A:145:PRO:HD2	1.95	0.48
1:A:142:HIS:CD2	1:A:155:GLY:HA2	2.49	0.48
1:A:217:TYR:HB2	1:A:300:LYS:HD3	1.96	0.47
1:A:65:TRP:HE1	1:A:304:CYS:HB3	1.79	0.47
1:A:65:TRP:NE1	1:A:304:CYS:HB3	2.31	0.46
1:A:527:PRO:HG2	7:A:1386:HOH:O	2.15	0.46
1:A:43:ARG:HD3	7:A:1487:HOH:O	2.15	0.46
1:A:646:SER:HB3	1:A:655:LEU:HD22	1.97	0.46
1:A:618:ARG:HD3	7:A:1233:HOH:O	2.16	0.45
3:A:827:THD:C2	3:A:827:THD:H4'2	2.29	0.45
1:A:326:SER:HB2	1:A:333:HIS:ND1	2.31	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:583:MET:HA	1:A:617:ALA:HB1	2.00	0.44
1:A:748:ARG:HA	1:A:753:ASP:OD2	2.18	0.43
1:A:657:ALA:HA	1:A:801:TRP:CZ2	2.54	0.43
1:A:554:GLN:HE22	1:A:738:GLY:HA3	1.84	0.43
1:A:635:GLU:HB3	1:A:636:ASN:ND2	2.34	0.42
1:A:602:PHE:N	1:A:602:PHE:CD2	2.88	0.42
1:A:28:TYR:CD2	1:A:76:ALA:HB2	2.55	0.41
1:A:60:ARG:HD3	1:A:60:ARG:HH11	1.64	0.41
1:A:170:ASN:ND2	1:A:431:ARG:HH11	2.17	0.41
1:A:635:GLU:HB2	1:A:639:GLU:OE2	2.20	0.41
1:A:435:PRO:O	1:A:436:ASP:HB3	2.22	0.40
1:A:85:HIS:HB3	7:A:857:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	800/845 (95%)	767 (96%)	32 (4%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	302	TRP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	677/712 (95%)	666 (98%)	11 (2%)	62 69

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	ARG
1	A	105	GLN
1	A	134	SER
1	A	147	THR
1	A	267	PHE
1	A	560	SER
1	A	605	LYS
1	A	635	GLU
1	A	639	GLU
1	A	684	GLU
1	A	754	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	132	GLN
1	A	170	ASN
1	A	408	GLN
1	A	441	ASN
1	A	472	GLN
1	A	554	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	EDO	A	838	-	3,3,3	0.76	0	2,2,2	0.64	0
5	EDO	A	833	-	3,3,3	0.56	0	2,2,2	0.94	0
5	EDO	A	839	-	3,3,3	0.36	0	2,2,2	0.72	0
5	EDO	A	837	-	3,3,3	0.11	0	2,2,2	1.27	0
5	EDO	A	830	-	3,3,3	0.53	0	2,2,2	0.39	0
5	EDO	A	829	-	3,3,3	0.65	0	2,2,2	0.21	0
5	EDO	A	831	-	3,3,3	0.56	0	2,2,2	0.37	0
5	EDO	A	834	-	3,3,3	0.37	0	2,2,2	0.98	0
6	2PE	A	840	-	27,27,27	1.34	1 (3%)	26,26,26	1.35	3 (11%)
5	EDO	A	835	-	3,3,3	0.47	0	2,2,2	0.90	0
3	THD	A	827	2	26,31,31	1.97	6 (23%)	30,46,46	2.96	8 (26%)
5	EDO	A	832	-	3,3,3	0.48	0	2,2,2	0.58	0
5	EDO	A	836	-	3,3,3	0.72	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	838	-	-	0/1/1/1	-
5	EDO	A	833	-	-	0/1/1/1	-
5	EDO	A	839	-	-	1/1/1/1	-
5	EDO	A	837	-	-	1/1/1/1	-
5	EDO	A	830	-	-	1/1/1/1	-
5	EDO	A	829	-	-	0/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	831	-	-	1/1/1/1	-
5	EDO	A	834	-	-	1/1/1/1	-
6	2PE	A	840	-	-	13/25/25/25	-
5	EDO	A	835	-	-	0/1/1/1	-
3	THD	A	827	2	-	5/16/23/23	0/2/2/2
5	EDO	A	832	-	-	0/1/1/1	-
5	EDO	A	836	-	-	0/1/1/1	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	827	THD	C7'-C5'	4.90	1.61	1.51
3	A	827	THD	C4'-N3'	3.87	1.40	1.35
3	A	827	THD	C9-C8	3.37	1.57	1.50
3	A	827	THD	O9-C8	3.32	1.41	1.32
3	A	827	THD	C6-C5	3.13	1.52	1.50
3	A	827	THD	C5-S1	-2.92	1.68	1.74
6	A	840	2PE	O7-C6	2.22	1.51	1.42

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	827	THD	C6-C5-C4	10.49	135.85	127.43
3	A	827	THD	C2-N3-C4	6.58	112.30	108.64
3	A	827	THD	C5-C4-N3	4.81	117.71	107.66
3	A	827	THD	C6'-N1'-C2'	4.68	123.93	115.96
3	A	827	THD	N1'-C2'-N3'	-3.30	119.85	125.54
6	A	840	2PE	C17-O16-C15	3.30	127.59	113.29
3	A	827	THD	CM2-C2'-N1'	2.92	120.35	117.14
6	A	840	2PE	O4-C5-C6	2.69	122.52	110.39
3	A	827	THD	CM4-C4-C5	-2.67	121.76	127.60
3	A	827	THD	PA-O3A-PB	-2.30	124.94	132.83
6	A	840	2PE	O10-C9-C8	2.04	119.58	110.39

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	827	THD	C4'-C5'-C7'-N3
6	A	840	2PE	O19-C20-C21-O22
6	A	840	2PE	O4-C5-C6-O7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
6	A	840	2PE	O22-C23-C24-O25
6	A	840	2PE	O25-C26-C27-O28
6	A	840	2PE	O7-C8-C9-O10
5	A	837	EDO	O1-C1-C2-O2
5	A	831	EDO	O1-C1-C2-O2
5	A	834	EDO	O1-C1-C2-O2
6	A	840	2PE	O13-C14-C15-O16
6	A	840	2PE	O1-C2-C3-O4
6	A	840	2PE	O10-C11-C12-O13
5	A	839	EDO	O1-C1-C2-O2
3	A	827	THD	PA-O3A-PB-O1B
6	A	840	2PE	C6-C5-O4-C3
6	A	840	2PE	C14-C15-O16-C17
6	A	840	2PE	C24-C23-O22-C21
3	A	827	THD	C6'-C5'-C7'-N3
6	A	840	2PE	C17-C18-O19-C20
6	A	840	2PE	C2-C3-O4-C5
5	A	830	EDO	O1-C1-C2-O2
3	A	827	THD	PA-O3A-PB-O2B
3	A	827	THD	PA-O3A-PB-O3B


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	829	EDO	1	0
3	A	827	THD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	802/845 (94%)	-0.56	2 (0%) 95 95	11, 19, 33, 50	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	507	VAL	2.6
1	A	508	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	2PE	A	840	28/28	0.64	0.21	36,50,59,61	0
5	EDO	A	836	4/4	0.73	0.23	51,52,56,57	0
5	EDO	A	831	4/4	0.79	0.20	51,53,53,54	0
5	EDO	A	832	4/4	0.80	0.27	47,48,48,51	0
5	EDO	A	830	4/4	0.84	0.17	40,40,43,45	0

*Continued on next page...*




*Continued from previous page...*


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	EDO	A	834	4/4	0.89	0.14	34,34,37,44	0
5	EDO	A	835	4/4	0.89	0.13	36,38,38,40	0
4	NA	A	828	1/1	0.93	0.10	18,18,18,18	1
5	EDO	A	833	4/4	0.94	0.10	32,33,33,36	0
5	EDO	A	829	4/4	0.94	0.14	33,36,39,42	0
5	EDO	A	837	4/4	0.94	0.19	38,39,40,45	0
2	MG	A	826	1/1	0.97	0.12	15,15,15,15	0
3	THD	A	827	30/30	0.98	0.14	12,17,25,31	4
5	EDO	A	839	4/4	0.98	0.07	26,29,29,30	0
5	EDO	A	838	4/4	0.98	0.06	17,17,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 2PE A 840:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.