



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 09:16 am BST

PDB ID : 6AHG
Title : Trimeric structure of concanavalin A from Canavalia ensiformis
Authors : Park, J.H.; Kim, D.S.; Park, Y.R.; Lee, S.J.
Deposited on : 2018-08-18
Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	<div> <div>20%</div> <div>92%</div> <div>6%</div> </div>
1	B	237	<div> <div>16%</div> <div>95%</div> <div></div> </div>
1	C	237	<div> <div>19%</div> <div>92%</div> <div>6%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5404 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Concanavalin-A,Concanavalin-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	235	Total	C	N	O	S	0	0	0
			1792	1131	298	361	2			
1	B	235	Total	C	N	O	S	0	0	0
			1792	1131	298	361	2			
1	C	235	Total	C	N	O	S	0	0	0
			1792	1131	298	361	2			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is CADMIUM ION (three-letter code: CD) (formula: Cd) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Cd	0	0
			2	2		
3	A	3	Total	Cd	0	0
			3	3		
3	C	1	Total	Cd	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total 6	O 6	0	0
4	B	7	Total 7	O 7	0	0
4	C	6	Total 6	O 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Concanavalin-A,Concanavalin-A



- Molecule 1: Concanavalin-A,Concanavalin-A



- Molecule 1: Concanavalin-A,Concanavalin-A



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	136.46Å 136.46Å 193.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.29 – 2.83 48.29 – 2.83	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.29-2.83) 99.8 (48.29-2.83)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.243 , 0.279 0.244 , 0.279	Depositor DCC
R_{free} test set	1141 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	50.6	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 6.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.477 for -1/2*h-1/2*k-1/2*l,-1/2*h-1/2*k+1/2*l,-h+k 0.459 for -1/2*h+1/2*k-1/2*l,1/2*h-1/2*k-1/2*l,-h-k	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	5404	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/1832	0.81	0/2494
1	B	0.47	0/1832	0.80	0/2494
1	C	0.48	0/1832	0.81	0/2494
All	All	0.49	0/5496	0.81	0/7482

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1792	0	1740	15	0
1	B	1792	0	1740	3	0
1	C	1792	0	1739	8	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	3	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
4	A	6	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	7	0	0	0	0
4	C	6	0	0	0	0
All	All	5404	0	5219	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:LEU:HD21	1:A:65:VAL:HG21	1.31	1.06
1:A:9:LEU:HD23	1:A:25:ILE:HG22	1.41	1.00
1:A:9:LEU:HD23	1:A:25:ILE:CG2	1.99	0.92
1:A:183:GLU:HG2	1:A:184:SER:H	1.41	0.85
1:A:9:LEU:HD21	1:A:65:VAL:CG2	2.19	0.66
1:A:181:ILE:O	1:A:182:TRP:HB2	1.99	0.61
1:A:9:LEU:CD2	1:A:65:VAL:HG21	2.21	0.59
1:A:183:GLU:HG2	1:A:184:SER:N	2.16	0.57
1:C:100:TYR:HB2	1:C:207:ALA:HB2	1.89	0.55
1:A:183:GLU:CG	1:A:184:SER:H	2.15	0.54
1:B:100:TYR:HB2	1:B:207:ALA:HB2	1.91	0.52
1:B:135:LYS:HG2	1:B:149:GLY:HA3	1.93	0.51
1:A:100:TYR:HB2	1:A:207:ALA:HB2	1.95	0.47
1:C:17:ILE:HG12	1:C:228:ARG:HD2	1.96	0.47
1:A:83:ASN:HD22	1:B:33:ARG:HH22	1.64	0.46
1:C:135:LYS:HG2	1:C:149:GLY:HA3	1.99	0.45
1:A:9:LEU:CD2	1:A:25:ILE:CG2	2.84	0.43
1:A:9:LEU:HB2	1:A:210:ILE:O	2.19	0.43
1:C:53:ILE:HG22	1:C:192:GLU:HG2	2.01	0.42
1:A:12:TYR:HB2	1:A:207:ALA:HA	2.02	0.42
1:C:44:ASN:HD21	1:C:201:SER:HB3	1.84	0.41
1:C:104:ASN:HD22	1:C:210:ILE:HG13	1.85	0.41
1:A:33:ARG:HH22	1:C:83:ASN:HD22	1.69	0.41
1:C:12:TYR:HB2	1:C:207:ALA:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/237 (98%)	225 (97%)	6 (3%)	0	100	100
1	B	231/237 (98%)	225 (97%)	5 (2%)	1 (0%)	34	56
1	C	231/237 (98%)	224 (97%)	7 (3%)	0	100	100
All	All	693/711 (98%)	674 (97%)	18 (3%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	208	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/203 (99%)	197 (98%)	4 (2%)	55	76
1	B	201/203 (99%)	198 (98%)	3 (2%)	65	82
1	C	201/203 (99%)	198 (98%)	3 (2%)	65	82
All	All	603/609 (99%)	593 (98%)	10 (2%)	60	80

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	ASP
1	A	184	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	203	ASP
1	A	235	ASP
1	B	107	LEU
1	B	183	GLU
1	B	188	VAL
1	C	53	ILE
1	C	58	ASP
1	C	188	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	104	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	235/237 (99%)	1.10	48 (20%) 1 0	28, 47, 74, 93	0
1	B	235/237 (99%)	1.09	39 (16%) 1 1	34, 47, 73, 97	0
1	C	235/237 (99%)	1.16	46 (19%) 1 0	32, 47, 73, 93	0
All	All	705/711 (99%)	1.12	133 (18%) 1 1	28, 47, 73, 97	0

All (133) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	136	ASP	10.3
1	C	185	SER	8.2
1	B	185	SER	7.9
1	B	184	SER	7.5
1	C	216	ASN	7.3
1	A	1	ALA	7.1
1	B	99	LEU	6.9
1	C	119	SER	6.3
1	B	204	SER	6.3
1	B	162	ASN	6.2
1	C	215	SER	6.0
1	C	208	ASP	5.9
1	B	69	ASN	5.9
1	A	224	GLY	5.9
1	C	134	SER	5.5
1	A	208	ASP	5.4
1	B	31	SER	5.2
1	C	51	HIS	5.0
1	C	100	TYR	4.7
1	B	203	ASP	4.6
1	A	14	ASN	4.6
1	C	182	TRP	4.5
1	C	148	THR	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	100	TYR	4.4
1	C	186	ALA	4.3
1	C	204	SER	4.3
1	C	123	THR	4.3
1	A	118	ASN	4.3
1	C	168	SER	4.3
1	B	135	LYS	4.3
1	B	208	ASP	4.2
1	B	235	ASP	4.2
1	C	99	LEU	4.2
1	A	227	GLY	4.1
1	B	136	ASP	4.0
1	B	237	ASN	4.0
1	B	100	TYR	3.9
1	A	230	LEU	3.9
1	A	167	GLY	3.9
1	B	149	GLY	3.8
1	A	69	ASN	3.8
1	B	205	HIS	3.8
1	C	150	THR	3.8
1	A	74	THR	3.7
1	A	148	THR	3.6
1	B	188	VAL	3.6
1	A	184	SER	3.5
1	C	169	SER	3.4
1	C	147	THR	3.4
1	A	73	ALA	3.4
1	B	3	THR	3.3
1	C	132	GLN	3.3
1	B	171	GLY	3.2
1	A	12	TYR	3.2
1	C	23	PRO	3.2
1	B	151	ASP	3.1
1	B	150	THR	3.1
1	C	97	THR	3.1
1	A	185	SER	3.1
1	C	41	ASN	3.1
1	A	116	LYS	3.1
1	C	191	PHE	3.0
1	B	1	ALA	3.0
1	A	80	ASP	3.0
1	B	98	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	106	ILE	2.9
1	A	236	ALA	2.9
1	A	228	ARG	2.9
1	C	135	LYS	2.9
1	A	81	LEU	2.9
1	B	192	GLU	2.9
1	C	226	THR	2.9
1	C	164	SER	2.8
1	B	189	ALA	2.8
1	C	224	GLY	2.8
1	A	51	HIS	2.8
1	C	184	SER	2.7
1	A	226	THR	2.7
1	C	183	GLU	2.7
1	C	151	ASP	2.7
1	A	205	HIS	2.7
1	A	210	ILE	2.7
1	C	146	ALA	2.7
1	C	217	ILE	2.7
1	A	207	ALA	2.7
1	C	214	ILE	2.6
1	B	81	LEU	2.6
1	C	96	SER	2.6
1	C	231	GLY	2.6
1	A	162	ASN	2.6
1	A	209	GLY	2.5
1	A	150	THR	2.5
1	A	119	SER	2.5
1	B	198	LEU	2.5
1	C	227	GLY	2.5
1	A	102	GLU	2.4
1	C	145	ASP	2.4
1	B	148	THR	2.4
1	A	71	ASP	2.4
1	C	76	SER	2.4
1	B	58	ASP	2.4
1	A	4	ILE	2.3
1	B	229	LEU	2.3
1	A	204	SER	2.3
1	C	209	GLY	2.3
1	C	12	TYR	2.3
1	A	223	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	79	VAL	2.3
1	B	105	THR	2.3
1	A	200	LYS	2.2
1	B	220	SER	2.2
1	C	225	SER	2.2
1	A	206	PRO	2.2
1	A	7	VAL	2.2
1	B	154	LEU	2.2
1	C	1	ALA	2.2
1	B	186	ALA	2.2
1	C	52	ILE	2.1
1	A	212	PHE	2.1
1	C	112	THR	2.1
1	B	224	GLY	2.1
1	A	84	VAL	2.1
1	A	151	ASP	2.1
1	A	149	GLY	2.1
1	A	68	PRO	2.1
1	A	96	SER	2.0
1	A	214	ILE	2.0
1	B	94	SER	2.0
1	B	163	GLY	2.0
1	C	207	ALA	2.0
1	B	183	GLU	2.0
1	A	104	ASN	2.0
1	A	232	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CA	C	301	1/1	0.94	0.30	18,18,18,18	0
2	CA	B	301	1/1	0.95	0.33	16,16,16,16	0
2	CA	A	301	1/1	0.96	0.26	17,17,17,17	0
3	CD	B	302	1/1	0.98	0.19	41,41,41,41	0
3	CD	A	304	1/1	0.99	0.15	54,54,54,54	0
3	CD	C	302	1/1	0.99	0.16	42,42,42,42	0
3	CD	A	302	1/1	0.99	0.13	39,39,39,39	0
3	CD	A	303	1/1	0.99	0.12	53,53,53,53	0
3	CD	B	303	1/1	0.99	0.13	52,52,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CA B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CA A 301:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CD B 302:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around CD C 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CD A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CD A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CD B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.