



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 21, 2020 – 05:20 PM BST

PDB ID : 3AIC
Title : Crystal Structure of Glucansucrase from Streptococcus mutans
Authors : Ito, K.; Ito, S.; Shimamura, T.; Iwata, S.
Deposited on : 2010-05-12
Resolution : 3.11 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1292 (3.14-3.10)
Clashscore	141614	1389 (3.14-3.10)
Ramachandran outliers	138981	1337 (3.14-3.10)
Sidechain outliers	138945	1337 (3.14-3.10)
RSRZ outliers	127900	1260 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	844	<div> <div>%</div> <div> <div></div> <div>65%</div> <div>31%</div> <div>•</div> </div> </div>
1	B	844	<div> <div>2%</div> <div> <div></div> <div>57%</div> <div>29%</div> <div>•</div> <div>11%</div> </div> </div>
1	C	844	<div> <div></div> <div> <div></div> <div>66%</div> <div>31%</div> <div>•</div> </div> </div>
1	D	844	<div> <div>%</div> <div> <div></div> <div>63%</div> <div>32%</div> <div>5%</div> </div> </div>
1	E	844	<div> <div>2%</div> <div> <div></div> <div>66%</div> <div>28%</div> <div>5%</div> <div>•</div> </div> </div>
1	F	844	<div> <div>%</div> <div> <div></div> <div>62%</div> <div>34%</div> <div>•</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	844	<div> <div></div> <div>64%</div> <div>31%</div> <div>.</div> </div>
1	H	844	<div> <div></div> <div>64%</div> <div>27%</div> <div>...</div> </div>
2	I	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>
2	J	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>
2	K	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>
2	L	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>
2	M	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>
2	N	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>
2	O	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>
2	P	3	<div> <div></div> <div>33%</div> <div>67%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	AC1	I	3	-	-	X	-
2	AC1	K	3	-	-	X	-
2	AC1	M	3	-	-	X	-
4	MES	F	5001	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 52865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucosyltransferase-SI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	842	Total	C	N	O	S	0	0	0
			6643	4184	1141	1302	16			
1	B	747	Total	C	N	O	S	0	0	0
			5854	3684	1005	1151	14			
1	C	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	D	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	E	843	Total	C	N	O	S	0	0	0
			6654	4193	1142	1303	16			
1	F	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	G	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	H	807	Total	C	N	O	S	0	0	0
			6372	4019	1093	1244	16			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	597	ASP	ASN	SEE REMARK 999	UNP P13470
A	600	LYS	ARG	SEE REMARK 999	UNP P13470
A	727	ILE	THR	SEE REMARK 999	UNP P13470
A	734	VAL	ALA	SEE REMARK 999	UNP P13470
B	597	ASP	ASN	SEE REMARK 999	UNP P13470
B	600	LYS	ARG	SEE REMARK 999	UNP P13470
B	727	ILE	THR	SEE REMARK 999	UNP P13470
B	734	VAL	ALA	SEE REMARK 999	UNP P13470
C	597	ASP	ASN	SEE REMARK 999	UNP P13470
C	600	LYS	ARG	SEE REMARK 999	UNP P13470
C	727	ILE	THR	SEE REMARK 999	UNP P13470
C	734	VAL	ALA	SEE REMARK 999	UNP P13470
D	597	ASP	ASN	SEE REMARK 999	UNP P13470

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	600	LYS	ARG	SEE REMARK 999	UNP P13470
D	727	ILE	THR	SEE REMARK 999	UNP P13470
D	734	VAL	ALA	SEE REMARK 999	UNP P13470
E	597	ASP	ASN	SEE REMARK 999	UNP P13470
E	600	LYS	ARG	SEE REMARK 999	UNP P13470
E	727	ILE	THR	SEE REMARK 999	UNP P13470
E	734	VAL	ALA	SEE REMARK 999	UNP P13470
F	597	ASP	ASN	SEE REMARK 999	UNP P13470
F	600	LYS	ARG	SEE REMARK 999	UNP P13470
F	727	ILE	THR	SEE REMARK 999	UNP P13470
F	734	VAL	ALA	SEE REMARK 999	UNP P13470
G	597	ASP	ASN	SEE REMARK 999	UNP P13470
G	600	LYS	ARG	SEE REMARK 999	UNP P13470
G	727	ILE	THR	SEE REMARK 999	UNP P13470
G	734	VAL	ALA	SEE REMARK 999	UNP P13470
H	597	ASP	ASN	SEE REMARK 999	UNP P13470
H	600	LYS	ARG	SEE REMARK 999	UNP P13470
H	727	ILE	THR	SEE REMARK 999	UNP P13470
H	734	VAL	ALA	SEE REMARK 999	UNP P13470


- Molecule 2 is an oligosaccharide called 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	J	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	K	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	L	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	M	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	N	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	O	3	Total	C	N	O	0	0	0
			44	25	1	18			
2	P	3	Total	C	N	O	0	0	0
			44	25	1	18			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	1	Total Ca 1 1	0	0
3	D	1	Total Ca 1 1	0	0
3	E	1	Total Ca 1 1	0	0
3	H	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0
3	C	1	Total Ca 1 1	0	0
3	A	1	Total Ca 1 1	0	0
3	F	1	Total Ca 1 1	0	0

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O S 12 6 1 4 1	0	0
4	B	1	Total C N O S 12 6 1 4 1	0	0
4	C	1	Total C N O S 12 6 1 4 1	0	0


Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	D	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	E	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	F	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	G	1	Total 12	C 6	N 1	O 4	S 1	0	0
4	H	1	Total 12	C 6	N 1	O 4	S 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	33	Total 33	O 33	0	0
5	B	20	Total 20	O 20	0	0
5	C	32	Total 32	O 32	0	0
5	D	40	Total 40	O 40	0	0
5	E	48	Total 48	O 48	0	0
5	F	15	Total 15	O 15	0	0
5	G	34	Total 34	O 34	0	0
5	H	24	Total 24	O 24	0	0






Molecule 1: Glucosyltransferase-SI







• Molecule 1: Glucosyltransferase-SI



• Molecule 1: Glucosyltransferase-SI





- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl}]\text{amino}\}$ - α -D-glucopyranose-(1-4)- α -D-glucopyranose-(1-4)- α -D-glucopyranose

Chain I: 33% 67%



- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl}]\text{amino}\}$ - α -D-glucopyranose-(1-4)- α -D-glucopyranose-(1-4)- α -D-glucopyranose

Chain J: 33% 67%



- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl}]\text{amino}\}$ - α -D-glucopyranose-(1-4)- α -D-glucopyranose-(1-4)- α -D-glucopyranose

Chain K: 33% 67%



- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl}]\text{amino}\}$ - α -D-glucopyranose-(1-4)- α -D-glucopyranose-(1-4)- α -D-glucopyranose

Chain L: 33% 67%



- Molecule 2: 4,6-dideoxy-4- $\{[(1S,4R,5S,6S)-4,5,6\text{-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl}]\text{amino}\}$ - α -D-glucopyranose-(1-4)- α -D-glucopyranose-(1-4)- α -D-glucopyranose

Chain M: 33% 67%



- Molecule 2: 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	295.52Å 214.41Å 220.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.77 – 3.11 61.77 – 3.11	Depositor EDS
% Data completeness (in resolution range)	100.0 (61.77-3.11) 95.3 (61.77-3.11)	Depositor EDS
R_{merge}	0.37	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 3.13Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.211 , 0.244 0.208 , 0.239	Depositor DCC
R_{free} test set	12026 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	65.5	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	52865	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.08 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.9554e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC, AC1, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.99	9/6784 (0.1%)	0.95	3/9213 (0.0%)
1	B	0.88	12/5976 (0.2%)	0.88	7/8113 (0.1%)
1	C	0.94	1/6802 (0.0%)	0.94	8/9237 (0.1%)
1	D	0.95	4/6802 (0.1%)	0.92	11/9237 (0.1%)
1	E	1.04	14/6796 (0.2%)	0.95	10/9229 (0.1%)
1	F	0.81	1/6802 (0.0%)	0.85	3/9237 (0.0%)
1	G	0.97	6/6802 (0.1%)	0.95	7/9237 (0.1%)
1	H	0.91	4/6510 (0.1%)	0.93	9/8837 (0.1%)
All	All	0.94	51/53274 (0.1%)	0.92	58/72340 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	H	0	1
All	All	0	2

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	320	HIS	CE1-NE2	14.48	1.66	1.32
1	E	245	PHE	N-CA	12.06	1.70	1.46
1	E	245	PHE	CE2-CZ	11.00	1.58	1.37
1	E	273	ARG	CZ-NH1	10.60	1.46	1.33
1	H	249	ASN	CG-OD1	10.12	1.46	1.24

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	245	PHE	CB-CG-CD1	-11.95	112.44	120.80
1	H	245	PHE	CG-CD2-CE2	-10.93	108.77	120.80
1	E	273	ARG	NE-CZ-NH2	-9.82	115.39	120.30
1	D	793	ASP	CB-CA-C	-8.51	93.38	110.40
1	E	727	ILE	CG1-CB-CG2	-8.07	93.64	111.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	370	TRP	Peptide
1	H	245	PHE	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6643	0	6475	332	0
1	B	5854	0	5691	243	0
1	C	6660	0	6489	258	0
1	D	6660	0	6489	323	0
1	E	6654	0	6484	337	0
1	F	6660	0	6489	318	0
1	G	6660	0	6489	287	0
1	H	6372	0	6196	334	0
2	I	44	0	30	10	0
2	J	44	0	30	7	0
2	K	44	0	30	9	0
2	L	44	0	30	6	0
2	M	44	0	30	10	0
2	N	44	0	30	6	0
2	O	44	0	30	7	0
2	P	44	0	30	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	A	12	0	12	2	0
4	B	12	0	12	0	0
4	C	12	0	12	2	0
4	D	12	0	12	2	0
4	E	12	0	12	1	0
4	F	12	0	12	0	0
4	G	12	0	12	1	0
4	H	12	0	12	0	0
5	A	33	0	0	6	0
5	B	20	0	0	6	0
5	C	32	0	0	2	0
5	D	40	0	0	2	0
5	E	48	0	0	2	0
5	F	15	0	0	10	0
5	G	34	0	0	2	0
5	H	24	0	0	4	0
All	All	52865	0	51138	2464	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 2464 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:245:PHE:N	1:E:245:PHE:CA	1.70	1.54
1:E:277:ILE:CD1	1:E:291:ASP:HB3	1.38	1.54
1:A:290:LYS:NZ	1:A:290:LYS:CE	1.71	1.53
1:E:279:LYS:CE	1:E:279:LYS:NZ	1.73	1.51
1:E:273:ARG:CD	1:E:287:SER:HB2	1.40	1.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	840/844 (100%)	772 (92%)	57 (7%)	11 (1%)	12	41
1	B	743/844 (88%)	678 (91%)	53 (7%)	12 (2%)	9	36
1	C	842/844 (100%)	777 (92%)	57 (7%)	8 (1%)	15	48
1	D	842/844 (100%)	763 (91%)	66 (8%)	13 (2%)	10	38
1	E	841/844 (100%)	758 (90%)	70 (8%)	13 (2%)	10	38
1	F	842/844 (100%)	771 (92%)	65 (8%)	6 (1%)	22	56
1	G	842/844 (100%)	776 (92%)	56 (7%)	10 (1%)	13	43
1	H	803/844 (95%)	717 (89%)	62 (8%)	24 (3%)	4	23
All	All	6595/6752 (98%)	6012 (91%)	486 (7%)	97 (2%)	10	38

5 of 97 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	250	GLN
1	C	321	GLN
1	C	327	THR
1	C	351	LYS
1	D	278	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	713/715 (100%)	669 (94%)	44 (6%)	18	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	627/715 (88%)	596 (95%)	31 (5%)	25	57
1	C	715/715 (100%)	671 (94%)	44 (6%)	18	48
1	D	715/715 (100%)	671 (94%)	44 (6%)	18	48
1	E	714/715 (100%)	658 (92%)	56 (8%)	12	40
1	F	715/715 (100%)	662 (93%)	53 (7%)	13	41
1	G	715/715 (100%)	666 (93%)	49 (7%)	15	44
1	H	682/715 (95%)	640 (94%)	42 (6%)	18	48
All	All	5596/5720 (98%)	5233 (94%)	363 (6%)	17	47

5 of 363 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	247	GLN
1	E	740	ASN
1	H	368	SER
1	E	286	GLN
1	E	332	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 178 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	700	GLN
1	E	440	ASN
1	H	407	ASN
1	D	759	HIS
1	E	258	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

24 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	I	1	2	12,12,12	0.69	0	17,17,17	2.03	6 (35%)
2	GLC	I	2	2	11,11,12	1.19	0	15,15,17	2.72	5 (33%)
2	AC1	I	3	2	21,22,23	1.48	5 (23%)	22,32,34	2.28	4 (18%)
2	GLC	J	1	2	12,12,12	0.79	0	17,17,17	1.92	5 (29%)
2	GLC	J	2	2	11,11,12	1.15	1 (9%)	15,15,17	2.33	5 (33%)
2	AC1	J	3	2	21,22,23	1.35	2 (9%)	22,32,34	1.97	4 (18%)
2	GLC	K	1	2	12,12,12	0.79	0	17,17,17	1.92	5 (29%)
2	GLC	K	2	2	11,11,12	1.14	1 (9%)	15,15,17	2.34	5 (33%)
2	AC1	K	3	2	21,22,23	1.34	2 (9%)	22,32,34	1.98	4 (18%)
2	GLC	L	1	2	12,12,12	0.80	0	17,17,17	1.92	5 (29%)
2	GLC	L	2	2	11,11,12	1.14	1 (9%)	15,15,17	2.33	5 (33%)
2	AC1	L	3	2	21,22,23	1.35	2 (9%)	22,32,34	1.98	4 (18%)
2	GLC	M	1	2	12,12,12	0.79	0	17,17,17	1.92	5 (29%)
2	GLC	M	2	2	11,11,12	1.15	1 (9%)	15,15,17	2.33	5 (33%)
2	AC1	M	3	2	21,22,23	1.36	3 (14%)	22,32,34	1.97	4 (18%)
2	GLC	N	1	2	12,12,12	0.79	0	17,17,17	1.92	5 (29%)
2	GLC	N	2	2	11,11,12	1.15	1 (9%)	15,15,17	2.33	5 (33%)
2	AC1	N	3	2	21,22,23	1.34	2 (9%)	22,32,34	1.98	4 (18%)
2	GLC	O	1	2	12,12,12	0.79	0	17,17,17	1.92	5 (29%)
2	GLC	O	2	2	11,11,12	1.13	1 (9%)	15,15,17	2.34	5 (33%)
2	AC1	O	3	2	21,22,23	1.35	2 (9%)	22,32,34	1.97	4 (18%)
2	GLC	P	1	2	12,12,12	0.79	0	17,17,17	1.92	5 (29%)
2	GLC	P	2	2	11,11,12	1.14	1 (9%)	15,15,17	2.33	5 (33%)
2	AC1	P	3	2	21,22,23	1.35	2 (9%)	22,32,34	1.97	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	I	1	2	-	2/2/22/22	0/1/1/1
2	GLC	I	2	2	-	2/2/19/22	0/1/1/1
2	AC1	I	3	2	-	3/6/43/46	0/2/2/2
2	GLC	J	1	2	-	2/2/22/22	0/1/1/1
2	GLC	J	2	2	-	2/2/19/22	0/1/1/1
2	AC1	J	3	2	-	3/6/43/46	0/2/2/2
2	GLC	K	1	2	-	2/2/22/22	0/1/1/1
2	GLC	K	2	2	-	2/2/19/22	0/1/1/1
2	AC1	K	3	2	-	3/6/43/46	0/2/2/2
2	GLC	L	1	2	-	2/2/22/22	0/1/1/1
2	GLC	L	2	2	-	2/2/19/22	0/1/1/1
2	AC1	L	3	2	-	3/6/43/46	0/2/2/2
2	GLC	M	1	2	-	2/2/22/22	0/1/1/1
2	GLC	M	2	2	-	2/2/19/22	0/1/1/1
2	AC1	M	3	2	-	3/6/43/46	0/2/2/2
2	GLC	N	1	2	-	2/2/22/22	0/1/1/1
2	GLC	N	2	2	-	2/2/19/22	0/1/1/1
2	AC1	N	3	2	-	3/6/43/46	0/2/2/2
2	GLC	O	1	2	-	2/2/22/22	0/1/1/1
2	GLC	O	2	2	-	2/2/19/22	0/1/1/1
2	AC1	O	3	2	-	3/6/43/46	0/2/2/2
2	GLC	P	1	2	-	2/2/22/22	0/1/1/1
2	GLC	P	2	2	-	2/2/19/22	0/1/1/1
2	AC1	P	3	2	-	3/6/43/46	0/2/2/2

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	3	AC1	C4A-C5B	-3.16	1.48	1.51
2	L	3	AC1	C4A-C5B	-2.73	1.49	1.51
2	J	3	AC1	C4A-C5B	-2.71	1.49	1.51
2	M	3	AC1	C4A-C5B	-2.70	1.49	1.51
2	P	3	AC1	C4A-C5B	-2.70	1.49	1.51

The worst 5 of 113 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	3	AC1	C1-C2-C3	8.17	119.71	109.67
2	L	3	AC1	C2-C3-C4	-6.23	105.13	110.63
2	K	3	AC1	C2-C3-C4	-6.20	105.16	110.63
2	N	3	AC1	C2-C3-C4	-6.20	105.17	110.63
2	P	3	AC1	C2-C3-C4	-6.18	105.18	110.63

There are no chirality outliers.

5 of 56 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	K	3	AC1	C7B-C1B-N4A-C4
2	K	3	AC1	C4A-C5B-C6B-O6B
2	K	3	AC1	C7B-C5B-C6B-O6B
2	L	3	AC1	C7B-C1B-N4A-C4
2	L	3	AC1	C4A-C5B-C6B-O6B


There are no ring outliers.


16 monomers are involved in 61 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	3	AC1	9	0
2	L	3	AC1	6	0
2	N	2	GLC	4	0
2	J	2	GLC	4	0
2	I	3	AC1	10	0
2	N	3	AC1	6	0
2	O	2	GLC	4	0
2	I	2	GLC	4	0
2	O	3	AC1	7	0
2	M	2	GLC	4	0
2	K	2	GLC	4	0
2	L	2	GLC	4	0
2	M	3	AC1	10	0
2	P	3	AC1	6	0
2	J	3	AC1	7	0
2	P	2	GLC	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




















5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MES	A	5001	-	12,12,12	1.49	2 (16%)	14,16,16	8.44	10 (71%)
4	MES	B	5001	-	12,12,12	1.68	4 (33%)	14,16,16	8.11	11 (78%)
4	MES	D	5001	-	12,12,12	1.39	2 (16%)	14,16,16	6.50	9 (64%)
4	MES	F	5001	-	12,12,12	1.84	3 (25%)	14,16,16	7.65	11 (78%)
4	MES	C	5001	-	12,12,12	1.49	3 (25%)	14,16,16	6.95	9 (64%)
4	MES	E	5001	-	12,12,12	1.38	4 (33%)	14,16,16	7.57	11 (78%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MES	G	5001	-	12,12,12	1.44	2 (16%)	14,16,16	7.31	10 (71%)
4	MES	H	5001	-	12,12,12	1.05	0	14,16,16	7.36	11 (78%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	A	5001	-	-	2/6/14/14	0/1/1/1
4	MES	B	5001	-	-	2/6/14/14	0/1/1/1
4	MES	D	5001	-	-	5/6/14/14	0/1/1/1
4	MES	F	5001	-	-	4/6/14/14	0/1/1/1
4	MES	C	5001	-	-	4/6/14/14	0/1/1/1
4	MES	E	5001	-	-	1/6/14/14	0/1/1/1
4	MES	G	5001	-	-	5/6/14/14	0/1/1/1
4	MES	H	5001	-	-	1/6/14/14	0/1/1/1

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	5001	MES	C8-S	-4.00	1.71	1.77
4	A	5001	MES	O2S-S	3.67	1.55	1.45
4	B	5001	MES	C8-S	-3.27	1.72	1.77
4	B	5001	MES	O2S-S	3.04	1.54	1.45
4	F	5001	MES	O1S-S	2.81	1.53	1.45

The worst 5 of 82 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	5001	MES	O2S-S-C8	-26.11	75.47	106.92
4	B	5001	MES	O1S-S-C8	-21.78	80.69	106.92
4	F	5001	MES	O2S-S-C8	-18.51	84.62	106.92
4	C	5001	MES	O1S-S-C8	-18.46	84.69	106.92
4	E	5001	MES	O2S-S-C8	-18.26	84.93	106.92

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	5001	MES	C8-C7-N4-C5
4	B	5001	MES	N4-C7-C8-S
4	D	5001	MES	C8-C7-N4-C5
4	D	5001	MES	N4-C7-C8-S
4	F	5001	MES	C8-C7-N4-C5

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	5001	MES	2	0
4	D	5001	MES	2	0
4	C	5001	MES	2	0
4	E	5001	MES	1	0
4	G	5001	MES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	842/844 (99%)	-0.19	7 (0%) 86 74	23, 38, 90, 113	0
1	B	747/844 (88%)	-0.11	15 (2%) 65 45	31, 50, 88, 136	0
1	C	844/844 (100%)	-0.31	2 (0%) 95 91	24, 35, 60, 70	0
1	D	844/844 (100%)	-0.23	11 (1%) 77 60	26, 39, 83, 101	0
1	E	843/844 (99%)	-0.18	16 (1%) 66 47	22, 36, 101, 126	0
1	F	844/844 (100%)	-0.17	6 (0%) 87 77	37, 53, 73, 89	0
1	G	844/844 (100%)	-0.26	6 (0%) 87 77	24, 35, 68, 79	0
1	H	807/844 (95%)	-0.19	13 (1%) 72 52	24, 42, 95, 139	0
All	All	6615/6752 (97%)	-0.21	76 (1%) 80 65	22, 41, 83, 139	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1087	PRO	5.8
1	A	1087	PRO	5.3
1	G	1087	PRO	5.0
1	E	285	THR	4.7
1	H	1087	PRO	4.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	L	1	12/12	0.80	0.28	48,49,49,50	0
2	GLC	M	1	12/12	0.81	0.30	48,49,49,50	0
2	GLC	O	1	12/12	0.83	0.23	48,49,49,50	0
2	GLC	N	1	12/12	0.83	0.26	48,49,49,50	0
2	GLC	N	2	11/12	0.84	0.28	46,47,47,47	0
2	GLC	O	2	11/12	0.86	0.26	46,47,47,47	0
2	GLC	K	1	12/12	0.86	0.25	48,49,49,50	0
2	GLC	P	1	12/12	0.86	0.26	48,49,49,50	0
2	GLC	M	2	11/12	0.87	0.27	46,47,47,47	0
2	AC1	M	3	21/22	0.89	0.29	35,50,53,53	0
2	GLC	J	1	12/12	0.89	0.21	48,49,49,50	0
2	GLC	J	2	11/12	0.89	0.22	46,47,47,47	0
2	GLC	I	1	12/12	0.90	0.21	48,49,49,50	0
2	AC1	N	3	21/22	0.90	0.28	35,50,53,53	0
2	GLC	L	2	11/12	0.90	0.23	46,47,47,47	0
2	AC1	K	3	21/22	0.91	0.27	35,50,53,53	0
2	GLC	K	2	11/12	0.91	0.27	46,47,47,47	0
2	GLC	P	2	11/12	0.91	0.22	46,47,47,47	0
2	GLC	I	2	11/12	0.91	0.20	46,47,47,47	0
2	AC1	P	3	21/22	0.92	0.22	35,50,53,53	0
2	AC1	I	3	21/22	0.92	0.24	35,50,53,53	0
2	AC1	L	3	21/22	0.94	0.24	35,50,53,53	0
2	AC1	J	3	21/22	0.94	0.23	35,50,53,53	0
2	AC1	O	3	21/22	0.95	0.26	35,50,53,53	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain I:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain J:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain L:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain M:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around Chain N:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain O:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain P:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CA	F	4001	1/1	0.94	0.15	38,38,38,38	0
4	MES	B	5001	12/12	0.94	0.18	39,41,43,44	0
4	MES	F	5001	12/12	0.94	0.17	36,37,37,37	0
4	MES	A	5001	12/12	0.94	0.20	29,31,33,33	0
4	MES	C	5001	12/12	0.95	0.16	29,31,32,32	0
4	MES	E	5001	12/12	0.96	0.18	30,33,35,36	0
4	MES	D	5001	12/12	0.96	0.16	31,33,35,35	0
3	CA	A	4001	1/1	0.96	0.16	28,28,28,28	0
4	MES	G	5001	12/12	0.96	0.15	30,31,33,33	0
4	MES	H	5001	12/12	0.97	0.17	28,30,32,32	0
3	CA	B	4001	1/1	0.97	0.14	33,33,33,33	0
3	CA	C	4001	1/1	0.98	0.13	29,29,29,29	0
3	CA	E	4001	1/1	0.98	0.18	32,32,32,32	0
3	CA	D	4001	1/1	0.98	0.16	32,32,32,32	0
3	CA	G	4001	1/1	0.98	0.14	28,28,28,28	0
3	CA	H	4001	1/1	0.99	0.17	32,32,32,32	0

6.5 Other polymers

There are no such residues in this entry.