



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:58 am BST

PDB ID : 2AJP
Title : Crystal structure of a human pyridoxal kinase
Authors : Ismail, S.; Dimov, S.; Atanassova, A.; Tempel, W.; Arrowsmith, C.; Edwards, A.; Sundstrom, M.; Weigelt, J.; Bochkarev, A.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2005-08-02
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	<div> <div>5%</div> <div> <div></div> <div>81%</div> <div>12%</div> <div>7%</div> </div> </div>
1	B	326	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>11%</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNX	A	1404	-	-	-	X
3	UNX	A	1409	-	-	-	X
3	UNX	A	1411	-	-	-	X
3	UNX	A	1412	-	-	-	X
3	UNX	A	402	-	-	-	X
3	UNX	A	403	-	-	-	X
3	UNX	A	405	-	-	-	X
3	UNX	A	406	-	-	-	X
3	UNX	A	407	-	-	-	X
3	UNX	A	408	-	-	-	X
3	UNX	A	410	-	-	-	X
3	UNX	A	413	-	-	-	X
3	UNX	A	414	-	-	-	X
3	UNX	A	415	-	-	-	X
3	UNX	A	416	-	-	-	X
3	UNX	A	417	-	-	-	X
3	UNX	A	418	-	-	-	X
3	UNX	A	419	-	-	-	X
3	UNX	A	420	-	-	-	X
3	UNX	A	421	-	-	-	X
3	UNX	B	1402	-	-	-	X
3	UNX	B	1403	-	-	-	X
3	UNX	B	1405	-	-	-	X
3	UNX	B	1406	-	-	-	X
3	UNX	B	1407	-	-	-	X
3	UNX	B	1408	-	-	-	X
3	UNX	B	1410	-	-	-	X
3	UNX	B	1413	-	-	-	X
3	UNX	B	1414	-	-	-	X
3	UNX	B	1415	-	-	-	X
3	UNX	B	1416	-	-	-	X
3	UNX	B	1417	-	-	-	X
3	UNX	B	1418	-	-	-	X
3	UNX	B	1419	-	-	-	X
3	UNX	B	1420	-	-	-	X
3	UNX	B	1421	-	-	-	X
3	UNX	B	404	-	-	-	X
3	UNX	B	409	-	-	-	X
3	UNX	B	411	-	-	-	X
3	UNX	B	412	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4693 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridoxal kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	303	Total	C	N	O	S	0	0	0
			2313	1469	392	438	14			
1	B	301	Total	C	N	O	S	0	0	0
			2276	1451	382	430	13			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	CLONING ARTIFACT	UNP O00764
A	-12	GLY	-	CLONING ARTIFACT	UNP O00764
A	-11	SER	-	CLONING ARTIFACT	UNP O00764
A	-10	SER	-	CLONING ARTIFACT	UNP O00764
A	-9	HIS	-	EXPRESSION TAG	UNP O00764
A	-8	HIS	-	EXPRESSION TAG	UNP O00764
A	-7	HIS	-	EXPRESSION TAG	UNP O00764
A	-6	HIS	-	EXPRESSION TAG	UNP O00764
A	-5	HIS	-	EXPRESSION TAG	UNP O00764
A	-4	HIS	-	EXPRESSION TAG	UNP O00764
A	-3	SER	-	CLONING ARTIFACT	UNP O00764
A	-2	SER	-	CLONING ARTIFACT	UNP O00764
A	-1	GLY	-	CLONING ARTIFACT	UNP O00764
A	0	LEU	-	CLONING ARTIFACT	UNP O00764
A	1	VAL	-	CLONING ARTIFACT	UNP O00764
A	2	PRO	-	CLONING ARTIFACT	UNP O00764
A	3	ARG	-	CLONING ARTIFACT	UNP O00764
A	4	GLY	-	CLONING ARTIFACT	UNP O00764
A	5	SER	-	CLONING ARTIFACT	UNP O00764
B	-13	MET	-	CLONING ARTIFACT	UNP O00764
B	-12	GLY	-	CLONING ARTIFACT	UNP O00764
B	-11	SER	-	CLONING ARTIFACT	UNP O00764
B	-10	SER	-	CLONING ARTIFACT	UNP O00764
B	-9	HIS	-	EXPRESSION TAG	UNP O00764
B	-8	HIS	-	EXPRESSION TAG	UNP O00764

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	HIS	-	EXPRESSION TAG	UNP O00764
B	-6	HIS	-	EXPRESSION TAG	UNP O00764
B	-5	HIS	-	EXPRESSION TAG	UNP O00764
B	-4	HIS	-	EXPRESSION TAG	UNP O00764
B	-3	SER	-	CLONING ARTIFACT	UNP O00764
B	-2	SER	-	CLONING ARTIFACT	UNP O00764
B	-1	GLY	-	CLONING ARTIFACT	UNP O00764
B	0	LEU	-	CLONING ARTIFACT	UNP O00764
B	1	VAL	-	CLONING ARTIFACT	UNP O00764
B	2	PRO	-	CLONING ARTIFACT	UNP O00764
B	3	ARG	-	CLONING ARTIFACT	UNP O00764
B	4	GLY	-	CLONING ARTIFACT	UNP O00764
B	5	SER	-	CLONING ARTIFACT	UNP O00764


- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	20	Total X 20 20	0	0
3	A	20	Total X 20 20	0	0

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
4	B	1	Total 31	C 10	N 6	O 12	P 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyridoxal kinase



• Molecule 1: Pyridoxal kinase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.13Å 114.97Å 169.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.08 – 2.50 46.07 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.08-2.50) 99.4 (46.07-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.40 (at 2.51Å)	Xtriage
Refinement program	REFMAC refmac_5.2.0005	Depositor
R, R_{free}	0.241 , 0.283 0.244 , 0.287	Depositor DCC
R_{free} test set	936 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	41.3	Xtriage
Anisotropy	0.235	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4693	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/2355	0.64	0/3204
1	B	0.75	0/2318	0.63	0/3157
All	All	0.76	0/4673	0.63	0/6361

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2313	0	2258	26	0
1	B	2276	0	2201	23	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	20	0	0	0	0
3	B	20	0	0	0	0
4	A	31	0	13	1	0
4	B	31	0	13	1	0
All	All	4693	0	4485	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:187:SER:HB2	1:B:199:LEU:HD11	1.62	0.81
1:B:226:VAL:H	4:B:1501:ANP:HN61	1.32	0.77
1:B:12:SER:HB2	1:B:41:VAL:HG22	1.67	0.77
1:B:192:SER:HB2	1:B:200:ILE:HD11	1.77	0.65
1:B:204:SER:HB2	1:B:218:ARG:HG2	1.79	0.64
1:A:69:LEU:HD23	1:A:74:MET:HE3	1.81	0.63
1:B:187:SER:CB	1:B:199:LEU:HD11	2.28	0.63
1:A:172:MET:CE	1:A:204:SER:HB2	2.30	0.61
1:B:154:ALA:O	1:B:158:SER:HB2	2.01	0.60
1:B:149:PRO:O	1:B:185:ILE:HA	2.06	0.56
1:B:12:SER:HB2	1:B:41:VAL:CG2	2.36	0.56
1:A:133:LEU:HB3	1:A:134:PRO:HD3	1.89	0.54
1:B:206:ARG:HH11	1:B:216:MET:HE3	1.72	0.54
1:A:172:MET:HE2	1:A:204:SER:HB2	1.92	0.52
1:B:253:LYS:O	1:B:257:GLU:HG3	2.09	0.52
1:B:128:VAL:HG13	1:B:132:LEU:HD22	1.91	0.51
1:A:172:MET:SD	1:A:185:ILE:HD12	2.52	0.50
1:A:146:ILE:HG12	1:A:182:THR:HB	1.94	0.49
1:B:176:HIS:NE2	1:B:204:SER:OG	2.31	0.48
1:A:79:TYR:CD2	1:A:109:VAL:HB	2.48	0.48
1:A:197:ASN:ND2	1:A:224:ARG:CD	2.77	0.48
1:A:192:SER:HB2	1:A:200:ILE:HD11	1.95	0.47
1:A:226:VAL:H	4:A:501:ANP:HN61	1.62	0.47
1:A:142:PRO:HA	1:A:178:MET:O	2.14	0.47
1:A:220:ARG:NH1	1:A:222:ASP:OD2	2.48	0.47
1:A:236:LEU:HD11	1:A:240:MET:CE	2.45	0.46
1:B:136:TYR:HA	1:B:140:VAL:HB	1.97	0.46
1:A:112:CYS:O	1:A:147:ILE:HA	2.15	0.46
1:A:119:LYS:HA	1:A:124:GLY:HA2	1.98	0.45
1:A:240:MET:HB3	1:A:259:THR:HG23	1.99	0.45
1:A:131:ASP:O	1:A:134:PRO:HD2	2.17	0.45
1:B:220:ARG:HD3	1:B:222:ASP:OD1	2.17	0.44
1:A:236:LEU:CD1	1:A:240:MET:CE	2.95	0.44
1:B:221:MET:CE	1:B:256:CYS:HB3	2.49	0.43
1:A:12:SER:HB2	1:A:41:VAL:HG22	2.01	0.43
1:A:55:GLN:NE2	1:B:61:GLU:HB3	2.34	0.42
1:B:124:GLY:HA3	1:B:152:PHE:CE1	2.54	0.42
1:A:69:LEU:HD23	1:A:74:MET:CE	2.48	0.42
1:A:165:GLN:O	1:A:165:GLN:HG2	2.20	0.42
1:B:69:LEU:HD23	1:B:74:MET:CE	2.49	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:ARG:NH1	1:A:222:ASP:OD1	2.53	0.42
1:A:221:MET:CE	1:A:260:VAL:HG21	2.50	0.42
1:B:158:SER:OG	1:B:171:VAL:HG22	2.20	0.42
1:B:74:MET:HB2	1:B:74:MET:HE2	1.83	0.42
1:A:149:PRO:O	1:A:185:ILE:HA	2.21	0.41
1:B:259:THR:O	1:B:263:LEU:HD23	2.21	0.41
1:B:7:VAL:HG22	1:B:79:TYR:HB2	2.03	0.41
1:A:236:LEU:HD11	1:A:240:MET:HE1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/326 (92%)	286 (96%)	10 (3%)	3 (1%)	15	28
1	B	297/326 (91%)	284 (96%)	11 (4%)	2 (1%)	22	39
All	All	596/652 (91%)	570 (96%)	21 (4%)	5 (1%)	19	35

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121	ASP
1	A	75	ASN
1	B	190	LEU
1	B	179	GLY
1	A	122	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/286 (86%)	245 (99%)	2 (1%)	81	93
1	B	238/286 (83%)	229 (96%)	9 (4%)	33	58
All	All	485/572 (85%)	474 (98%)	11 (2%)	50	76

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	GLU
1	A	280	GLU
1	B	45	ASN
1	B	85	THR
1	B	89	SER
1	B	128	VAL
1	B	194	GLN
1	B	199	LEU
1	B	220	ARG
1	B	222	ASP
1	B	226	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 44 ligands modelled in this entry, 40 are unknown and 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ANP	A	501	2	29,33,33	2.37	8 (27%)	31,52,52	1.82	8 (25%)
4	ANP	B	1501	2	29,33,33	2.34	8 (27%)	31,52,52	2.27	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	A	501	2	-	2/14/38/38	0/3/3/3
4	ANP	B	1501	2	-	2/14/38/38	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1501	ANP	PG-O1G	7.81	1.58	1.46
4	A	501	ANP	PG-O1G	6.45	1.56	1.46
4	A	501	ANP	PG-N3B	5.71	1.78	1.63
4	B	1501	ANP	PG-N3B	5.63	1.78	1.63
4	A	501	ANP	PB-N3B	4.91	1.76	1.63
4	B	1501	ANP	PB-N3B	3.45	1.72	1.63
4	B	1501	ANP	PB-O1B	3.42	1.51	1.46
4	B	1501	ANP	C5-C4	3.10	1.49	1.40
4	A	501	ANP	PB-O1B	2.96	1.50	1.46
4	A	501	ANP	PB-O3A	2.93	1.62	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	ANP	O4'-C1'	2.83	1.45	1.41
4	A	501	ANP	PB-O2B	-2.66	1.49	1.56
4	A	501	ANP	C5-C4	2.65	1.47	1.40
4	B	1501	ANP	C2-N3	2.64	1.36	1.32
4	B	1501	ANP	C2-N1	2.47	1.38	1.33
4	B	1501	ANP	C2'-C1'	-2.04	1.50	1.53

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1501	ANP	O1B-PB-N3B	-7.98	100.01	111.77
4	A	501	ANP	N3-C2-N1	-4.81	121.16	128.68
4	B	1501	ANP	O2B-PB-O1B	4.29	118.92	109.92
4	A	501	ANP	O2B-PB-O1B	4.28	118.89	109.92
4	B	1501	ANP	N3-C2-N1	-3.46	123.26	128.68
4	A	501	ANP	O3G-PG-O2G	3.39	116.67	107.64
4	B	1501	ANP	C3'-C2'-C1'	3.39	106.08	100.98
4	A	501	ANP	C3'-C2'-C1'	3.30	105.94	100.98
4	B	1501	ANP	O2G-PG-O1G	-2.96	106.00	113.45
4	B	1501	ANP	N6-C6-N1	2.87	124.53	118.57
4	B	1501	ANP	PA-O3A-PB	-2.74	122.97	132.62
4	A	501	ANP	PA-O3A-PB	-2.65	123.29	132.62
4	A	501	ANP	C2-N1-C6	2.28	122.66	118.75
4	A	501	ANP	O2'-C2'-C1'	-2.20	102.75	110.85
4	B	1501	ANP	C5-C6-N6	-2.18	117.04	120.35
4	A	501	ANP	O1G-PG-N3B	-2.15	108.60	111.77

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	ANP	PG-N3B-PB-O1B
4	B	1501	ANP	PB-N3B-PG-O1G
4	B	1501	ANP	PG-N3B-PB-O1B
4	A	501	ANP	PB-O3A-PA-O1A

There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	ANP	1	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1501	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/326 (92%)	0.22	16 (5%) 26 28	20, 37, 62, 89	0
1	B	301/326 (92%)	-0.02	8 (2%) 54 58	18, 35, 56, 73	0
All	All	604/652 (92%)	0.10	24 (3%) 38 41	18, 36, 60, 89	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	GLY	5.3
1	A	280	GLU	4.6
1	A	275	LYS	4.4
1	B	210	PRO	3.7
1	A	282	VAL	3.2
1	A	215	VAL	3.2
1	A	281	GLY	3.0
1	B	212	GLY	2.9
1	A	274	ALA	2.8
1	A	103	GLN	2.8
1	B	120	TRP	2.7
1	A	198	TYR	2.6
1	B	71	LEU	2.5
1	A	278	ALA	2.5
1	B	227	ASP	2.4
1	A	283	ARG	2.4
1	A	227	ASP	2.3
1	A	285	SER	2.3
1	A	120	TRP	2.2
1	B	163	HIS	2.2
1	A	276	ALA	2.1
1	B	52	TRP	2.1
1	B	209	ASN	2.1
1	A	122	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	UNX	B	1419	1/1	-0.33	4.02	2,2,2,2	1
3	UNX	B	1415	1/1	-0.22	2.07	2,2,2,2	1
3	UNX	A	408	1/1	-0.21	1.62	2,2,2,2	1
3	UNX	A	420	1/1	-0.02	5.15	2,2,2,2	1
3	UNX	B	1402	1/1	0.00	1.91	2,2,2,2	1
3	UNX	B	412	1/1	0.01	1.24	2,2,2,2	1
3	UNX	A	421	1/1	0.03	1.11	2,2,2,2	1
3	UNX	A	419	1/1	0.04	2.44	2,2,2,2	1
3	UNX	B	1416	1/1	0.07	2.50	2,2,2,2	1
3	UNX	A	1412	1/1	0.12	1.51	2,2,2,2	1
3	UNX	B	404	1/1	0.13	1.25	2,2,2,2	1
3	UNX	B	1417	1/1	0.14	0.96	2,2,2,2	1
3	UNX	B	1414	1/1	0.15	3.69	2,2,2,2	1
3	UNX	A	417	1/1	0.16	2.43	2,2,2,2	1
3	UNX	B	1418	1/1	0.17	1.48	2,2,2,2	1
3	UNX	A	416	1/1	0.20	2.07	2,2,2,2	1
3	UNX	A	410	1/1	0.21	1.59	2,2,2,2	1
3	UNX	B	1410	1/1	0.21	1.69	2,2,2,2	1
3	UNX	A	405	1/1	0.28	1.94	2,2,2,2	1
3	UNX	B	409	1/1	0.28	1.70	2,2,2,2	1
3	UNX	A	418	1/1	0.30	2.88	2,2,2,2	1
3	UNX	A	413	1/1	0.33	1.08	2,2,2,2	1
3	UNX	A	402	1/1	0.33	3.32	2,2,2,2	1
3	UNX	B	1413	1/1	0.37	1.10	2,2,2,2	1
3	UNX	B	1421	1/1	0.38	1.19	2,2,2,2	1
3	UNX	A	1409	1/1	0.39	1.24	2,2,2,2	1

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	UNX	A	414	1/1	0.39	1.91	2,2,2,2	1
3	UNX	B	1407	1/1	0.42	1.51	2,2,2,2	1
3	UNX	B	1406	1/1	0.43	0.88	2,2,2,2	1
3	UNX	A	406	1/1	0.45	0.69	2,2,2,2	1
3	UNX	A	415	1/1	0.48	1.07	2,2,2,2	1
3	UNX	A	1411	1/1	0.51	1.15	2,2,2,2	1
3	UNX	A	1404	1/1	0.52	1.30	2,2,2,2	1
3	UNX	A	403	1/1	0.53	1.16	2,2,2,2	1
3	UNX	A	407	1/1	0.53	1.19	2,2,2,2	1
3	UNX	B	1408	1/1	0.54	1.52	2,2,2,2	1
3	UNX	B	1420	1/1	0.56	2.37	2,2,2,2	1
3	UNX	B	1405	1/1	0.63	0.97	2,2,2,2	1
3	UNX	B	1403	1/1	0.69	1.07	2,2,2,2	1
3	UNX	B	411	1/1	0.70	1.00	2,2,2,2	1
2	MG	B	1401	1/1	0.74	0.21	29,29,29,29	0
2	MG	A	401	1/1	0.87	0.24	42,42,42,42	0
4	ANP	A	501	31/31	0.92	0.16	18,29,34,37	0
4	ANP	B	1501	31/31	0.92	0.18	20,33,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP B 1501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.