



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 06:00 pm BST

PDB ID : 3AJI
Title : Structure of Gankyrin-S6ATPase photo-cross-linked site-specifically, and incorporated by genetic code expansion
Authors : Sato, S.; Mimasu, S.; Sato, A.; Hino, N.; Sakamoto, K.; Umehara, T.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2010-06-07
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>19%</div> <div>••</div> </div> </div>
1	C	231	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>16%</div> <div>••</div> </div> </div>
2	B	83	<div> <div>2%</div> <div> <div></div> <div>42%</div> <div>28%</div> <div>14%</div> <div>•</div> <div>12%</div> </div> </div>
2	D	83	<div> <div>6%</div> <div> <div></div> <div>36%</div> <div>39%</div> <div>10%</div> <div>•</div> <div>12%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 26S proteasome non-ATPase regulatory subunit 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	0	0
			1745	1092	306	337	10			
1	C	229	Total	C	N	O	S	0	0	0
			1745	1092	306	337	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	85	PBF	ARG	ENGINEERED MUTATION	UNP Q9Z2X2
C	85	PBF	ARG	ENGINEERED MUTATION	UNP Q9Z2X2

- Molecule 2 is a protein called Proteasome (Prosome, macropain) 26S subunit, ATPase, 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	73	Total	C	N	O	S	0	0	0
			589	368	103	114	4			
2	D	73	Total	C	N	O	S	0	0	0
			589	368	103	114	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	336	MET	-	EXPRESSION TAG	UNP Q6ZWN9
D	336	MET	-	EXPRESSION TAG	UNP Q6ZWN9

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	144	Total	O	0	0
			144	144		
3	B	22	Total	O	0	0
			22	22		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	142	Total	O	0	0
			142	142		
3	D	19	Total	O	0	0
			19	19		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 26S proteasome non-ATPase regulatory subunit 10




- Molecule 1: 26S proteasome non-ATPase regulatory subunit 10



- Molecule 2: Proteasome (Prosome, macropain) 26S subunit, ATPase, 4



- Molecule 2: Proteasome (Prosome, macropain) 26S subunit, ATPase, 4



GLU
GLN
GLU
HIS
GLU
PHE
TYR
LYS

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	103.16Å 103.16Å 154.99Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.93 – 2.05 42.92 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.93-2.05) 99.8 (42.92-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 2.05Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.171 , 0.229 0.164 , 0.224	Depositor DCC
R_{free} test set	1921 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	29.9	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.477 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4995	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PBF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.49	13/1753 (0.7%)	1.25	10/2368 (0.4%)
1	C	1.48	11/1753 (0.6%)	1.22	12/2368 (0.5%)
2	B	1.39	4/594 (0.7%)	1.58	10/794 (1.3%)
2	D	1.53	5/594 (0.8%)	1.72	13/794 (1.6%)
All	All	1.48	33/4694 (0.7%)	1.35	45/6324 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
2	D	0	2
All	All	0	4

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	356	GLU	C-N	10.45	1.58	1.34
2	D	356	GLU	CD-OE2	9.96	1.36	1.25
2	B	356	GLU	C-N	9.17	1.55	1.34
2	D	356	GLU	CG-CD	-8.99	1.38	1.51
1	A	205	GLU	CB-CG	7.90	1.67	1.52

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	391	ARG	NE-CZ-NH1	-16.09	112.26	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	25	ARG	NE-CZ-NH1	-10.80	114.90	120.30
2	D	391	ARG	NE-CZ-NH2	10.55	125.57	120.30
2	D	390	ASN	N-CA-C	-10.42	82.86	111.00
1	A	25	ARG	NE-CZ-NH1	-9.88	115.36	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	199	TYR	Sidechain
2	B	392	TYR	Sidechain
2	D	363	TYR	Sidechain
2	D	399	PHE	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1731	39	0
1	C	1745	0	1729	31	0
2	B	589	0	604	68	0
2	D	589	0	605	51	0
3	A	144	0	0	3	0
3	B	22	0	0	3	0
3	C	142	0	0	5	1
3	D	19	0	0	2	0
All	All	4995	0	4669	178	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 178 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:340:GLN:HE21	2:D:340:GLN:HA	1.11	1.13

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:390:ASN:CG	2:B:390:ASN:O	1.84	1.11
2:D:391:ARG:O	2:D:392:TYR:HD2	1.36	1.09
2:D:391:ARG:N	2:D:391:ARG:HE	1.58	1.01
1:C:105:ASN:HB2	3:D:157:HOH:O	1.61	0.99

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:312:HOH:O	3:C:369:HOH:O[8_654]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/231 (98%)	222 (98%)	4 (2%)	0	100	100
1	C	226/231 (98%)	222 (98%)	4 (2%)	0	100	100
2	B	71/83 (86%)	68 (96%)	3 (4%)	0	100	100
2	D	71/83 (86%)	68 (96%)	3 (4%)	0	100	100
All	All	594/628 (95%)	580 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/183 (99%)	171 (94%)	10 (6%)	21	13
1	C	181/183 (99%)	170 (94%)	11 (6%)	18	10
2	B	66/76 (87%)	51 (77%)	15 (23%)	1	0
2	D	66/76 (87%)	59 (89%)	7 (11%)	6	2
All	All	494/518 (95%)	451 (91%)	43 (9%)	10	4

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	388	ARG
2	B	405	THR
2	D	367	PRO
2	B	391	ARG
2	B	392	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	380	GLN
1	C	38	GLN
2	D	340	GLN
2	B	353	ASN
1	C	194	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PBF	C	85	1,2	18,19,21	2.24	2 (11%)	21,24,28	5.20	6 (28%)
1	PBF	A	85	1,2	18,19,21	2.55	2 (11%)	21,24,28	7.15	8 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PBF	C	85	1,2	-	2/9/10/16	0/2/2/2
1	PBF	A	85	1,2	-	3/9/10/16	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	85	PBF	CB-CG	9.55	1.74	1.51
1	C	85	PBF	CB-CG	-6.81	1.34	1.51
1	C	85	PBF	CB-CA	4.36	1.63	1.53
1	A	85	PBF	CB-CA	4.04	1.62	1.53

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	85	PBF	CB-CA-C	24.03	156.52	111.47
1	C	85	PBF	CB-CA-C	18.15	145.49	111.47
1	A	85	PBF	CG-CB-CA	17.89	150.32	114.10
1	C	85	PBF	CG-CB-CA	13.62	141.68	114.10
1	A	85	PBF	CB-CG-CD2	-9.39	102.26	120.91

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	85	PBF	CA-CB-CG-CD2
1	C	85	PBF	CA-CB-CG-CD1
1	A	85	PBF	CA-CB-CG-CD1
1	A	85	PBF	CA-CB-CG-CD2
1	A	85	PBF	C-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	85	PBF	4	0
1	A	85	PBF	5	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	228/231 (98%)	-0.65	2 (0%) 84 86	20, 28, 48, 104	0
1	C	228/231 (98%)	-0.68	2 (0%) 84 86	20, 27, 48, 104	0
2	B	73/83 (87%)	0.01	2 (2%) 54 59	28, 52, 81, 91	0
2	D	73/83 (87%)	0.17	5 (6%) 17 18	29, 52, 84, 101	0
All	All	602/628 (95%)	-0.48	11 (1%) 68 71	20, 30, 69, 104	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	MET	5.0
1	C	231	MET	4.7
1	C	3	GLY	3.0
2	D	387	VAL	2.9
2	B	392	TYR	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PBF	A	85	18/20	0.87	0.15	21,39,73,76	0
1	PBF	C	85	18/20	0.92	0.14	19,45,70,78	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.