



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 12:24 am BST

PDB ID : 3AK9  
Title : Crystal structure of the SEp22 dodecamer, a Dps-like protein from *Salmonella enterica* subsp. *enterica* serovar Enteritidis, FE-soaked form  
Authors : Miyamoto, T.; Asahina, Y.; Miyazaki, S.; Shimizu, H.; Ohto, U.; Noguchi, S.; Satow, Y.  
Deposited on : 2010-07-08  
Resolution : 1.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>
1	B	167	<div> <div>4%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>•</div> </div> </div>
1	C	167	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>7%</div> </div> </div>
1	D	167	<div> <div>•</div> <div> <div></div> <div>84%</div> <div>8%</div> <div>8%</div> </div> </div>
1	E	167	<div> <div>•</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>
1	F	167	<div> <div>•</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>7%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	167	<div><div>%</div><div><div></div><div>86%</div><div>7%</div><div>7%</div></div></div>
1	H	167	<div><div>%</div><div><div></div><div>85%</div><div>8%</div><div>7%</div></div></div>
1	I	167	<div><div>%</div><div><div></div><div>84%</div><div>8%</div><div>7%</div></div></div>
1	J	167	<div><div>%</div><div><div></div><div>82%</div><div>11%</div><div>7%</div></div></div>
1	K	167	<div><div>%</div><div><div></div><div>86%</div><div>7%</div><div>7%</div></div></div>
1	L	167	<div><div>%</div><div><div></div><div>86%</div><div>7%</div><div>7%</div></div></div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA protection during starvation protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	155	Total	C	N	O	S	0	1	0
			1236	778	215	240	3			
1	B	155	Total	C	N	O	S	0	1	0
			1234	775	215	241	3			
1	C	155	Total	C	N	O	S	0	3	0
			1241	781	215	242	3			
1	D	154	Total	C	N	O	S	0	2	0
			1229	773	213	240	3			
1	E	155	Total	C	N	O	S	0	3	0
			1242	782	215	242	3			
1	F	155	Total	C	N	O	S	0	0	0
			1231	773	215	240	3			
1	G	155	Total	C	N	O	S	0	1	0
			1234	775	215	241	3			
1	H	155	Total	C	N	O	S	0	2	0
			1239	780	215	241	3			
1	I	155	Total	C	N	O	S	0	0	0
			1231	773	215	240	3			
1	J	155	Total	C	N	O	S	0	0	0
			1231	773	215	240	3			
1	K	155	Total	C	N	O	S	0	3	0
			1242	782	215	242	3			
1	L	155	Total	C	N	O	S	0	3	0
			1242	782	215	242	3			

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Fe	0	0
			1	1		
2	J	1	Total	Fe	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Fe	0	0
			1	1		
2	K	1	Total	Fe	0	0
			1	1		
2	E	1	Total	Fe	0	0
			1	1		
2	H	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		
2	I	1	Total	Fe	0	0
			1	1		
2	C	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		
2	L	1	Total	Fe	0	0
			1	1		
2	F	1	Total	Fe	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	O	S	0	0
			5	4	1		
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		
3	J	1	Total	O	S	0	0
			5	4	1		
3	K	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	1	Total	Mg	0	0
			1	1		
4	G	1	Total	Mg	0	0
			1	1		
4	I	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	97	Total	O	0	0
			97	97		
5	B	102	Total	O	0	0
			102	102		
5	C	116	Total	O	0	0
			116	116		
5	D	118	Total	O	0	0
			118	118		
5	E	105	Total	O	0	0
			105	105		
5	F	123	Total	O	0	0
			123	123		
5	G	107	Total	O	0	0
			107	107		
5	H	116	Total	O	0	0
			116	116		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	I	114	Total 114	O 114	0	0
5	J	101	Total 101	O 101	0	0
5	K	107	Total 107	O 107	0	0
5	L	115	Total 115	O 115	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA protection during starvation protein




- Molecule 1: DNA protection during starvation protein




- Molecule 1: DNA protection during starvation protein



- Molecule 1: DNA protection during starvation protein




- Molecule 1: DNA protection during starvation protein




- Molecule 1: DNA protection during starvation protein







- Molecule 1: DNA protection during starvation protein




- Molecule 1: DNA protection during starvation protein




- Molecule 1: DNA protection during starvation protein




- Molecule 1: DNA protection during starvation protein



- Molecule 1: DNA protection during starvation protein



- Molecule 1: DNA protection during starvation protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.47Å 98.00Å 208.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.60 – 1.30 48.60 – 1.30	Depositor EDS
% Data completeness (in resolution range)	90.8 (48.60-1.30) 90.8 (48.60-1.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 1.30Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.153 , 0.172 0.161 , 0.181	Depositor DCC
$R_{free}$ test set	21511 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.2	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 46.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	16199	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FE2, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/1258	0.65	0/1703
1	B	0.48	0/1256	0.70	1/1700 (0.1%)
1	C	0.49	0/1269	0.67	1/1718 (0.1%)
1	D	0.47	0/1254	0.69	0/1697
1	E	0.49	0/1270	0.64	0/1719
1	F	0.50	0/1250	0.68	2/1692 (0.1%)
1	G	0.46	0/1256	0.63	0/1700
1	H	0.48	0/1264	0.64	0/1711
1	I	0.48	0/1250	0.68	0/1692
1	J	0.49	0/1250	0.67	0/1692
1	K	0.47	0/1270	0.68	0/1719
1	L	0.49	0/1270	0.65	0/1719
All	All	0.48	0/15117	0.67	4/20462 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	70	ARG	NE-CZ-NH2	-6.19	117.20	120.30
1	B	70	ARG	NE-CZ-NH2	-5.85	117.38	120.30
1	F	70	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	F	70	ARG	NE-CZ-NH2	-5.43	117.58	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1236	0	1230	8	0
1	B	1234	0	1224	3	0
1	C	1241	0	1238	14	0
1	D	1229	0	1223	11	0
1	E	1242	0	1240	7	0
1	F	1231	0	1219	7	0
1	G	1234	0	1224	9	0
1	H	1239	0	1235	8	0
1	I	1231	0	1219	11	0
1	J	1231	0	1219	17	0
1	K	1242	0	1240	6	0
1	L	1242	0	1240	6	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	1	0
3	F	5	0	0	0	0
3	J	5	0	0	0	0
3	K	5	0	0	1	0
4	D	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
5	A	97	0	0	2	0
5	B	102	0	0	0	0
5	C	116	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	118	0	0	1	0
5	E	105	0	0	0	0
5	F	123	0	0	1	0
5	G	107	0	0	0	0
5	H	116	0	0	0	0
5	I	114	0	0	1	0
5	J	101	0	0	0	0
5	K	107	0	0	0	0
5	L	115	0	0	0	0
All	All	16199	0	14751	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 103 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:35:ARG:HH22	1:C:131:ASP:HB3	1.29	0.95
1:C:35:ARG:NH2	1:C:131:ASP:HB3	1.89	0.88
1:C:35:ARG:HH22	1:C:131:ASP:CB	1.87	0.86
1:J:43:LEU:HD22	1:J:128:VAL:HG11	1.60	0.82
1:K:74:THR:HG21	3:K:169:SO4:O4	1.86	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/167 (92%)	152 (99%)	2 (1%)	0	100	100
1	B	154/167 (92%)	152 (99%)	2 (1%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	156/167 (93%)	154 (99%)	2 (1%)	0	100	100
1	D	154/167 (92%)	153 (99%)	1 (1%)	0	100	100
1	E	156/167 (93%)	155 (99%)	1 (1%)	0	100	100
1	F	153/167 (92%)	152 (99%)	1 (1%)	0	100	100
1	G	154/167 (92%)	152 (99%)	2 (1%)	0	100	100
1	H	155/167 (93%)	155 (100%)	0	0	100	100
1	I	153/167 (92%)	151 (99%)	2 (1%)	0	100	100
1	J	153/167 (92%)	152 (99%)	1 (1%)	0	100	100
1	K	156/167 (93%)	155 (99%)	1 (1%)	0	100	100
1	L	156/167 (93%)	155 (99%)	1 (1%)	0	100	100
All	All	1854/2004 (92%)	1838 (99%)	16 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/143 (94%)	133 (99%)	1 (1%)	84	61
1	B	134/143 (94%)	134 (100%)	0	100	100
1	C	136/143 (95%)	136 (100%)	0	100	100
1	D	134/143 (94%)	134 (100%)	0	100	100
1	E	136/143 (95%)	136 (100%)	0	100	100
1	F	133/143 (93%)	133 (100%)	0	100	100
1	G	134/143 (94%)	134 (100%)	0	100	100
1	H	135/143 (94%)	135 (100%)	0	100	100
1	I	133/143 (93%)	133 (100%)	0	100	100
1	J	133/143 (93%)	133 (100%)	0	100	100
1	K	136/143 (95%)	136 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	136/143 (95%)	136 (100%)	0	100	100
All	All	1614/1716 (94%)	1613 (100%)	1 (0%)	93	79

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	117	HIS
1	F	117	HIS
1	K	117	HIS
1	F	39	GLN
1	F	113	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 22 ligands modelled in this entry, 16 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	D	169	-	4,4,4	0.17	0	6,6,6	0.26	0
3	SO4	F	169	-	4,4,4	0.17	0	6,6,6	0.34	0
3	SO4	K	169	-	4,4,4	0.36	0	6,6,6	0.46	0
3	SO4	E	169	-	4,4,4	0.12	0	6,6,6	0.56	0
3	SO4	C	169	-	4,4,4	0.31	0	6,6,6	0.39	0
3	SO4	J	169	-	4,4,4	0.22	0	6,6,6	0.39	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	K	169	SO4	1	0
3	E	169	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	155/167 (92%)	-0.11	3 (1%) 66 67	19, 24, 35, 43	0
1	B	155/167 (92%)	-0.01	6 (3%) 39 36	18, 23, 33, 43	0
1	C	155/167 (92%)	-0.04	4 (2%) 56 54	20, 24, 32, 41	0
1	D	154/167 (92%)	-0.05	1 (0%) 89 88	19, 23, 30, 35	0
1	E	155/167 (92%)	-0.25	2 (1%) 77 79	21, 24, 33, 42	0
1	F	155/167 (92%)	-0.29	2 (1%) 77 79	19, 23, 31, 39	0
1	G	155/167 (92%)	-0.21	2 (1%) 77 79	19, 23, 32, 44	0
1	H	155/167 (92%)	-0.27	1 (0%) 89 88	19, 23, 31, 42	0
1	I	155/167 (92%)	-0.21	1 (0%) 89 88	18, 23, 31, 41	0
1	J	155/167 (92%)	-0.17	2 (1%) 77 79	18, 23, 32, 41	0
1	K	155/167 (92%)	-0.21	1 (0%) 89 88	20, 23, 31, 41	0
1	L	155/167 (92%)	-0.25	1 (0%) 89 88	20, 23, 33, 37	0
All	All	1859/2004 (92%)	-0.17	26 (1%) 75 77	18, 23, 32, 44	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	13	ASN	6.3
1	E	13	ASN	5.9
1	B	13	ASN	5.8
1	G	13	ASN	5.8
1	H	13	ASN	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	E	169	5/5	0.73	0.35	26,44,45,46	0
4	MG	I	169	1/1	0.91	0.21	52,52,52,52	0
3	SO4	F	169	5/5	0.92	0.27	31,38,39,39	0
4	MG	H	169	1/1	0.92	0.20	56,56,56,56	0
3	SO4	K	169	5/5	0.93	0.26	33,39,40,40	0
3	SO4	D	169	5/5	0.94	0.22	31,37,39,40	0
4	MG	D	170	1/1	0.96	0.13	36,36,36,36	0
3	SO4	C	169	5/5	0.96	0.25	32,39,39,41	0
4	MG	G	169	1/1	0.97	0.21	36,36,36,36	0
2	FE2	K	168	1/1	0.97	0.04	25,25,25,25	0
3	SO4	J	169	5/5	0.97	0.17	33,38,39,40	0
2	FE2	C	168	1/1	0.98	0.04	24,24,24,24	0
2	FE2	E	168	1/1	0.98	0.04	24,24,24,24	0
2	FE2	B	168	1/1	0.98	0.04	24,24,24,24	0
2	FE2	L	168	1/1	0.99	0.05	23,23,23,23	0
2	FE2	J	168	1/1	0.99	0.04	24,24,24,24	0
2	FE2	D	168	1/1	0.99	0.03	24,24,24,24	0
2	FE2	F	168	1/1	0.99	0.03	23,23,23,23	0
2	FE2	A	168	1/1	0.99	0.05	25,25,25,25	0
2	FE2	G	168	1/1	1.00	0.05	25,25,25,25	0
2	FE2	H	168	1/1	1.00	0.04	23,23,23,23	0
2	FE2	I	168	1/1	1.00	0.04	25,25,25,25	0

### 6.5 Other polymers ⓘ

There are no such residues in this entry.