



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 09:52 am BST

PDB ID : 4AKY  
Title : CRYSTAL STRUCTURE OF VIRB8 FROM BRUCELLA SUIS IN COM-  
PLEX WITH INTERACTION INHIBITOR 2-(butylamino)-8-quinolinol  
Authors : Coincon, M.; Smith, M.A.; Sygusch, J.; Baron, C.  
Deposited on : 2012-02-29  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	<div> <div>2%</div> <div>87%</div> <div>10%</div> <div>.</div> </div>
1	B	138	<div> <div>4%</div> <div>78%</div> <div>17%</div> <div>.</div> </div>
1	C	138	<div> <div>9%</div> <div>80%</div> <div>17%</div> <div>.</div> </div>
1	D	138	<div> <div>13%</div> <div>80%</div> <div>17%</div> <div>.</div> </div>
1	E	138	<div> <div>14%</div> <div>78%</div> <div>20%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	4LL	C	1235	-	-	-	X
2	4LL	D	1235	-	X	-	-
2	4LL	E	1235	-	X	-	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TYPE IV SECRETION SYSTEM PROTEIN VIRB8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	134	Total	C	N	O	S	0	0	0
			1065	673	176	213	3			
1	B	134	Total	C	N	O	S	0	0	0
			1066	673	176	214	3			
1	C	134	Total	C	N	O	S	0	0	0
			1066	673	176	214	3			
1	D	134	Total	C	N	O	S	0	0	0
			1066	673	176	214	3			
1	E	134	Total	C	N	O	S	0	0	0
			1066	673	176	214	3			

- Molecule 2 is 2-(butylamino)quinolin-8-ol (three-letter code: 4LL) (formula: C<sub>13</sub>H<sub>16</sub>N<sub>2</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	13	2	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			16	13	2	1		
2	C	1	Total	C	N	O	0	0
			16	13	2	1		
2	D	1	Total	C	N	O	0	0
			16	13	2	1		
2	E	1	Total	C	N	O	0	0
			16	13	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	126	Total	O	0	0
			126	126		
3	B	97	Total	O	0	0
			97	97		
3	C	97	Total	O	0	0
			97	97		
3	D	76	Total	O	0	0
			76	76		
3	E	84	Total	O	0	0
			84	84		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8





- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8




- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8




- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8



- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	198.44Å 198.44Å 103.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.56 – 2.60 48.56 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.56-2.60) 100.0 (48.56-2.60)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.80 (at 2.61Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.198 , 0.236 0.201 , 0.200	Depositor DCC
$R_{free}$ test set	2391 reflections (7.50%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.3	Xtriage
Anisotropy	0.320	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 86.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5889	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4LL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/1089	0.47	0/1480
1	B	0.43	0/1090	0.47	0/1481
1	C	0.38	0/1090	0.45	0/1481
1	D	0.34	0/1090	0.43	0/1481
1	E	0.47	0/1090	0.45	0/1481
All	All	0.41	0/5449	0.45	0/7404

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1065	0	1020	11	0
1	B	1066	0	1023	18	0
1	C	1066	0	1023	20	0
1	D	1066	0	1023	19	0
1	E	1066	0	1023	21	0
2	A	16	0	16	2	0
2	B	16	0	16	5	0
2	C	16	0	16	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	16	0	16	5	0
2	E	16	0	16	3	0
3	A	126	0	0	0	0
3	B	97	0	0	0	0
3	C	97	0	0	2	1
3	D	76	0	0	0	0
3	E	84	0	0	1	0
All	All	5889	0	5192	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:97:SER:HB3	3:C:2002:HOH:O	1.65	0.96
1:E:231:VAL:HG11	2:E:1235:4LL:H04	1.53	0.88
1:B:231:VAL:HG11	2:B:1235:4LL:H04	1.56	0.85
1:E:117:TYR:HB2	2:E:1235:4LL:H12	1.64	0.78
1:B:117:TYR:HB2	2:B:1235:4LL:H12	1.71	0.72
1:D:231:VAL:HG11	2:D:1235:4LL:H04	1.71	0.72
1:E:124:LYS:HB3	3:E:2016:HOH:O	1.88	0.71
1:D:155:TYR:HB3	1:D:159:VAL:HB	1.75	0.69
1:D:98:TYR:HH	1:E:98:TYR:HE1	1.41	0.69
1:A:117:TYR:HB2	2:A:1235:4LL:H12	1.74	0.68
1:D:141:TYR:HE1	2:D:1235:4LL:H10	1.58	0.68
1:B:102:MET:HE2	1:B:106:TRP:HE1	1.59	0.68
1:D:212:MET:CE	1:D:223:GLY:HA2	2.25	0.67
1:E:155:TYR:HB3	1:E:159:VAL:HB	1.76	0.67
1:A:212:MET:CE	1:A:223:GLY:HA2	2.27	0.64
1:C:212:MET:HE1	1:C:223:GLY:HA2	1.79	0.63
1:D:141:TYR:CE1	2:D:1235:4LL:H10	2.34	0.63
1:C:151:LEU:HD22	1:C:155:TYR:CE2	2.34	0.62
1:B:209:PRO:HA	1:B:212:MET:HE3	1.83	0.60
1:C:135:PRO:HB3	1:E:207:VAL:HG11	1.85	0.58
1:A:144:GLN:HE22	2:A:1235:4LL:C15	2.17	0.57
1:B:144:GLN:HE22	2:B:1235:4LL:C15	2.16	0.57
1:C:212:MET:CE	1:C:223:GLY:HA2	2.34	0.57
1:A:212:MET:HE1	1:A:223:GLY:HA2	1.86	0.56
1:E:144:GLN:HE22	2:E:1235:4LL:C16	2.19	0.54
1:E:161:THR:HG21	1:E:182:LYS:HE2	1.89	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:182:LYS:NZ	2:C:1235:4LL:H02A	2.21	0.54
1:C:124:LYS:HB3	3:C:2020:HOH:O	2.07	0.53
1:A:212:MET:HB3	1:A:216:ALA:HB3	1.92	0.52
1:E:151:LEU:HD22	1:E:155:TYR:CE2	2.45	0.52
1:D:212:MET:HE3	1:D:223:GLY:HA2	1.90	0.52
1:A:144:GLN:HG3	1:B:230:ARG:NH1	2.24	0.52
1:D:179:ARG:HG2	1:D:199:ILE:HG12	1.90	0.52
1:D:105:TYR:O	1:D:109:GLN:HG2	2.10	0.51
1:B:209:PRO:HA	1:B:212:MET:CE	2.40	0.51
1:D:222:LEU:HD11	1:E:102:MET:HE1	1.92	0.51
1:E:125:ASP:O	1:E:129:VAL:HG23	2.11	0.51
1:D:125:ASP:O	1:D:129:VAL:HG23	2.12	0.50
1:C:212:MET:HB3	1:C:216:ALA:HB3	1.93	0.50
1:B:102:MET:CE	1:B:106:TRP:HE1	2.24	0.50
1:D:114:ARG:O	2:D:1235:4LL:C10	2.60	0.49
1:A:98:TYR:HH	1:C:98:TYR:HH	1.57	0.49
1:B:179:ARG:NH2	1:B:234:GLU:OE1	2.46	0.49
1:D:151:LEU:HD22	1:D:155:TYR:CE2	2.47	0.49
1:E:102:MET:HG2	1:E:222:LEU:CD1	2.42	0.49
1:C:97:SER:HB2	1:C:100:THR:OG1	2.14	0.48
1:D:212:MET:HE1	1:D:223:GLY:HA2	1.94	0.47
1:E:212:MET:CE	1:E:223:GLY:HA2	2.44	0.47
1:C:158:ASN:HA	1:C:187:THR:HB	1.96	0.47
1:B:171:ASN:ND2	1:B:173:LYS:HB2	2.30	0.47
1:C:211:LEU:HD23	1:E:126:TYR:CE1	2.49	0.47
1:E:212:MET:HE1	1:E:223:GLY:HA2	1.96	0.46
1:A:125:ASP:O	1:A:129:VAL:HG23	2.15	0.46
1:C:152:ASP:N	1:C:152:ASP:OD1	2.49	0.46
1:C:111:VAL:HG21	1:C:178:VAL:HG11	1.98	0.46
1:C:117:TYR:HB2	2:C:1235:4LL:H12	1.98	0.46
1:C:135:PRO:HB3	1:E:207:VAL:CG1	2.46	0.46
1:B:100:THR:O	1:B:104:LYS:HG3	2.17	0.45
1:D:212:MET:HB3	1:D:216:ALA:HB3	1.99	0.44
1:D:104:LYS:HE2	1:D:168:ILE:O	2.18	0.43
1:B:155:TYR:HB3	1:B:159:VAL:HB	2.00	0.42
1:E:203:GLY:HA3	1:E:228:SER:OG	2.20	0.42
1:E:106:TRP:CZ2	1:E:221:PRO:HD2	2.54	0.42
1:E:178:VAL:HB	1:E:200:ALA:HB3	2.02	0.42
1:A:102:MET:HG2	1:A:222:LEU:HD11	2.02	0.42
1:D:118:ASP:HB3	1:D:121:THR:OG1	2.20	0.41
1:B:119:TRP:HB2	1:B:156:GLY:O	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:GLN:HE22	2:B:1235:4LL:H15	1.84	0.41
1:B:152:ASP:OD1	1:B:152:ASP:N	2.52	0.41
1:C:182:LYS:HZ3	2:C:1235:4LL:H02A	1.84	0.41
1:C:207:VAL:HG11	1:E:135:PRO:HB3	2.02	0.41
1:B:122:LEU:HD11	1:B:145:PHE:CZ	2.56	0.41
1:A:221:PRO:HG3	1:C:105:TYR:HB2	2.03	0.41
1:D:168:ILE:HG23	1:D:178:VAL:HG22	2.03	0.41
1:A:100:THR:O	1:A:104:LYS:HG3	2.21	0.40
1:D:117:TYR:HB2	2:D:1235:4LL:H12	2.03	0.40
1:E:137:GLU:HA	1:E:140:SER:HB3	2.02	0.40
1:C:208:ASN:HA	1:C:209:PRO:HD3	1.92	0.40
1:B:114:ARG:O	2:B:1235:4LL:C10	2.69	0.40
1:B:128:THR:O	1:B:132:LEU:HG	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:2048:HOH:O	3:C:2048:HOH:O[6_546]	1.71	0.49

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	130/138 (94%)	126 (97%)	4 (3%)	0	100	100
1	B	130/138 (94%)	126 (97%)	4 (3%)	0	100	100
1	C	130/138 (94%)	129 (99%)	1 (1%)	0	100	100
1	D	130/138 (94%)	128 (98%)	2 (2%)	0	100	100
1	E	130/138 (94%)	125 (96%)	5 (4%)	0	100	100
All	All	650/690 (94%)	634 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/121 (97%)	116 (99%)	1 (1%)	78	91
1	B	118/121 (98%)	114 (97%)	4 (3%)	37	63
1	C	118/121 (98%)	116 (98%)	2 (2%)	60	81
1	D	118/121 (98%)	118 (100%)	0	100	100
1	E	118/121 (98%)	118 (100%)	0	100	100
All	All	589/605 (97%)	582 (99%)	7 (1%)	71	87

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	231	VAL
1	B	124	LYS
1	B	144	GLN
1	B	186	ARG
1	B	231	VAL
1	C	136	SER
1	C	231	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	4LL	E	1235	-	17,17,17	5.93	12 (70%)	21,22,22	1.16	3 (14%)
2	4LL	D	1235	-	17,17,17	5.89	12 (70%)	21,22,22	1.19	3 (14%)
2	4LL	C	1235	-	17,17,17	5.85	12 (70%)	21,22,22	1.03	1 (4%)
2	4LL	B	1235	-	17,17,17	5.93	12 (70%)	21,22,22	1.19	3 (14%)
2	4LL	A	1235	-	17,17,17	5.91	12 (70%)	21,22,22	1.14	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4LL	E	1235	-	-	4/5/5/5	0/2/2/2
2	4LL	D	1235	-	-	4/5/5/5	0/2/2/2
2	4LL	C	1235	-	-	3/5/5/5	0/2/2/2
2	4LL	B	1235	-	-	3/5/5/5	0/2/2/2
2	4LL	A	1235	-	-	3/5/5/5	0/2/2/2

All (60) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1235	4LL	C12-C13	8.76	1.54	1.37
2	B	1235	4LL	C12-C13	8.69	1.54	1.37
2	A	1235	4LL	C12-C13	8.67	1.54	1.37
2	D	1235	4LL	C12-C13	8.65	1.54	1.37
2	C	1235	4LL	C12-C13	8.52	1.54	1.37
2	E	1235	4LL	C09-C08	8.24	1.54	1.42
2	C	1235	4LL	C09-C08	8.22	1.54	1.42
2	A	1235	4LL	C09-C08	8.21	1.54	1.42
2	D	1235	4LL	C09-C08	8.19	1.54	1.42
2	B	1235	4LL	C09-C08	8.19	1.54	1.42
2	D	1235	4LL	C13-C08	7.94	1.54	1.42
2	E	1235	4LL	C13-C08	7.94	1.54	1.42
2	B	1235	4LL	C13-C08	7.93	1.54	1.42
2	A	1235	4LL	C13-C08	7.76	1.54	1.42
2	B	1235	4LL	C11-C10	7.75	1.54	1.36
2	A	1235	4LL	C11-C10	7.71	1.54	1.36
2	E	1235	4LL	C11-C10	7.71	1.54	1.36
2	D	1235	4LL	C11-C10	7.66	1.54	1.36
2	C	1235	4LL	C11-C10	7.59	1.54	1.36
2	C	1235	4LL	C13-C08	7.53	1.54	1.42
2	A	1235	4LL	C08-N07	7.40	1.51	1.37
2	B	1235	4LL	C08-N07	7.36	1.51	1.37
2	C	1235	4LL	C08-N07	7.35	1.51	1.37
2	D	1235	4LL	C08-N07	7.34	1.51	1.37
2	E	1235	4LL	C08-N07	7.33	1.51	1.37
2	B	1235	4LL	C11-C12	7.31	1.54	1.38
2	A	1235	4LL	C11-C12	7.29	1.54	1.38
2	E	1235	4LL	C11-C12	7.23	1.54	1.38
2	C	1235	4LL	C11-C12	7.15	1.54	1.38
2	D	1235	4LL	C11-C12	7.11	1.53	1.38
2	B	1235	4LL	C16-C06	7.02	1.55	1.39
2	C	1235	4LL	C16-C06	7.01	1.55	1.39
2	E	1235	4LL	C16-C06	7.00	1.55	1.39
2	A	1235	4LL	C16-C06	6.97	1.55	1.39
2	D	1235	4LL	C16-C06	6.94	1.55	1.39
2	B	1235	4LL	C15-C16	6.80	1.51	1.36
2	D	1235	4LL	C06-N05	6.77	1.46	1.36
2	E	1235	4LL	C06-N05	6.77	1.46	1.36
2	C	1235	4LL	C15-C16	6.74	1.50	1.36
2	C	1235	4LL	C06-N05	6.74	1.46	1.36
2	E	1235	4LL	C15-C16	6.73	1.50	1.36
2	A	1235	4LL	C15-C16	6.68	1.50	1.36
2	A	1235	4LL	C06-N05	6.67	1.46	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1235	4LL	C15-C16	6.67	1.50	1.36
2	B	1235	4LL	C06-N05	6.60	1.46	1.36
2	A	1235	4LL	C06-N07	5.99	1.50	1.33
2	B	1235	4LL	C06-N07	5.94	1.50	1.33
2	E	1235	4LL	C06-N07	5.93	1.50	1.33
2	D	1235	4LL	C06-N07	5.88	1.50	1.33
2	C	1235	4LL	C06-N07	5.86	1.50	1.33
2	B	1235	4LL	C10-C09	5.47	1.55	1.41
2	A	1235	4LL	C10-C09	5.43	1.55	1.41
2	E	1235	4LL	C10-C09	5.43	1.55	1.41
2	C	1235	4LL	C10-C09	5.36	1.54	1.41
2	D	1235	4LL	C10-C09	5.35	1.54	1.41
2	B	1235	4LL	C15-C09	4.10	1.51	1.41
2	C	1235	4LL	C15-C09	4.03	1.51	1.41
2	A	1235	4LL	C15-C09	4.01	1.51	1.41
2	E	1235	4LL	C15-C09	3.98	1.51	1.41
2	D	1235	4LL	C15-C09	3.95	1.51	1.41

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1235	4LL	C13-C08-N07	2.93	120.28	117.33
2	E	1235	4LL	C13-C08-N07	2.88	120.23	117.33
2	B	1235	4LL	C13-C08-N07	2.77	120.11	117.33
2	A	1235	4LL	C13-C08-N07	2.75	120.10	117.33
2	B	1235	4LL	C04-N05-C06	-2.55	119.35	123.38
2	D	1235	4LL	C04-N05-C06	-2.27	119.78	123.38
2	D	1235	4LL	C09-C08-N07	-2.25	119.64	122.64
2	C	1235	4LL	C13-C08-N07	2.22	119.56	117.33
2	A	1235	4LL	C09-C08-N07	-2.19	119.73	122.64
2	B	1235	4LL	C09-C08-N07	-2.15	119.78	122.64
2	E	1235	4LL	C09-C08-N07	-2.13	119.81	122.64
2	A	1235	4LL	C04-N05-C06	-2.12	120.03	123.38
2	E	1235	4LL	C04-N05-C06	-2.01	120.20	123.38

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1235	4LL	N07-C06-N05-C04
2	E	1235	4LL	C16-C06-N05-C04
2	D	1235	4LL	N07-C06-N05-C04

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	D	1235	4LL	C16-C06-N05-C04
2	C	1235	4LL	N07-C06-N05-C04
2	C	1235	4LL	C16-C06-N05-C04
2	B	1235	4LL	N07-C06-N05-C04
2	B	1235	4LL	C16-C06-N05-C04
2	A	1235	4LL	N07-C06-N05-C04
2	A	1235	4LL	C16-C06-N05-C04
2	C	1235	4LL	C02-C03-C04-N05
2	A	1235	4LL	C02-C03-C04-N05
2	B	1235	4LL	C01-C02-C03-C04
2	E	1235	4LL	C01-C02-C03-C04
2	D	1235	4LL	C02-C03-C04-N05
2	E	1235	4LL	C03-C04-N05-C06
2	D	1235	4LL	C01-C02-C03-C04

There are no ring outliers.

5 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1235	4LL	3	0
2	D	1235	4LL	5	0
2	C	1235	4LL	3	0
2	B	1235	4LL	5	0
2	A	1235	4LL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	134/138 (97%)	0.16	3 (2%) 62 56	30, 48, 92, 116	0
1	B	134/138 (97%)	0.17	5 (3%) 41 34	27, 44, 84, 106	0
1	C	134/138 (97%)	0.33	12 (8%) 9 6	33, 52, 98, 128	0
1	D	134/138 (97%)	0.69	18 (13%) 3 2	46, 67, 110, 140	0
1	E	134/138 (97%)	0.63	20 (14%) 2 1	48, 68, 123, 135	0
All	All	670/690 (97%)	0.39	58 (8%) 10 7	27, 58, 103, 140	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	147	GLY	7.0
1	E	148	ASP	6.8
1	E	154	GLN	6.8
1	D	148	ASP	6.6
1	D	149	LYS	5.8
1	E	149	LYS	5.4
1	E	153	LYS	4.9
1	C	192	ASP	4.9
1	E	186	ARG	4.8
1	D	154	GLN	4.7
1	E	150	ALA	4.4
1	E	194	GLU	4.0
1	C	149	LYS	4.0
1	D	155	TYR	4.0
1	D	147	GLY	4.0
1	E	159	VAL	3.9
1	C	187	THR	3.9
1	D	187	THR	3.9
1	C	186	ARG	3.7
1	C	150	ALA	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	171	ASN	3.5
1	D	186	ARG	3.3
1	E	120	TYR	3.3
1	D	153	LYS	3.2
1	E	185	LYS	3.2
1	E	155	TYR	3.2
1	E	145	PHE	3.2
1	E	119	TRP	3.1
1	E	151	LEU	3.1
1	D	146	GLN	3.1
1	A	148	ASP	2.9
1	E	146	GLN	2.9
1	D	98	TYR	2.9
1	D	230	ARG	2.9
1	E	187	THR	2.8
1	D	119	TRP	2.8
1	C	146	GLN	2.8
1	D	151	LEU	2.8
1	C	173	LYS	2.8
1	C	194	GLU	2.7
1	D	185	LYS	2.7
1	D	194	GLU	2.7
1	A	192	ASP	2.7
1	C	193	GLY	2.6
1	C	98	TYR	2.5
1	C	151	LEU	2.5
1	E	184	THR	2.4
1	D	159	VAL	2.4
1	B	98	TYR	2.3
1	B	186	ARG	2.3
1	B	119	TRP	2.3
1	E	182	LYS	2.3
1	A	187	THR	2.2
1	E	158	ASN	2.1
1	B	155	TYR	2.1
1	D	150	ALA	2.0
1	B	148	ASP	2.0
1	D	184	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	4LL	E	1235	16/16	0.72	0.34	64,78,82,84	16
2	4LL	C	1235	16/16	0.76	0.46	56,68,79,80	16
2	4LL	D	1235	16/16	0.80	0.27	64,73,80,88	16
2	4LL	A	1235	16/16	0.83	0.35	52,60,71,72	16
2	4LL	B	1235	16/16	0.85	0.33	52,58,66,66	16

### 6.5 Other polymers [i](#)

There are no such residues in this entry.