



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 03:25 am BST

PDB ID : 2AL1
Title : Crystal Structure Analysis of Enolase Mg Subunit Complex at pH 8.0
Authors : Sims, P.A.; Menefee, A.L.; Larsen, T.M.; Mansoorabadi, S.O.; Reed, G.H.
Deposited on : 2005-08-04
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	<div> <div>8%</div> <div> <div></div> <div>86%</div> <div>14%</div> <div>.</div> </div> </div>
1	B	436	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>..</div> </div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7298 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called enolase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	0	0
			3292	2079	570	637	6			
1	B	431	Total	C	N	O	S	0	0	0
			3263	2063	563	631	6			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is PHOSPHOENOLPYRUVATE (three-letter code: PEP) (formula: C₃H₅O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	1
			10	3	6	1		
3	B	1	Total	C	O	P	0	0
			10	3	6	1		

- Molecule 4 is 2-PHOSPHOGLYCERIC ACID (three-letter code: 2PG) (formula: $C_3H_7O_7P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	1
			11	3	7	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	K	0	0
			2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	352	Total 352	O 352	0	0
7	B	353	Total 353	O 353	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: enolase 1



• Molecule 1: enolase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.00Å 65.00Å 85.90Å 90.00° 99.50° 90.00°	Depositor
Resolution (Å)	10.00 – 1.50 19.75 – 1.50	Depositor EDS
% Data completeness (in resolution range)	5.0 (10.00-1.50) 96.8 (19.75-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.71 (at 1.50Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.153 , 0.217 0.190 , 0.189	Depositor DCC
R_{free} test set	6042 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	12.3	Xtriage
Anisotropy	0.410	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 66.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7298	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PEP, K, 2PG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	1/3352 (0.0%)	0.45	0/4534
1	B	0.27	1/3321 (0.0%)	0.43	0/4491
All	All	0.28	2/6673 (0.0%)	0.44	0/9025

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	436	LEU	C-OXT	7.34	1.37	1.23
1	B	436	LEU	C-OXT	-6.76	1.10	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3292	0	3301	47	0
1	B	3263	0	3275	41	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	10	0	2	0	0
3	B	10	0	2	0	0
4	A	11	0	4	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	1	0	0	0	0
6	B	2	0	0	0	0
7	A	352	0	0	1	0
7	B	353	0	0	7	0
All	All	7298	0	6584	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:HIS:HA	1:B:286:MET:HE3	1.57	0.87
1:B:140:LYS:HE2	1:B:391:ARG:NH2	2.04	0.72
1:B:278:LEU:HD12	1:B:281:LEU:HD22	1.71	0.72
1:B:57:MET:HB2	1:B:59:LYS:HZ2	1.61	0.66
1:A:140:LYS:HE2	1:A:142:SER:O	1.97	0.65
1:A:278:LEU:HD12	1:A:281:LEU:HD22	1.79	0.65
1:A:163:ALA:H	1:A:217:ASN:ND2	1.98	0.61
1:A:345:LYS:HZ3	4:A:441[B]:2PG:H2	1.66	0.60
1:A:257:LYS:HB3	1:A:273:LEU:O	2.04	0.58
1:B:45:ALA:HA	1:B:101:ASN:ND2	2.18	0.58
1:A:196:LEU:O	1:A:200:ARG:HG3	2.04	0.58
1:B:78:LYS:HD3	7:B:1255:HOH:O	2.04	0.57
1:A:269:LYS:HA	1:A:272:TRP:CD1	2.40	0.57
1:A:403:SER:HB2	1:B:404:GLU:HB3	1.88	0.56
1:A:269:LYS:HA	1:A:272:TRP:NE1	2.22	0.55
1:A:269:LYS:HZ2	1:A:272:TRP:HH2	1.53	0.55
1:B:297:PRO:HD2	1:B:306:TRP:CH2	2.41	0.54
1:A:157:GLY:HA2	1:A:263:LYS:HE2	1.88	0.54
1:B:146:LEU:HD13	1:B:416:GLU:HB2	1.89	0.54
1:B:131:LYS:HG2	1:B:144:TYR:OH	2.06	0.54
1:A:345:LYS:NZ	4:A:441[B]:2PG:H2	2.23	0.54
1:A:373:HIS:CG	1:A:397:THR:HA	2.43	0.54
1:A:404:GLU:HB3	1:B:403:SER:HB2	1.90	0.54
1:B:155:ASN:ND2	7:B:1216:HOH:O	2.44	0.51
1:A:73:ALA:HB3	1:A:74:PRO:HD3	1.93	0.51
1:A:155:ASN:ND2	1:A:213:GLY:HA3	2.26	0.51
1:A:140:LYS:HD3	1:A:391:ARG:NH2	2.25	0.51
1:A:264:ASN:O	1:A:267:SER:HB2	2.10	0.50
1:B:252:PHE:HB3	1:B:262:PHE:CD2	2.47	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:284:SER:CB	1:A:288:ARG:HH21	2.24	0.50
1:B:326:ASN:OD1	1:B:328:LYS:HE2	2.11	0.49
1:B:17:PRO:HG3	1:B:56:TRP:CD1	2.48	0.48
1:A:296:ASP:HA	1:A:306:TRP:CH2	2.49	0.48
1:B:274:THR:HG23	1:B:277:GLN:OE1	2.14	0.47
1:B:296:ASP:OD2	1:B:320:ASP:HB3	2.14	0.47
1:A:7:ALA:HB2	1:A:68:VAL:HG11	1.96	0.47
1:B:101:ASN:ND2	7:B:1601:HOH:O	2.47	0.47
1:B:281:LEU:HD23	1:B:282:TYR:N	2.30	0.47
1:A:140:LYS:HG2	1:A:142:SER:H	1.79	0.47
1:B:217:ASN:ND2	7:B:1308:HOH:O	2.47	0.47
1:B:308:HIS:HA	7:B:1683:HOH:O	2.15	0.47
1:B:278:LEU:CD1	1:B:281:LEU:HD22	2.41	0.46
1:A:15:GLY:O	1:B:191:HIS:HE1	1.99	0.46
1:A:362:SER:O	1:A:367:TRP:HB2	2.16	0.46
1:A:269:LYS:HA	1:A:272:TRP:CE2	2.51	0.46
1:A:296:ASP:OD2	1:A:320:ASP:HB3	2.16	0.45
1:B:296:ASP:HA	1:B:306:TRP:CH2	2.51	0.45
1:A:268:ASP:HB3	1:A:271:LYS:HD2	1.98	0.45
1:B:269:LYS:HA	1:B:272:TRP:CE2	2.51	0.45
1:B:66:LYS:NZ	1:B:70:ASP:OD2	2.50	0.45
1:B:163:ALA:HB3	1:B:219:GLN:HA	1.98	0.45
1:A:252:PHE:HB2	1:A:259:ASP:O	2.16	0.44
1:B:164:LEU:HD21	1:B:169:PHE:CE2	2.53	0.44
1:A:297:PRO:HD2	1:A:306:TRP:CH2	2.53	0.44
1:B:101:ASN:ND2	7:B:1194:HOH:O	2.50	0.44
1:B:128:PRO:HD2	1:B:131:LYS:HD2	2.00	0.44
1:B:101:ASN:HA	1:B:101:ASN:HD22	1.59	0.44
1:B:373:HIS:CG	1:B:397:THR:HA	2.53	0.44
1:A:278:LEU:HG	1:A:282:TYR:CE2	2.53	0.44
1:A:306:TRP:HB3	1:A:338:ALA:HB1	2.00	0.43
1:A:269:LYS:HD3	1:A:272:TRP:CH2	2.54	0.42
1:B:73:ALA:O	1:B:77:VAL:HG22	2.19	0.42
1:A:164:LEU:HD21	1:A:169:PHE:CE2	2.54	0.42
1:A:257:LYS:HG2	1:A:274:THR:HG22	2.01	0.42
1:A:161:GLY:H	1:B:207:ASN:HD21	1.65	0.42
1:A:44:GLU:HG2	1:A:348:GLN:HG2	2.01	0.42
1:B:276:PRO:HD3	7:B:1518:HOH:O	2.18	0.42
1:A:403:SER:CB	1:B:404:GLU:HB3	2.50	0.42
1:A:284:SER:OG	1:A:288:ARG:NH2	2.50	0.42
1:A:227:LEU:HD12	1:A:227:LEU:HA	1.85	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:325:THR:HG22	1:B:325:THR:O	2.19	0.42
1:B:249:SER:HA	1:B:252:PHE:CE2	2.55	0.42
1:B:256:GLY:O	1:B:257:LYS:HD3	2.20	0.42
1:B:57:MET:HB2	1:B:59:LYS:NZ	2.30	0.42
1:A:130:TYR:CE1	1:A:419:LEU:HD21	2.54	0.41
1:A:398:GLY:HA3	1:A:405:ARG:HD2	2.02	0.41
1:A:404:GLU:HB3	1:B:403:SER:CB	2.51	0.41
1:A:374:ARG:O	1:A:377:GLU:HG2	2.21	0.41
1:A:275:GLY:N	1:A:276:PRO:HD2	2.35	0.41
1:A:325:THR:O	1:A:325:THR:HG22	2.22	0.40
1:A:269:LYS:HG2	1:A:272:TRP:CZ2	2.57	0.40
1:A:345:LYS:NZ	7:A:1002:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/436 (100%)	423 (98%)	10 (2%)	1 (0%)	47	23
1	B	427/436 (98%)	419 (98%)	6 (1%)	2 (0%)	29	9
All	All	861/872 (99%)	842 (98%)	16 (2%)	3 (0%)	41	18

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	402	ARG
1	B	402	ARG
1	B	324	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/344 (100%)	339 (98%)	5 (2%)	65	39
1	B	342/344 (99%)	327 (96%)	15 (4%)	28	5
All	All	686/688 (100%)	666 (97%)	20 (3%)	42	13

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	217	ASN
1	A	281	LEU
1	A	347	ASN
1	A	373	HIS
1	A	406	LEU
1	B	80	ASN
1	B	101	ASN
1	B	142	SER
1	B	155	ASN
1	B	219	GLN
1	B	233	LYS
1	B	238	ASP
1	B	255	ASP
1	B	263	LYS
1	B	267	SER
1	B	281	LEU
1	B	287	LYS
1	B	347	ASN
1	B	373	HIS
1	B	392	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	ASN
1	A	126	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	152	ASN
1	A	167	GLN
1	A	217	ASN
1	A	219	GLN
1	A	422	ASN
1	B	80	ASN
1	B	101	ASN
1	B	126	ASN
1	B	152	ASN
1	B	155	ASN
1	B	207	ASN
1	B	217	ASN
1	B	219	GLN
1	B	422	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 7 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEP	A	440[A]	2	6,9,9	1.67	1 (16%)	8,13,13	1.03	1 (12%)
4	2PG	A	441[B]	2	7,10,10	1.17	1 (14%)	8,14,14	0.55	0
3	PEP	B	440	2	6,9,9	1.69	1 (16%)	8,13,13	0.95	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEP	A	440[A]	2	-	0/5/9/9	-
4	2PG	A	441[B]	2	-	0/7/11/11	-
3	PEP	B	440	2	-	0/5/9/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	440	PEP	C3-C2	3.31	1.39	1.33
3	A	440[A]	PEP	C3-C2	3.26	1.39	1.33
4	A	441[B]	2PG	O1P-C2	-2.05	1.43	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	440[A]	PEP	O2-C2-C3	-2.40	120.16	124.79
3	B	440	PEP	O2-C2-C3	-2.04	120.85	124.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	441[B]	2PG	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/436 (100%)	0.47	37 (8%) 10 11	8, 15, 42, 82	0
1	B	431/436 (98%)	0.31	16 (3%) 41 46	8, 15, 35, 58	0
All	All	867/872 (99%)	0.39	53 (6%) 21 23	8, 15, 36, 82	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	269	LYS	9.9
1	A	276	PRO	9.7
1	A	272	TRP	7.3
1	A	266	ASN	6.1
1	B	164	LEU	5.5
1	A	275	GLY	5.4
1	B	270	SER	4.8
1	B	163	ALA	4.7
1	A	274	THR	4.6
1	B	219	GLN	4.5
1	A	262	PHE	4.5
1	A	273	LEU	4.5
1	A	267	SER	4.3
1	A	265	PRO	4.2
1	A	268	ASP	4.2
1	A	264	ASN	4.0
1	A	164	LEU	4.0
1	B	271	LYS	3.7
1	B	255	ASP	3.7
1	B	272	TRP	3.7
1	A	260	LEU	3.6
1	A	271	LYS	3.6
1	A	335	GLU	3.5
1	A	281	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	42	VAL	3.4
1	B	39	SER	3.4
1	A	278	LEU	3.4
1	A	284	SER	3.2
1	A	277	GLN	3.2
1	A	219	GLN	3.1
1	A	258	TYR	3.0
1	A	279	ALA	2.8
1	A	280	ASP	2.8
1	B	268	ASP	2.8
1	B	262	PHE	2.7
1	A	257	LYS	2.7
1	A	161	GLY	2.7
1	B	269	LYS	2.6
1	A	263	LYS	2.6
1	A	203	ALA	2.5
1	B	261	ASP	2.5
1	B	267	SER	2.5
1	B	217	ASN	2.5
1	A	261	ASP	2.5
1	B	421	ASP	2.5
1	A	256	GLY	2.3
1	A	312	THR	2.3
1	A	160	ALA	2.2
1	A	40	THR	2.1
1	A	311	LYS	2.1
1	A	270	SER	2.1
1	A	288	ARG	2.1
1	B	142	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	B	439	1/1	0.89	0.12	35,35,35,35	0
2	MG	A	439	1/1	0.95	0.14	24,24,24,24	0
6	K	B	961	1/1	0.96	0.10	25,25,25,25	0
6	K	B	960	1/1	0.97	0.09	29,29,29,29	0
3	PEP	B	440	10/10	0.97	0.07	10,11,13,13	0
5	CL	B	950	1/1	0.98	0.06	15,15,15,15	0
2	MG	B	438	1/1	0.98	0.08	15,15,15,15	0
4	2PG	A	441[B]	11/11	0.98	0.07	10,11,12,14	11
2	MG	A	438	1/1	0.98	0.07	12,12,12,12	0
3	PEP	A	440[A]	10/10	0.98	0.07	9,11,12,14	10

6.5 Other polymers

There are no such residues in this entry.