



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 22, 2020 – 10:16 am GMT

PDB ID : 7AL4
Title : Ancestral Flavin-containing monooxygenase (FMO) 1 (mammalian)
Authors : Nicoll, C.R.; Mattevi, A.
Deposited on : 2020-10-05
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.16
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	531	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 77%, yellow 77%, yellow 89%, orange 89%, orange 97%, grey 97%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> % 77% 18% .. </div> </div>
1	B	531	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 4%, green 4%, green 79%, yellow 79%, yellow 87%, orange 87%, orange 95%, grey 95%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 4% 79% 18% .. </div> </div>
1	C	531	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 11%, green 11%, green 78%, yellow 78%, yellow 86%, orange 86%, orange 94%, grey 94%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> 11% 78% 18% .. </div> </div>
1	D	531	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, green 1%, green 76%, yellow 76%, yellow 84%, orange 84%, orange 92%, grey 92%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> % 76% 20% .. </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	LMT	A	603	-	-	-	X

2 Entry composition ⓘ


There are 7 unique types of molecules in this entry. The entry contains 17079 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ancestral Flavin-containing monooxygenase 1 (mammalian).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	524	Total	C	N	O	S	0	0	0
			4153	2693	687	748	25			
1	C	520	Total	C	N	O	S	0	0	0
			4081	2633	681	744	23			
1	B	523	Total	C	N	O	S	0	0	0
			4105	2650	685	747	23			
1	A	517	Total	C	N	O	S	0	0	0
			4071	2630	679	739	23			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is DODECYL-BETA-D-MALTOSE (three-letter code: LMT) (formula: C₂₄H₄₆O₁₁).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Cl	0	0
			1	1		
6	A	1	Total	Cl	0	0
			1	1		
6	D	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	16	Total	O	0	0
			16	16		
7	C	5	Total	O	0	0
			5	5		
7	B	8	Total	O	0	0
			8	8		
7	A	11	Total	O	0	0
			11	11		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ancestral Flavin-containing monooxygenase 1 (mammalian)




- Molecule 1: Ancestral Flavin-containing monooxygenase 1 (mammalian)



• Molecule 1: Ancestral Flavin-containing monooxygenase 1 (mammalian)



• Molecule 1: Ancestral Flavin-containing monooxygenase 1 (mammalian)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	115.92Å 92.45Å 156.69Å 90.00° 95.12° 90.00°	Depositor
Resolution (Å)	49.10 – 3.00 49.10 – 3.00	Depositor EDS
% Data completeness (in resolution range)	60.1 (49.10-3.00) 60.1 (49.10-3.00)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.228 , 0.264 0.228 , 0.265	Depositor DCC
R_{free} test set	817 reflections (2.05%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtriage
Anisotropy	(Not available)	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 61.0	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	17079	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: (*Not available*)

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAP, LMT, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/4176	0.76	2/5672 (0.0%)
1	B	0.68	0/4211	0.76	2/5722 (0.0%)
1	C	0.69	0/4185	0.75	3/5684 (0.1%)
1	D	0.68	0/4260	0.76	0/5786
All	All	0.68	0/16832	0.76	7/22864 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	517	PRO	N-CA-CB	6.55	111.16	103.30
1	C	515	PRO	N-CA-CB	5.98	110.47	103.30
1	B	528	PRO	N-CA-CB	5.82	110.28	103.30
1	C	513	PRO	N-CA-CB	5.72	110.17	103.30
1	C	526	PRO	N-CA-CB	5.69	110.13	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4071	0	4033	54	0
1	B	4105	0	4049	53	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	4081	0	4011	45	0
1	D	4153	0	4191	52	0
2	A	53	0	31	4	0
2	B	53	0	31	3	0
2	C	53	0	31	3	0
2	D	53	0	31	2	0
3	A	48	0	25	3	0
3	B	48	0	25	6	0
3	C	48	0	25	1	0
3	D	48	0	25	1	0
4	A	105	0	138	0	0
4	B	70	0	92	2	0
4	D	35	0	46	0	0
5	D	12	0	16	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	A	11	0	0	0	0
7	B	8	0	0	0	0
7	C	5	0	0	1	0
7	D	16	0	0	0	0
All	All	17079	0	16800	194	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:379:ILE:HB	1:A:380:PRO:HD3	1.66	0.77
1:D:379:ILE:HB	1:D:380:PRO:HD3	1.68	0.74
1:B:379:ILE:HB	1:B:380:PRO:HD3	1.69	0.73
1:C:379:ILE:HB	1:C:380:PRO:HD3	1.70	0.72
1:D:344:LYS:O	1:D:345:VAL:HG22	1.88	0.72

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	511/531 (96%)	442 (86%)	57 (11%)	12 (2%)	6	30
1	B	519/531 (98%)	445 (86%)	63 (12%)	11 (2%)	7	33
1	C	512/531 (96%)	437 (85%)	61 (12%)	14 (3%)	5	26
1	D	520/531 (98%)	438 (84%)	69 (13%)	13 (2%)	5	28
All	All	2062/2124 (97%)	1762 (86%)	250 (12%)	50 (2%)	6	29

5 of 50 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	66	MET
1	D	422	LEU
1	D	425	CYS
1	D	514	LEU
1	C	66	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/469 (94%)	402 (91%)	41 (9%)	9	33
1	B	444/469 (95%)	407 (92%)	37 (8%)	11	39
1	C	441/469 (94%)	401 (91%)	40 (9%)	9	34
1	D	461/469 (98%)	415 (90%)	46 (10%)	7	29
All	All	1789/1876 (95%)	1625 (91%)	164 (9%)	9	34

5 of 164 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	412	ARG
1	B	147	THR
1	A	402	GLN
1	C	426	LYS
1	C	462	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 44 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	347	ASN
1	B	84	ASN
1	A	347	ASN
1	C	472	GLN
1	C	491	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 3 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	B	602	-	45,52,52	0.90	1 (2%)	56,80,80	1.37	8 (14%)
3	NAP	A	602	-	45,52,52	0.89	2 (4%)	56,80,80	1.50	9 (16%)
3	NAP	D	602	-	45,52,52	0.88	2 (4%)	56,80,80	1.25	7 (12%)
2	FAD	D	601	-	51,58,58	1.10	2 (3%)	60,89,89	1.70	6 (10%)
4	LMT	A	603	-	36,36,36	0.76	1 (2%)	47,47,47	1.36	7 (14%)
5	GOL	D	606	-	5,5,5	0.14	0	5,5,5	0.37	0
5	GOL	D	604	-	5,5,5	0.21	0	5,5,5	0.57	0
4	LMT	D	603	-	36,36,36	0.77	1 (2%)	47,47,47	1.11	4 (8%)
4	LMT	B	604	-	36,36,36	0.73	1 (2%)	47,47,47	1.41	6 (12%)
4	LMT	A	605	-	36,36,36	0.67	1 (2%)	47,47,47	0.88	2 (4%)
2	FAD	B	601	-	51,58,58	1.09	2 (3%)	60,89,89	1.78	7 (11%)
4	LMT	B	603	-	36,36,36	0.63	1 (2%)	47,47,47	0.88	2 (4%)
2	FAD	C	601	-	51,58,58	1.11	2 (3%)	60,89,89	1.74	7 (11%)
4	LMT	A	606	-	36,36,36	0.73	1 (2%)	47,47,47	1.60	5 (10%)
2	FAD	A	601	-	51,58,58	1.12	3 (5%)	60,89,89	1.78	9 (15%)
3	NAP	C	602	-	45,52,52	0.90	3 (6%)	56,80,80	1.20	6 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	B	602	-	-	9/31/67/67	0/5/5/5
3	NAP	A	602	-	-	10/31/67/67	0/5/5/5
3	NAP	D	602	-	-	14/31/67/67	0/5/5/5
2	FAD	D	601	-	-	12/30/50/50	0/6/6/6
4	LMT	A	603	-	-	11/21/61/61	0/2/2/2
5	GOL	D	606	-	-	0/4/4/4	-
5	GOL	D	604	-	-	2/4/4/4	-
4	LMT	D	603	-	-	8/21/61/61	0/2/2/2
4	LMT	B	604	-	-	10/21/61/61	0/2/2/2
4	LMT	A	605	-	-	10/21/61/61	0/2/2/2
2	FAD	B	601	-	-	12/30/50/50	0/6/6/6
4	LMT	B	603	-	-	9/21/61/61	0/2/2/2
2	FAD	C	601	-	-	12/30/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LMT	A	606	-	-	12/21/61/61	0/2/2/2
2	FAD	A	601	-	-	12/30/50/50	0/6/6/6
3	NAP	C	602	-	-	13/31/67/67	0/5/5/5

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	601	FAD	C4X-C10	5.85	1.44	1.38
2	D	601	FAD	C4X-C10	5.62	1.44	1.38
2	A	601	FAD	C4X-C10	5.49	1.44	1.38
2	B	601	FAD	C4X-C10	5.31	1.44	1.38
4	A	606	LMT	O1'-C1'	2.88	1.45	1.40

The worst 5 of 85 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	FAD	C4-N3-C2	8.00	121.90	115.14
2	C	601	FAD	C4-N3-C2	7.93	121.83	115.14
2	A	601	FAD	C4-N3-C2	7.88	121.80	115.14
2	D	601	FAD	C4-N3-C2	7.77	121.70	115.14
4	A	606	LMT	O1'-C1'-C2'	7.03	119.28	108.30

There are no chirality outliers.

5 of 156 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	602	NAP	C5B-O5B-PA-O1A
3	B	602	NAP	C2N-C3N-C7N-O7N
3	B	602	NAP	C2N-C3N-C7N-N7N
3	A	602	NAP	O4B-C4B-C5B-O5B
3	A	602	NAP	C5D-O5D-PN-O3

There are no ring outliers.

9 monomers are involved in 20 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	602	NAP	6	0
3	A	602	NAP	3	0
3	D	602	NAP	1	0
2	D	601	FAD	2	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	604	LMT	2	0
2	B	601	FAD	3	0
2	C	601	FAD	3	0
2	A	601	FAD	4	0
3	C	602	NAP	1	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.















Ligand FAD B 601




Ligand LMT B 603









5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2
1	B	1
1	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	505:GLN	C	513:GLU	N	16.04
1	B	506:GLU	C	514:SER	N	15.75
1	C	506:GLU	C	512:SER	N	11.59

Continued on next page...

Continued from previous page...

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	521:LEU	C	527:LEU	N	11.20

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	517/531 (97%)	-0.31	6 (1%) 79 54	28, 61, 116, 167	0
1	B	523/531 (98%)	-0.10	21 (4%) 38 15	38, 83, 144, 190	0
1	C	520/531 (97%)	0.51	61 (11%) 4 1	62, 136, 181, 217	0
1	D	524/531 (98%)	-0.35	6 (1%) 80 56	26, 56, 116, 222	0
All	All	2084/2124 (98%)	-0.06	94 (4%) 33 12	26, 77, 162, 222	0

The worst 5 of 94 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	277	THR	6.0
1	C	340	GLU	4.9
1	A	277	THR	4.8
1	C	341	SER	4.4
1	C	142	ALA	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	LMT	A	605	35/35	0.68	0.29	109,147,183,190	0
4	LMT	A	603	35/35	0.69	0.42	95,139,166,169	0
4	LMT	B	603	35/35	0.72	0.29	84,146,161,163	0
4	LMT	D	603	35/35	0.75	0.27	88,129,141,146	0
4	LMT	A	606	35/35	0.75	0.24	96,153,175,179	0
4	LMT	B	604	35/35	0.76	0.36	103,134,154,157	0
5	GOL	D	606	6/6	0.84	0.40	89,93,96,96	0
3	NAP	C	602	48/48	0.84	0.22	121,138,148,151	0
2	FAD	C	601	53/53	0.92	0.20	101,125,144,148	0
6	CL	A	604	1/1	0.92	0.14	84,84,84,84	0
3	NAP	A	602	48/48	0.93	0.19	48,65,106,117	0
2	FAD	B	601	53/53	0.94	0.17	44,52,71,82	0
3	NAP	B	602	48/48	0.94	0.14	73,89,115,121	0
3	NAP	D	602	48/48	0.94	0.17	65,86,92,98	0
5	GOL	D	604	6/6	0.95	0.14	43,45,46,46	0
6	CL	B	605	1/1	0.96	0.07	56,56,56,56	0
2	FAD	A	601	53/53	0.96	0.17	33,42,56,61	0
6	CL	D	605	1/1	0.98	0.10	43,43,43,43	0
2	FAD	D	601	53/53	0.98	0.17	32,36,43,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LMT A 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT A 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT B 603:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT D 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LMT A 606:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LMT B 604:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP C 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD C 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP D 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD D 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.