



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 01:20 pm BST

PDB ID : 2AMO  
Title : Loose Dimer of a Bacillus subtilis Nitric Oxide Synthase  
Authors : Pant, K.; Crane, B.R.  
Deposited on : 2005-08-09  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	362	
1	B	362	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6103 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase oxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	S	0	0	0
			2925	1863	502	552	8			
1	B	362	Total	C	N	O	S	0	0	0
			2812	1785	490	531	6			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	INSERTION	UNP O34453
A	-1	HIS	-	INSERTION	UNP O34453
A	0	ALA	-	INSERTION	UNP O34453
A	1	ALA	-	INSERTION	UNP O34453
A	2	ILE	-	INSERTION	UNP O34453
A	3	LEU	-	INSERTION	UNP O34453
A	4	TRP	-	INSERTION	UNP O34453
A	5	ASN	-	INSERTION	UNP O34453
A	6	GLU	-	INSERTION	UNP O34453
A	7	ALA	-	INSERTION	UNP O34453
A	8	LYS	-	INSERTION	UNP O34453
A	9	ALA	-	INSERTION	UNP O34453
A	10	PHE	-	INSERTION	UNP O34453
A	11	ILE	-	INSERTION	UNP O34453
A	12	ALA	-	INSERTION	UNP O34453
A	13	GLU	-	INSERTION	UNP O34453
A	14	CYS	-	INSERTION	UNP O34453
A	15	TYR	-	INSERTION	UNP O34453
A	16	ALA	-	INSERTION	UNP O34453
A	17	GLU	-	INSERTION	UNP O34453
A	18	LEU	-	INSERTION	UNP O34453
A	19	GLY	-	INSERTION	UNP O34453
A	20	LYS	-	INSERTION	UNP O34453
A	21	ALA	-	INSERTION	UNP O34453
A	22	GLU	-	INSERTION	UNP O34453

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	GLU	-	INSERTION	UNP O34453
A	24	VAL	-	INSERTION	UNP O34453
A	25	ALA	-	INSERTION	UNP O34453
A	29	ASP	ALA	CONFLICT	UNP O34453
A	30	SER	ASP	CONFLICT	UNP O34453
A	84	ASP	GLU	CONFLICT	UNP O34453
A	105	SER	THR	CONFLICT	UNP O34453
A	134	ALA	SER	CONFLICT	UNP O34453
A	135	ALA	ASP	CONFLICT	UNP O34453
A	138	ALA	ARG	CONFLICT	UNP O34453
A	153	GLN	GLU	CONFLICT	UNP O34453
A	264	SER	ALA	CONFLICT	UNP O34453
A	265	THR	ALA	CONFLICT	UNP O34453
A	266	ASN	ASP	CONFLICT	UNP O34453
A	286	TYR	HIS	CONFLICT	UNP O34453
B	-2	SER	-	INSERTION	UNP O34453
B	-1	HIS	-	INSERTION	UNP O34453
B	0	ALA	-	INSERTION	UNP O34453
B	1	ALA	-	INSERTION	UNP O34453
B	2	ILE	-	INSERTION	UNP O34453
B	3	LEU	-	INSERTION	UNP O34453
B	4	TRP	-	INSERTION	UNP O34453
B	5	ASN	-	INSERTION	UNP O34453
B	6	GLU	-	INSERTION	UNP O34453
B	7	ALA	-	INSERTION	UNP O34453
B	8	LYS	-	INSERTION	UNP O34453
B	9	ALA	-	INSERTION	UNP O34453
B	10	PHE	-	INSERTION	UNP O34453
B	11	ILE	-	INSERTION	UNP O34453
B	12	ALA	-	INSERTION	UNP O34453
B	13	GLU	-	INSERTION	UNP O34453
B	14	CYS	-	INSERTION	UNP O34453
B	15	TYR	-	INSERTION	UNP O34453
B	16	ALA	-	INSERTION	UNP O34453
B	17	GLU	-	INSERTION	UNP O34453
B	18	LEU	-	INSERTION	UNP O34453
B	19	GLY	-	INSERTION	UNP O34453
B	20	LYS	-	INSERTION	UNP O34453
B	21	ALA	-	INSERTION	UNP O34453
B	22	GLU	-	INSERTION	UNP O34453
B	23	GLU	-	INSERTION	UNP O34453
B	24	VAL	-	INSERTION	UNP O34453

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	25	ALA	-	INSERTION	UNP O34453
B	29	ASP	ALA	CONFLICT	UNP O34453
B	30	SER	ASP	CONFLICT	UNP O34453
B	84	ASP	GLU	CONFLICT	UNP O34453
B	105	SER	THR	CONFLICT	UNP O34453
B	134	ALA	SER	CONFLICT	UNP O34453
B	135	ALA	ASP	CONFLICT	UNP O34453
B	138	ALA	ARG	CONFLICT	UNP O34453
B	153	GLN	GLU	CONFLICT	UNP O34453
B	264	SER	ALA	CONFLICT	UNP O34453
B	265	THR	ALA	CONFLICT	UNP O34453
B	266	ASN	ASP	CONFLICT	UNP O34453
B	286	TYR	HIS	CONFLICT	UNP O34453

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	158	Total 158	O 158	0	0
3	B	122	Total 122	O 122	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Nitric oxide synthase oxygenase



- Molecule 1: Nitric oxide synthase oxygenase



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.31Å 93.21Å 118.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.62 – 2.60	Depositor
% Data completeness (in resolution range)	95.0 (29.62-2.60)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REFMAC 5.1.9999	Depositor
R, $R_{free}$	0.274 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6103	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	1/3002 (0.0%)	0.63	0/4073
1	B	0.33	0/2886	0.61	3/3929 (0.1%)
All	All	0.40	1/5888 (0.0%)	0.62	3/8002 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	359	GLU	C-OXT	-10.41	1.03	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	336	HIS	N-CA-C	6.77	129.28	111.00
1	B	45	LYS	N-CA-C	6.38	128.22	111.00
1	B	73	VAL	N-CA-C	-5.50	96.16	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2925	0	2828	59	0
1	B	2812	0	2623	95	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	43	0	30	1	0
2	B	43	0	30	2	0
3	A	158	0	0	8	0
3	B	122	0	0	3	0
All	All	6103	0	5511	152	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 152 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:ASN:HD21	1:A:169:ILE:H	1.16	0.92
1:B:63:ILE:HG13	1:B:326:LEU:HG	1.54	0.88
1:B:44:THR:HG22	1:B:45:LYS:H	1.42	0.84
1:A:56:TRP:CE3	1:A:68:TRP:HA	2.13	0.84
1:B:57:ARG:HD2	1:B:317:ARG:HH21	1.41	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/362 (99%)	339 (94%)	20 (6%)	1 (0%)	41	64
1	B	360/362 (99%)	292 (81%)	55 (15%)	13 (4%)	3	4
All	All	720/724 (99%)	631 (88%)	75 (10%)	14 (2%)	8	15

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	71	LEU
1	B	265	THR
1	B	322	ASP
1	B	330	ILE
1	B	334	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/308 (100%)	301 (98%)	6 (2%)	55	78
1	B	278/308 (90%)	259 (93%)	19 (7%)	16	32
All	All	585/616 (95%)	560 (96%)	25 (4%)	29	54

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	78	ASP
1	B	87	ASP
1	B	336	HIS
1	B	79	VAL
1	B	162	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	225	HIS
1	A	311	GLN
1	B	123	ASN
1	A	124	HIS
1	B	98	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	1901	1	27,50,50	1.86	7 (25%)	17,82,82	1.41	3 (17%)
2	HEM	B	1902	1	27,50,50	1.54	5 (18%)	17,82,82	1.44	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1901	1	-	1/6/54/54	-
2	HEM	B	1902	1	-	0/6/54/54	-

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1901	HEM	C3B-CAB	-5.66	1.36	1.47
2	B	1902	HEM	C3B-CAB	-4.21	1.39	1.47
2	A	1901	HEM	C3B-C2B	-3.55	1.35	1.40
2	B	1902	HEM	C3C-CAC	-3.47	1.40	1.47
2	A	1901	HEM	C4D-C3D	2.99	1.49	1.42

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	1902	HEM	CMC-C2C-C3C	3.28	130.82	124.68
2	A	1901	HEM	CBD-CAD-C3D	-3.17	106.65	112.48
2	B	1902	HEM	CBA-CAA-C2A	3.04	118.08	112.49
2	A	1901	HEM	C4C-C3C-C2C	-2.31	105.28	106.90
2	A	1901	HEM	C1D-C2D-C3D	-2.08	105.55	107.00

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1901	HEM	C1A-C2A-CAA-CBA


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1901	HEM	1	0
2	B	1902	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.