



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 7, 2020 – 02:27 AM BST

PDB ID : 6ANV
Title : Crystal structure of anti-CRISPR protein AcrF1
Authors : Yang, H.; Patel, D.J.
Deposited on : 2017-08-14
Resolution : 2.27 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 88% 11% </div> </div>
1	B	452	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 87% 13% </div> </div>
2	C	4	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 50%, yellow 50%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 50% 50% </div> </div>
2	D	4	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 50%, yellow 50%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 50% 50% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PEG	B	511	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7488 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti-CRISPR protein AcrF1 fused with C-terminal MBP tag.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	451	Total	C	N	O	S	0	0	0
			3507	2246	579	673	9			
1	B	451	Total	C	N	O	S	0	0	0
			3507	2246	579	673	9			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP L7P7M1
A	0	SER	-	expression tag	UNP L7P7M1
A	79	LEU	-	linker	UNP L7P7M1
A	80	GLU	-	linker	UNP L7P7M1
A	447	ASN	ARG	engineered mutation	UNP P0AEY0
A	448	ALA	ILE	engineered mutation	UNP P0AEY0
A	449	ALA	THR	engineered mutation	UNP P0AEY0
A	450	ALA	LYS	engineered mutation	UNP P0AEY0
B	-1	GLY	-	expression tag	UNP L7P7M1
B	0	SER	-	expression tag	UNP L7P7M1
B	79	LEU	-	linker	UNP L7P7M1
B	80	GLU	-	linker	UNP L7P7M1
B	447	ASN	ARG	engineered mutation	UNP P0AEY0
B	448	ALA	ILE	engineered mutation	UNP P0AEY0
B	449	ALA	THR	engineered mutation	UNP P0AEY0
B	450	ALA	LYS	engineered mutation	UNP P0AEY0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	4	Total	C	O	0	0	0
			45	24	21			
2	D	4	Total	C	O	0	0	0
			45	24	21			

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	127	Total	O	0	0
			127	127		
7	B	126	Total	O	0	0
			126	126		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: anti-CRISPR protein AcrF1 fused with C-terminal MBP tag



- Molecule 1: anti-CRISPR protein AcrF1 fused with C-terminal MBP tag



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.39Å 63.24Å 96.73Å 90.00° 93.21° 90.00°	Depositor
Resolution (Å)	45.87 – 2.27 45.87 – 2.26	Depositor EDS
% Data completeness (in resolution range)	96.4 (45.87-2.27) 96.4 (45.87-2.26)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.27Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.165 , 0.223 0.165 , 0.223	Depositor DCC
R_{free} test set	2310 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7488	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, PGE, PEG, EDO, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3588	0.55	0/4864
1	B	0.44	1/3588 (0.0%)	0.56	0/4864
All	All	0.42	1/7176 (0.0%)	0.55	0/9728

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	256	TYR	CE1-CZ	-5.21	1.31	1.38

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3507	0	3465	33	0
1	B	3507	0	3465	38	1
2	C	45	0	39	0	0
2	D	45	0	39	0	0
3	A	12	0	12	2	0
3	B	12	0	12	0	0
4	A	10	0	14	0	0
4	B	20	0	28	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	12	0	18	0	0
5	B	16	0	24	0	0
6	A	28	0	40	2	0
6	B	21	0	30	6	0
7	A	127	0	0	5	1
7	B	126	0	0	3	0
All	All	7488	0	7186	71	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:415:GLN:OE1	7:A:601:HOH:O	1.88	0.90
1:B:190:VAL:O	7:B:601:HOH:O	2.12	0.67
1:A:358:GLU:OE2	1:A:362:ASN:ND2	2.31	0.64
1:A:398:ALA:O	1:A:402:GLU:HG3	1.98	0.63
1:B:411:PRO:HB2	1:B:416:MET:HE3	1.81	0.63

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:48:GLU:OE2	7:A:698:HOH:O 2_356	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	449/452 (99%)	440 (98%)	9 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	449/452 (99%)	438 (98%)	11 (2%)	0	100	100
All	All	898/904 (99%)	878 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/363 (100%)	361 (99%)	2 (1%)	86	91
1	B	363/363 (100%)	356 (98%)	7 (2%)	57	66
All	All	726/726 (100%)	717 (99%)	9 (1%)	71	80

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	109	LYS
1	B	417	SER
1	B	276	VAL
1	B	50	LEU
1	B	259	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	C	1	2	12,12,12	0.39	0	17,17,17	0.77	0
2	GLC	C	2	2	11,11,12	0.95	0	15,15,17	0.97	0
2	GLC	C	3	2	11,11,12	0.65	0	15,15,17	1.18	1 (6%)
2	GLC	C	4	2	11,11,12	0.85	0	15,15,17	1.40	3 (20%)
2	GLC	D	1	2	12,12,12	0.61	0	17,17,17	0.97	1 (5%)
2	GLC	D	2	2	11,11,12	0.50	0	15,15,17	0.85	0
2	GLC	D	3	2	11,11,12	0.83	0	15,15,17	1.07	1 (6%)
2	GLC	D	4	2	11,11,12	0.88	0	15,15,17	1.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	C	3	2	-	0/2/19/22	0/1/1/1
2	GLC	C	4	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	4	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3	GLC	C6-C5-C4	-3.14	105.66	113.00
2	C	4	GLC	C2-C3-C4	-2.66	106.30	110.89
2	D	3	GLC	C6-C5-C4	-2.42	107.33	113.00
2	C	4	GLC	C1-O5-C5	2.35	115.38	112.19
2	D	1	GLC	C6-C5-C4	-2.14	108.00	113.00


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PEG	B	509	-	6,6,6	0.45	0	5,5,5	0.43	0
5	EDO	B	508	-	3,3,3	0.48	0	2,2,2	0.45	0
4	PGE	B	503	-	9,9,9	0.35	0	8,8,8	0.32	0
5	EDO	B	506	-	3,3,3	0.48	0	2,2,2	0.21	0
4	PGE	A	503	-	9,9,9	0.37	0	8,8,8	0.31	0
5	EDO	B	505	-	3,3,3	0.46	0	2,2,2	0.20	0
6	PEG	A	507	-	6,6,6	0.51	0	5,5,5	0.23	0
3	MES	B	502	-	12,12,12	1.99	1 (8%)	14,16,16	1.95	4 (28%)
3	MES	A	502	-	12,12,12	1.94	1 (8%)	14,16,16	2.27	6 (42%)
5	EDO	A	504	-	3,3,3	0.45	0	2,2,2	0.26	0
6	PEG	A	509	-	6,6,6	0.46	0	5,5,5	0.31	0
5	EDO	A	505	-	3,3,3	0.47	0	2,2,2	0.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PEG	B	510	-	6,6,6	0.46	0	5,5,5	0.33	0
5	EDO	A	506	-	3,3,3	0.48	0	2,2,2	0.21	0
5	EDO	B	507	-	3,3,3	0.48	0	2,2,2	0.28	0
6	PEG	A	510	-	6,6,6	0.46	0	5,5,5	0.23	0
6	PEG	A	508	-	6,6,6	0.48	0	5,5,5	0.24	0
4	PGE	B	504	-	9,9,9	0.32	0	8,8,8	0.29	0
6	PEG	B	511	-	6,6,6	0.47	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	B	509	-	-	2/4/4/4	-
5	EDO	B	508	-	-	1/1/1/1	-
4	PGE	B	503	-	-	2/7/7/7	-
5	EDO	B	506	-	-	1/1/1/1	-
4	PGE	A	503	-	-	2/7/7/7	-
5	EDO	B	505	-	-	0/1/1/1	-
6	PEG	A	507	-	-	0/4/4/4	-
3	MES	B	502	-	-	4/6/14/14	0/1/1/1
3	MES	A	502	-	-	1/6/14/14	0/1/1/1
5	EDO	A	504	-	-	0/1/1/1	-
6	PEG	A	509	-	-	1/4/4/4	-
5	EDO	A	505	-	-	1/1/1/1	-
6	PEG	B	510	-	-	2/4/4/4	-
5	EDO	A	506	-	-	0/1/1/1	-
5	EDO	B	507	-	-	1/1/1/1	-
6	PEG	A	510	-	-	1/4/4/4	-
6	PEG	A	508	-	-	3/4/4/4	-
4	PGE	B	504	-	-	3/7/7/7	-
6	PEG	B	511	-	-	4/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	MES	C8-S	-6.64	1.68	1.77
3	A	502	MES	C8-S	-6.36	1.68	1.77

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	502	MES	C5-N4-C3	5.02	120.13	108.83
3	B	502	MES	O2S-S-C8	3.86	111.56	106.92
3	A	502	MES	C7-N4-C5	3.70	120.69	111.23
3	B	502	MES	C5-N4-C3	3.34	116.36	108.83
3	B	502	MES	C7-N4-C5	3.17	119.34	111.23

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	MES	C8-C7-N4-C5
3	B	502	MES	C7-C8-S-O1S
3	B	502	MES	C7-C8-S-O2S
3	A	502	MES	C8-C7-N4-C5
6	B	511	PEG	C1-C2-O2-C3

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	509	PEG	1	0
3	A	502	MES	2	0
6	A	509	PEG	1	0
6	B	510	PEG	1	0
6	A	510	PEG	1	0
4	B	504	PGE	1	0
6	B	511	PEG	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	451/452 (99%)	-0.32	5 (1%) 80 82	15, 26, 46, 97	0
1	B	451/452 (99%)	-0.25	4 (0%) 84 85	14, 26, 48, 81	0
All	All	902/904 (99%)	-0.29	9 (0%) 82 84	14, 26, 47, 97	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	GLY	4.4
1	A	253	ASN	4.3
1	A	252	GLU	4.0
1	B	253	ASN	3.0
1	B	14	TYR	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	4	11/12	0.94	0.11	30,35,39,41	0
2	GLC	C	4	11/12	0.95	0.12	25,34,41,45	0
2	GLC	D	3	11/12	0.96	0.11	18,21,23,24	0
2	GLC	D	1	12/12	0.97	0.11	16,23,27,28	0
2	GLC	C	3	11/12	0.97	0.14	16,21,27,27	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	C	1	12/12	0.98	0.11	18,21,25,27	0
2	GLC	D	2	11/12	0.98	0.10	15,19,23,25	0
2	GLC	C	2	11/12	0.98	0.14	16,18,24,25	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	B	508	4/4	0.76	0.18	47,47,52,56	0
5	EDO	A	504	4/4	0.80	0.16	41,47,51,52	0
5	EDO	A	506	4/4	0.83	0.20	38,47,48,52	0
6	PEG	B	511	7/7	0.84	0.14	45,49,54,54	0
5	EDO	B	505	4/4	0.85	0.16	30,35,41,43	0
6	PEG	A	509	7/7	0.86	0.16	47,51,53,53	0
5	EDO	A	505	4/4	0.86	0.14	50,50,52,55	0
4	PGE	B	504	10/10	0.87	0.23	38,49,56,58	0
6	PEG	B	510	7/7	0.87	0.17	37,41,47,51	0
6	PEG	A	510	7/7	0.88	0.22	42,46,59,60	0
6	PEG	B	509	7/7	0.91	0.10	37,51,54,60	0
5	EDO	B	506	4/4	0.91	0.15	35,37,44,47	0
4	PGE	A	503	10/10	0.91	0.16	28,34,38,43	0
5	EDO	B	507	4/4	0.92	0.10	47,51,51,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	PEG	A	508	7/7	0.92	0.14	31,37,42,45	0
4	PGE	B	503	10/10	0.94	0.14	20,32,46,47	0
3	MES	B	502	12/12	0.94	0.17	22,33,59,61	0
6	PEG	A	507	7/7	0.95	0.10	38,40,47,54	0
3	MES	A	502	12/12	0.96	0.14	18,28,46,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.