



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2021 – 09:09 AM EDT

PDB ID : 2AST
Title : Crystal structure of Skp1-Skp2-Cks1 in complex with a p27 peptide
Authors : Hao, B.; Zhang, N.; Schulman, B.A.; Wu, G.; Pagano, M.; Pavletich, N.P.
Deposited on : 2005-08-24
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	<div> <div>21%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>11%</div> </div> </div>
2	B	336	<div> <div>3%</div> <div> <div></div> <div>85%</div> <div>10%</div> </div> </div>
3	C	69	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>20%</div> </div> </div>
4	D	10	<div> <div>60%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-phase kinase-associated protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	142	Total	C	N	O	S	0	0	0
			1129	716	183	224	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1002	ALA	PRO	engineered mutation	UNP P63208

- Molecule 2 is a protein called S-phase kinase-associated protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	325	Total	C	N	O	S	0	0	0
			2572	1635	443	478	16			

- Molecule 3 is a protein called Cyclin-dependent kinases regulatory subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	69	Total	C	N	O	S	0	0	0
			599	387	102	107	3			

- Molecule 4 is a protein called Cyclin-dependent kinase inhibitor 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	10	Total	C	N	O	P	0	0	0
			77	44	13	19	1			

- Molecule 5 is BENZAMIDINE (three-letter code: BEN) (formula: C₇H₈N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	N	0	0
			9	7	2		
5	C	1	Total	C	N	0	0
			9	7	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	8	Total	O	0	0
			8	8		
6	B	202	Total	O	0	0
			202	202		
6	C	55	Total	O	0	0
			55	55		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: S-phase kinase-associated protein 1A




- Molecule 2: S-phase kinase-associated protein 2



- Molecule 3: Cyclin-dependent kinases regulatory subunit 1



- Molecule 4: Cyclin-dependent kinase inhibitor 1B



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	148.70Å 148.70Å 98.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.30 19.80 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (20.00-2.30) 97.3 (19.80-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.202 , 0.221 0.196 , 0.195	Depositor DCC
R_{free} test set	2760 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.002	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4660	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BEN, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/1146	0.62	1/1547 (0.1%)
2	B	0.42	0/2622	0.65	6/3558 (0.2%)
3	C	0.36	0/619	0.59	1/837 (0.1%)
4	D	0.44	0/65	0.47	0/81
All	All	0.43	0/4452	0.63	8/6023 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2404	LYS	O-C-N	-10.32	106.19	122.70
2	B	2135	SER	N-CA-CB	8.29	122.94	110.50
2	B	2134	GLU	C-N-CA	7.37	140.13	121.70
1	A	1148	GLU	N-CA-CB	6.29	121.93	110.60
2	B	2295	LYS	N-CA-C	5.65	126.26	111.00
2	B	2344	ARG	NE-CZ-NH2	-5.48	117.56	120.30
2	B	2404	LYS	CA-C-N	5.21	128.66	117.20
3	C	3037	LEU	CA-CB-CG	-5.01	103.77	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1129	0	1124	7	0
2	B	2572	0	2607	24	0
3	C	599	0	574	9	0
4	D	77	0	74	0	0
5	B	9	0	7	0	0
5	C	9	0	7	0	0
6	A	8	0	0	0	0
6	B	202	0	0	5	0
6	C	55	0	0	1	0
All	All	4660	0	4393	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1037:ASP:C	1:A:1038:PRO:N	2.25	0.90
2:B:2403:ASN:O	2:B:2405:LYS:N	2.14	0.79
2:B:2185:HIS:HD2	2:B:2210:ASN:HD22	1.32	0.74
2:B:2277:ALA:O	2:B:2281:VAL:HG22	1.89	0.71
2:B:2149:PRO:HA	2:B:2152:THR:HG23	1.71	0.70
3:C:3058:MET:HE1	6:C:182:HOH:O	1.94	0.68
1:A:1107:ALA:HB2	1:A:1115:LEU:HD12	1.76	0.67
2:B:2152:THR:HG21	2:B:2170:MET:SD	2.34	0.66
2:B:2208:LEU:H	2:B:2229:ASN:HD22	1.45	0.65
1:A:1083:ASP:O	1:A:1084:ASP:HB2	1.98	0.62
2:B:2344:ARG:NH2	3:C:3045:ASN:OD1	2.22	0.62
2:B:2277:ALA:O	2:B:2281:VAL:CG2	2.48	0.61
2:B:2133:ASP:O	2:B:2134:GLU:HB2	2.01	0.60
2:B:2110:SER:HA	2:B:2415:ARG:HH21	1.70	0.56
1:A:1108:ASN:HB2	2:B:2104:LEU:HD11	1.90	0.53
1:A:1107:ALA:HB2	1:A:1115:LEU:CD1	2.39	0.52
2:B:2185:HIS:CD2	2:B:2210:ASN:HD22	2.20	0.51
3:C:3057:TYR:CE2	3:C:3058:MET:CE	2.95	0.49
3:C:3057:TYR:CE2	3:C:3058:MET:HE3	2.48	0.48
2:B:2136:LEU:O	2:B:2415:ARG:NH1	2.45	0.48
3:C:3023:MET:HG2	3:C:3066:ILE:HG12	1.95	0.48
2:B:2167:ARG:NE	6:B:188:HOH:O	2.20	0.47
2:B:2167:ARG:NH2	6:B:188:HOH:O	2.44	0.46
2:B:2208:LEU:H	2:B:2229:ASN:ND2	2.11	0.45
2:B:2100:LEU:HA	2:B:2101:PRO:HD3	1.86	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:2306:VAL:CG1	2:B:2333:GLN:HB2	2.48	0.44
1:A:1146:THR:HG22	1:A:1147:GLU:H	1.83	0.44
3:C:3040:GLU:HG3	3:C:3043:TRP:CZ2	2.52	0.43
2:B:2134:GLU:HG2	6:B:241:HOH:O	2.17	0.43
3:C:3027:ASP:O	3:C:3030:LYS:HG2	2.19	0.43
2:B:2103:GLU:H	2:B:2103:GLU:CD	2.21	0.42
2:B:2415:ARG:NH2	6:B:86:HOH:O	2.51	0.42
2:B:2137:TRP:O	2:B:2160:VAL:HA	2.19	0.42
3:C:3020:ARG:HG2	3:C:3021:HIS:N	2.34	0.42
2:B:2228:LYS:HB2	2:B:2228:LYS:HE3	1.96	0.41
1:A:1062:ASP:HA	1:A:1063:PRO:HD3	1.93	0.41
2:B:2415:ARG:NH1	6:B:63:HOH:O	2.54	0.41
3:C:3057:TYR:CE2	3:C:3058:MET:HE2	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	136/159 (86%)	124 (91%)	8 (6%)	4 (3%)	4	3
2	B	323/336 (96%)	307 (95%)	11 (3%)	5 (2%)	10	10
3	C	67/69 (97%)	66 (98%)	1 (2%)	0	100	100
4	D	7/10 (70%)	6 (86%)	0	1 (14%)	0	0
All	All	533/574 (93%)	503 (94%)	20 (4%)	10 (2%)	8	7

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	2134	GLU
2	B	2404	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	2406	ASN
4	D	4183	SER
2	B	2295	LYS
2	B	2403	ASN
1	A	1084	ASP
1	A	1034	LEU
1	A	1083	ASP
1	A	1145	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	128/145 (88%)	122 (95%)	6 (5%)	26	37
2	B	301/311 (97%)	288 (96%)	13 (4%)	29	40
3	C	66/66 (100%)	62 (94%)	4 (6%)	18	25
4	D	7/7 (100%)	7 (100%)	0	100	100
All	All	502/529 (95%)	479 (95%)	23 (5%)	27	38

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1004	ILE
1	A	1045	ASN
1	A	1049	LEU
1	A	1083	ASP
1	A	1089	ASP
1	A	1145	PHE
2	B	2098	ASP
2	B	2146	ASN
2	B	2152	THR
2	B	2179	SER
2	B	2225	THR
2	B	2262	ASN
2	B	2281	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	2290	LEU
2	B	2302	LEU
2	B	2321	VAL
2	B	2354	LEU
2	B	2387	GLN
2	B	2415	ARG
3	C	3024	LEU
3	C	3034	LYS
3	C	3037	LEU
3	C	3050	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1045	ASN
1	A	1108	ASN
1	A	1125	ASN
2	B	2146	ASN
2	B	2185	HIS
2	B	2229	ASN
2	B	2249	GLN
2	B	2262	ASN
2	B	2366	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TPO	D	4187	4	8,10,11	0.70	0	10,14,16	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TPO	D	4187	4	-	1/9/11/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	4187	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	BEN	B	1002	-	9,9,9	0.78	0	7,11,11	0.75	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BEN	C	1001	-	9,9,9	0.90	1 (11%)	7,11,11	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BEN	B	1002	-	-	4/4/4/4	0/1/1/1
5	BEN	C	1001	-	-	0/4/4/4	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1001	BEN	C1-C	2.12	1.51	1.47

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1002	BEN	N2-C-C1-C2
5	B	1002	BEN	N2-C-C1-C6
5	B	1002	BEN	N1-C-C1-C2
5	B	1002	BEN	N1-C-C1-C6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	142/159 (89%)	1.15	33 (23%) 0 1	47, 83, 125, 135	0
2	B	325/336 (96%)	-0.08	10 (3%) 49 56	29, 42, 81, 127	0
3	C	69/69 (100%)	-0.06	1 (1%) 75 80	33, 45, 68, 78	0
4	D	9/10 (90%)	3.19	6 (66%) 0 0	57, 84, 100, 103	0
All	All	545/574 (94%)	0.29	50 (9%) 9 12	29, 50, 115, 135	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1082	THR	9.5
1	A	1064	PRO	8.2
4	D	4182	GLY	6.8
1	A	1063	PRO	6.3
2	B	2095	VAL	6.3
4	D	4181	ALA	5.8
2	B	2404	LYS	5.5
2	B	2403	ASN	5.4
1	A	1145	PHE	5.2
1	A	1083	ASP	5.1
4	D	4183	SER	4.9
2	B	2096	SER	4.8
1	A	1087	VAL	4.5
1	A	1147	GLU	4.4
1	A	1002	ALA	4.4
4	D	4189	LYS	4.2
1	A	1142	LYS	4.0
1	A	1133	GLU	3.9
1	A	1137	LYS	3.9
1	A	1033	ASP	3.9
4	D	4184	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1146	THR	3.7
4	D	4190	LYS	3.6
1	A	1148	GLU	3.5
1	A	1136	ARG	3.5
1	A	1061	ASP	3.4
1	A	1062	ASP	3.4
1	A	1084	ASP	3.4
2	B	2099	SER	3.3
1	A	1032	GLU	3.2
3	C	3015	GLU	3.1
2	B	2125	LYS	3.1
1	A	1130	LYS	3.0
2	B	2120	VAL	3.0
1	A	1149	GLU	2.9
1	A	1129	GLY	2.9
1	A	1101	PHE	2.9
2	B	2123	VAL	2.9
1	A	1034	LEU	2.8
1	A	1088	TRP	2.7
2	B	2419	GLN	2.6
1	A	1154	ARG	2.6
2	B	2405	LYS	2.5
1	A	1135	ILE	2.5
1	A	1091	GLU	2.4
1	A	1094	LYS	2.3
1	A	1052	VAL	2.3
1	A	1103	LEU	2.2
1	A	1158	GLN	2.0
1	A	1140	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	TPO	D	4187	11/12	0.98	0.08	53,54,58,60	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	BEN	C	1001	9/9	0.90	0.12	43,49,50,50	0
5	BEN	B	1002	9/9	0.98	0.09	37,38,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.