



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2020 – 06:59 AM BST

PDB ID : 3ASS
Title : Crystal structure of P domain from Norovirus Funabashi258 stain in the complex with Lewis-b
Authors : Kubota, T.; Kumagai, A.; Itoh, H.; Furukawa, S.; Narimatsu, H.; Wakita, T.; Ishii, K.; Takeda, N.; Someya, Y.; Shirato, H.
Deposited on : 2010-12-17
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	<div> <div>7%</div> <div>90%</div> <div>5%</div> <div>5%</div> </div>
1	B	326	<div> <div>9%</div> <div>86%</div> <div>7%</div> <div>7%</div> </div>
2	C	4	<div> <div>100%</div> </div>
2	D	4	<div> <div>100%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5218 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	1	0
			2387	1527	402	449	9			
1	B	302	Total	C	N	O	S	0	3	0
			2335	1496	394	436	9			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	216	GLY	-	expression tag	UNP Q8JW44
A	217	PRO	-	expression tag	UNP Q8JW44
A	218	LEU	-	expression tag	UNP Q8JW44
A	219	GLY	-	expression tag	UNP Q8JW44
A	220	SER	-	expression tag	UNP Q8JW44
B	216	GLY	-	expression tag	UNP Q8JW44
B	217	PRO	-	expression tag	UNP Q8JW44
B	218	LEU	-	expression tag	UNP Q8JW44
B	219	GLY	-	expression tag	UNP Q8JW44
B	220	SER	-	expression tag	UNP Q8JW44

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	4	Total	C	N	O	0	0	0
			45	26	1	18			
2	D	4	Total	C	N	O	0	0	0
			45	26	1	18			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is P-NITROPHENOL (three-letter code: NPO) (formula: C₆H₅NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			10	6	1	3		
4	B	1	Total	C	N	O	0	0
			10	6	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	202	Total	O	0	0
			202	202		
5	B	182	Total	O	0	0
			182	182		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Capsid protein




- Molecule 1: Capsid protein



- Molecule 2: alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: alpha-L-fucopyranose-(1-2)-beta-D-galactopyranose-(1-3)-[alpha-L-fucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	74.69 Å 74.69 Å 106.97 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	31.23 – 1.60 31.23 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (31.23-1.60) 99.8 (31.23-1.60)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.55 (at 1.60 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.185 , 0.202 0.184 , 0.200	Depositor DCC
R_{free} test set	4435 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.004	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 57.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.006 for -h,-k,l 0.026 for h,-h-k,-l 0.016 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5218	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NPO, GAL, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2463	0.53	0/3372
1	B	0.37	0/2415	0.53	0/3307
All	All	0.36	0/4878	0.53	0/6679

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2387	0	2316	10	0
1	B	2335	0	2260	13	0
2	C	45	0	40	0	0
2	D	45	0	40	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	4	0	0
4	B	10	0	4	0	0
5	A	202	0	0	0	0
5	B	182	0	0	0	0
All	All	5218	0	4664	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:529:LYS:HE2	1:B:530:PRO:HD2	1.73	0.70
1:A:315:PHE:HA	1:A:363:HIS:CE1	2.27	0.69
1:B:332:HIS:HE1	1:B:388:SER:OG	1.78	0.67
1:B:254:MET:HG2	1:B:507:PRO:HB2	1.79	0.63
1:A:332:HIS:HE1	1:A:388:SER:OG	1.81	0.62
1:A:290:ARG:NH2	1:B:248:PRO:O	2.36	0.58
1:B:363[A]:HIS:CE1	1:B:414:ALA:H	2.21	0.58
1:A:440:ASN:HD22	1:A:446:ASN:ND2	2.06	0.54
1:B:440:ASN:HD22	1:B:446:ASN:ND2	2.06	0.53
1:B:363[B]:HIS:CD2	1:B:407:GLY:HA2	2.48	0.49
1:B:332:HIS:HD2	1:B:352:SER:OG	1.94	0.49
1:A:315:PHE:HA	1:A:363:HIS:HE1	1.74	0.47
1:A:363:HIS:CE1	1:A:414:ALA:H	2.34	0.46
1:B:400:LEU:HD13	1:B:448:VAL:HG22	1.98	0.46
1:A:248:PRO:O	1:B:290:ARG:NH2	2.49	0.45
1:A:470:ASP:OD2	1:A:524:ARG:HG2	2.15	0.45
1:B:489:LYS:HG3	1:B:528:LEU:HD21	2.00	0.43
1:A:273:ILE:HG12	1:A:492:PRO:HA	2.00	0.42
1:B:433:VAL:CG2	1:B:447:ASP:HB3	2.50	0.41
1:A:332:HIS:HD2	1:A:352:SER:OG	2.04	0.40
1:B:273:ILE:HG12	1:B:492:PRO:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/326 (95%)	308 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	303/326 (93%)	300 (99%)	3 (1%)	0	100	100
All	All	613/652 (94%)	608 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/277 (96%)	263 (99%)	3 (1%)	73	57
1	B	261/277 (94%)	259 (99%)	2 (1%)	81	70
All	All	527/554 (95%)	522 (99%)	5 (1%)	76	65

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	317	SER
1	A	390	PRO
1	A	524	ARG
1	B	317	SER
1	B	529	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	296	ASN
1	A	332	HIS
1	A	363	HIS
1	A	374	ASN
1	A	375	HIS
1	A	415	ASN
1	A	446	ASN
1	B	296	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	332	HIS
1	B	354	GLN
1	B	415	ASN
1	B	446	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,4	14,14,15	0.52	0	17,19,21	0.78	0
2	GAL	C	2	2	11,11,12	0.62	0	15,15,17	0.68	0
2	FUC	C	3	2	10,10,11	0.57	0	14,14,16	0.50	0
2	FUC	C	4	2	10,10,11	0.56	0	14,14,16	0.60	0
2	NAG	D	1	2,4	14,14,15	0.52	0	17,19,21	0.78	0
2	GAL	D	2	2	11,11,12	0.53	0	15,15,17	0.76	0
2	FUC	D	3	2	10,10,11	0.59	0	14,14,16	0.53	0
2	FUC	D	4	2	10,10,11	0.54	0	14,14,16	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,4	-	0/6/23/26	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1
2	FUC	C	3	2	-	-	0/1/1/1
2	FUC	C	4	2	-	-	0/1/1/1
2	NAG	D	1	2,4	-	0/6/23/26	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	FUC	D	3	2	-	-	0/1/1/1
2	FUC	D	4	2	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NPO	A	1001	2	9,10,10	0.76	0	11,13,13	0.53	0
4	NPO	B	1001	2	9,10,10	0.77	0	11,13,13	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NPO	A	1001	2	-	0/2/4/4	0/1/1/1
4	NPO	B	1001	2	-	2/2/4/4	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1001	NPO	C2-C1-N1-O3
4	B	1001	NPO	C6-C1-N1-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	311/326 (95%)	0.22	22 (7%) 16 14	11, 18, 32, 42	0
1	B	302/326 (92%)	0.22	28 (9%) 8 7	11, 18, 32, 37	0
All	All	613/652 (94%)	0.22	50 (8%) 11 10	11, 18, 32, 42	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315	PHE	14.1
1	A	410	LEU	13.8
1	B	410	LEU	10.5
1	B	503	VAL	10.4
1	B	525	PHE	10.3
1	A	314	ALA	9.8
1	A	409	SER	9.2
1	A	225	ILE	8.5
1	B	315	PHE	8.3
1	B	409	SER	7.9
1	A	411	SER	7.1
1	A	412	GLN	7.0
1	B	504	GLY	6.9
1	A	503	VAL	6.4
1	B	314	ALA	6.3
1	A	408	SER	6.3
1	B	425	PHE	6.1
1	A	316	ASP	5.8
1	B	313	MET	5.6
1	A	504	GLY	5.0
1	B	412	GLN	4.7
1	B	501	ASN	4.4
1	B	230	ARG	4.4
1	B	524	ARG	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	224	THR	4.1
1	A	414	ALA	4.1
1	A	413	ALA	4.1
1	B	411	SER	3.8
1	B	502	GLY	3.8
1	A	317	SER	3.4
1	B	480	ASP	3.1
1	B	316	ASP	3.1
1	A	533	THR	3.1
1	A	534	ALA	3.1
1	B	505	ALA	3.1
1	B	414	ALA	2.9
1	B	483	ARG	2.8
1	B	408	SER	2.8
1	B	481	THR	2.7
1	A	358	GLN	2.7
1	A	415	ASN	2.6
1	B	317	SER	2.4
1	A	229	THR	2.4
1	B	231	ALA	2.3
1	B	358	GLN	2.2
1	A	407	GLY	2.2
1	B	413	ALA	2.2
1	B	482	ASN	2.1
1	B	527	GLN	2.0
1	A	227	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FUC	C	4	10/11	0.83	0.11	21,22,24,24	0
2	FUC	D	4	10/11	0.85	0.14	21,22,24,24	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	C	1	14/15	0.86	0.12	22,23,24,25	0
2	FUC	D	3	10/11	0.89	0.12	23,26,26,28	0
2	FUC	C	3	10/11	0.89	0.08	22,22,25,25	0
2	GAL	C	2	11/12	0.90	0.07	21,21,22,23	0
2	NAG	D	1	14/15	0.91	0.15	22,25,27,28	0
2	GAL	D	2	11/12	0.93	0.09	18,20,22,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NPO	B	1001	10/10	0.71	0.34	33,37,39,40	0
4	NPO	A	1001	10/10	0.92	0.19	26,28,30,30	0
3	NA	A	701	1/1	0.94	0.10	33,33,33,33	0
3	NA	B	701	1/1	0.99	0.08	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.