



wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 05:33 pm BST

PDB ID : 4AS3
Title : Pseudomonas Aeruginosa Phosphorylcholine Phosphatase. Orthorhombic form
Authors : Infantes, L.; Otero, L.H.; Albert, A.
Deposited on : 2012-04-27
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	<div> <div>83%</div> <div>17%</div> <div>.</div> </div>
1	B	327	<div> <div>82%</div> <div>17%</div> <div>.</div> </div>
1	C	327	<div> <div>82%</div> <div>18%</div> </div>
1	D	327	<div> <div>84%</div> <div>14%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	B	1329	-	-	X	-
4	BTB	A	1330	-	-	X	-
4	BTB	A	1331	-	-	X	-
4	BTB	C	1331	-	-	X	-
4	BTB	D	1330	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHORYLCHOLINE PHOSPHATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	0	0
			2615	1674	439	490	12			
1	B	327	Total	C	N	O	S	0	0	0
			2615	1674	439	490	12			
1	C	327	Total	C	N	O	S	0	0	0
			2615	1674	439	490	12			
1	D	327	Total	C	N	O	S	0	0	0
			2615	1674	439	490	12			


- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		

- Molecule 4 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	115	Total 115	O 115	0	0
5	B	97	Total 97	O 97	0	0
5	C	87	Total 87	O 87	0	0
5	D	110	Total 110	O 110	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PHOSPHORYLCHOLINE PHOSPHATASE




• Molecule 1: PHOSPHORYLCHOLINE PHOSPHATASE



• Molecule 1: PHOSPHORYLCHOLINE PHOSPHATASE



• Molecule 1: PHOSPHORYLCHOLINE PHOSPHATASE





4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	165.34Å 69.59Å 119.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	119.60 – 2.40 50.05 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (119.60-2.40) 99.9 (50.05-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.11 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.177 , 0.249 0.173 , 0.243	Depositor DCC
R_{free} test set	2773 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	38.4	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 33.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11017	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.18 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.9039e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, BTB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.82	1/2680 (0.0%)	0.84	1/3643 (0.0%)
1	B	0.83	3/2680 (0.1%)	0.83	1/3643 (0.0%)
1	C	0.77	2/2680 (0.1%)	0.81	0/3643
1	D	0.80	3/2680 (0.1%)	0.85	2/3643 (0.1%)
All	All	0.80	9/10720 (0.1%)	0.83	4/14572 (0.0%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	238	TRP	CD2-CE2	5.88	1.48	1.41
1	A	253	TRP	CD2-CE2	5.53	1.48	1.41
1	D	97	TRP	CD2-CE2	5.50	1.48	1.41
1	C	318	TRP	CD2-CE2	5.48	1.48	1.41
1	B	318	TRP	CD2-CE2	5.47	1.48	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	149	ARG	NE-CZ-NH1	6.71	123.66	120.30
1	D	137	LEU	CA-CB-CG	6.43	130.10	115.30
1	A	137	LEU	CA-CB-CG	5.97	129.03	115.30
1	B	55	LEU	CA-CB-CG	5.23	127.33	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2615	0	2581	34	0
1	B	2615	0	2581	43	0
1	C	2615	0	2581	44	0
1	D	2615	0	2583	43	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	1	0
3	B	1	0	0	2	0
3	C	1	0	0	1	0
3	D	1	0	0	1	0
4	A	42	0	57	19	0
4	B	28	0	38	10	0
4	C	42	0	57	20	0
4	D	28	0	38	15	0
5	A	115	0	0	9	0
5	B	97	0	0	8	0
5	C	87	0	0	8	0
5	D	110	0	0	9	0
All	All	11017	0	10516	200	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 200 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:1331:BTB:H71	4:D:1331:BTB:O6	1.47	1.12
1:D:285:ASN:HB2	1:D:291:MET:CE	1.83	1.09
1:A:265:ASP:HB3	5:A:2083:HOH:O	1.52	1.05
1:B:291:MET:HE1	1:B:320:ILE:HG21	1.43	1.00
4:A:1331:BTB:H31	5:A:2115:HOH:O	1.66	0.96

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/327 (99%)	305 (94%)	20 (6%)	0	100	100
1	B	325/327 (99%)	310 (95%)	15 (5%)	0	100	100
1	C	325/327 (99%)	308 (95%)	16 (5%)	1 (0%)	41	55
1	D	325/327 (99%)	314 (97%)	11 (3%)	0	100	100
All	All	1300/1308 (99%)	1237 (95%)	62 (5%)	1 (0%)	51	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	276	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/274 (100%)	266 (97%)	8 (3%)	42	62
1	B	274/274 (100%)	262 (96%)	12 (4%)	28	45
1	C	274/274 (100%)	265 (97%)	9 (3%)	38	57
1	D	274/274 (100%)	267 (97%)	7 (3%)	46	66
All	All	1096/1096 (100%)	1060 (97%)	36 (3%)	38	57

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	136	THR
1	C	55	LEU
1	D	137	LEU
1	B	181	ARG
1	C	57	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	23	ASN
1	B	278	ASN
1	C	23	ASN
1	C	72	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 8 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	BTB	C	1332	-	13,13,13	1.15	0	7,16,16	0.92	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BTB	A	1332	-	13,13,13	1.17	0	7,16,16	0.59	0
4	BTB	D	1331	-	13,13,13	0.95	0	7,16,16	1.51	1 (14%)
4	BTB	A	1331	-	13,13,13	0.99	0	7,16,16	1.97	2 (28%)
4	BTB	C	1331	-	13,13,13	0.85	0	7,16,16	1.14	1 (14%)
4	BTB	D	1330	-	13,13,13	1.34	1 (7%)	7,16,16	2.94	1 (14%)
4	BTB	A	1330	-	13,13,13	1.65	1 (7%)	7,16,16	1.75	1 (14%)
4	BTB	C	1330	-	13,13,13	0.95	1 (7%)	7,16,16	2.73	2 (28%)
4	BTB	B	1330	-	13,13,13	1.11	1 (7%)	7,16,16	1.95	1 (14%)
4	BTB	B	1331	-	13,13,13	1.02	0	7,16,16	1.21	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BTB	C	1332	-	-	14/21/21/21	-
4	BTB	A	1332	-	-	12/21/21/21	-
4	BTB	D	1331	-	-	3/21/21/21	-
4	BTB	A	1331	-	-	15/21/21/21	-
4	BTB	C	1331	-	-	10/21/21/21	-
4	BTB	D	1330	-	-	6/21/21/21	-
4	BTB	A	1330	-	-	5/21/21/21	-
4	BTB	C	1330	-	-	2/21/21/21	-
4	BTB	B	1330	-	-	4/21/21/21	-
4	BTB	B	1331	-	-	17/21/21/21	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1330	BTB	C1-C2	-4.99	1.46	1.53
4	D	1330	BTB	C3-C2	-4.02	1.48	1.53
4	B	1330	BTB	C3-C2	-3.16	1.49	1.53
4	C	1330	BTB	C3-C2	-2.93	1.49	1.53

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	D	1330	BTB	O3-C3-C2	-7.37	91.26	111.44
4	C	1330	BTB	O3-C3-C2	-6.19	94.48	111.44
4	B	1330	BTB	O3-C3-C2	-4.94	97.91	111.44
4	A	1331	BTB	O4-C4-C2	3.76	121.73	111.44
4	A	1330	BTB	O1-C1-C2	-3.23	102.60	111.44

There are no chirality outliers.

5 of 88 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	1332	BTB	O1-C1-C2-C3
4	C	1332	BTB	O1-C1-C2-C4
4	C	1332	BTB	O1-C1-C2-N
4	C	1332	BTB	C1-C2-C4-O4
4	C	1332	BTB	C3-C2-C4-O4

There are no ring outliers.

10 monomers are involved in 64 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1332	BTB	3	0
4	A	1332	BTB	5	0
4	D	1331	BTB	5	0
4	A	1331	BTB	7	0
4	C	1331	BTB	13	0
4	D	1330	BTB	10	0
4	A	1330	BTB	7	0
4	C	1330	BTB	4	0
4	B	1330	BTB	5	0
4	B	1331	BTB	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/327 (100%)	-0.65	1 (0%) 94 93	19, 35, 60, 92	0
1	B	327/327 (100%)	-0.67	1 (0%) 94 93	20, 34, 59, 90	0
1	C	327/327 (100%)	-0.63	0 100 100	24, 37, 62, 85	0
1	D	327/327 (100%)	-0.74	1 (0%) 94 93	20, 32, 58, 88	0
All	All	1308/1308 (100%)	-0.67	3 (0%) 95 94	19, 35, 60, 92	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	73	ALA	3.4
1	D	74	GLY	2.5
1	B	72	GLN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	BTB	A	1332	14/14	0.74	0.20	57,78,89,91	0
4	BTB	C	1332	14/14	0.80	0.23	94,112,124,127	0
4	BTB	B	1331	14/14	0.82	0.20	40,64,72,72	0
4	BTB	D	1331	14/14	0.84	0.21	38,60,66,71	0
4	BTB	C	1331	14/14	0.88	0.21	43,66,72,74	0
4	BTB	A	1331	14/14	0.88	0.16	40,63,68,71	0
3	CL	C	1329	1/1	0.92	0.08	67,67,67,67	0
4	BTB	B	1330	14/14	0.94	0.13	40,46,53,54	0
4	BTB	C	1330	14/14	0.94	0.13	38,50,59,61	0
2	MG	C	1328	1/1	0.94	0.08	26,26,26,26	0
4	BTB	D	1330	14/14	0.95	0.15	30,43,53,56	0
4	BTB	A	1330	14/14	0.95	0.12	32,41,47,48	0
2	MG	A	1328	1/1	0.97	0.12	30,30,30,30	0
3	CL	D	1329	1/1	0.98	0.08	46,46,46,46	0
3	CL	A	1329	1/1	0.99	0.08	47,47,47,47	0
3	CL	B	1329	1/1	0.99	0.07	46,46,46,46	0
2	MG	D	1328	1/1	0.99	0.13	22,22,22,22	0
2	MG	B	1328	1/1	0.99	0.09	18,18,18,18	0

6.5 Other polymers ⓘ

There are no such residues in this entry.