



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 09:20 am BST

PDB ID : 6ASB  
Title : CXXC and PHD-type zinc finger regions of FBXL19 in complex with DNA  
Authors : Liu, K.; Tempel, W.; Walker, J.R.; Arrowsmith, C.H.; Bountra, C.; Edwards, A.M.; Min, J.; Structural Genomics Consortium (SGC)  
Deposited on : 2017-08-24  
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	12	<div><div></div><div>100%</div></div>
1	B	12	<div><div></div><div>100%</div></div>
1	D	12	<div><div></div><div>92%</div><div>8%</div></div>
1	E	12	<div><div></div><div>100%</div></div>
1	G	12	<div><div></div><div>92%</div><div>8%</div></div>
1	H	12	<div><div></div><div>83%</div><div>17%</div></div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	12	<div><div></div><div>92%</div><div>8%</div></div>
1	K	12	<div><div></div><div>100%</div><div></div></div>
2	C	123	<div><div></div><div>%</div><div>89%</div><div>7%</div><div></div><div></div></div>
2	F	123	<div><div></div><div>7%</div><div>58%</div><div>10%</div><div></div><div>31%</div><div></div></div>
2	I	123	<div><div></div><div>2%</div><div>91%</div><div>7%</div><div></div><div></div></div>
2	L	123	<div><div></div><div>3%</div><div>81%</div><div>8%</div><div>11%</div><div></div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	B	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	D	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	E	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	G	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	H	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	J	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			
1	K	12	Total	C	N	O	P	0	0	0
			243	116	46	70	11			

- Molecule 2 is a protein called F-box/LRR-repeat protein 19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	118	Total	C	N	O	S	0	0	1
			819	500	152	147	20			
2	F	85	Total	C	N	O	S	0	0	0
			584	351	118	97	18			
2	I	121	Total	C	N	O	S	0	0	0
			837	511	154	152	20			
2	L	110	Total	C	N	O	S	0	0	1
			710	430	133	129	18			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	I	4	Total 4	Zn 4	0	0
3	L	4	Total 4	Zn 4	0	0
3	C	4	Total 4	Zn 4	0	0
3	F	4	Total 4	Zn 4	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain A:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain B:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain D:  92% 8%



- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain E:  100%


There are no outlier residues recorded for this chain.

- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain G:  92% 8%



- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain H:  83% 17%



- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain J:  92% 8%




- Molecule 1: DNA (5'-D(\*GP\*CP\*CP\*AP\*AP\*CP\*GP\*TP\*TP\*GP\*GP\*C)-3')

Chain K:  100%

There are no outlier residues recorded for this chain.


- Molecule 2: F-box/LRR-repeat protein 19

Chain C:  89% 7% . .




- Molecule 2: F-box/LRR-repeat protein 19

Chain F:  7% 58% 10% . 31%




- Molecule 2: F-box/LRR-repeat protein 19

Chain I:  2% 91% 7% .



- Molecule 2: F-box/LRR-repeat protein 19

Chain L:  3% 81% 8% 11%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.73Å 64.28Å 67.32Å 103.53° 104.66° 99.40°	Depositor
Resolution (Å)	47.91 – 2.85 45.32 – 2.85	Depositor EDS
% Data completeness (in resolution range)	97.3 (47.91-2.85) 97.3 (45.32-2.85)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 2.86Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.213 , 0.251 0.223 , 0.258	Depositor DCC
$R_{free}$ test set	984 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.3	Xtriage
Anisotropy	0.555	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 61.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4910	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.15	0/272	0.98	0/418
1	B	1.12	0/272	0.95	0/418
1	D	1.23	0/272	0.97	0/418
1	E	1.23	0/272	0.91	0/418
1	G	1.08	0/272	0.98	0/418
1	H	1.00	0/272	0.95	0/418
1	J	1.25	0/272	0.97	0/418
1	K	1.10	0/272	0.90	0/418
2	C	0.52	0/832	0.72	0/1125
2	F	0.51	0/591	0.73	0/790
2	I	0.49	0/849	0.71	0/1145
2	L	0.50	0/719	0.66	0/971
All	All	0.84	0/5167	0.83	0/7375

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	243	0	136	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	243	0	136	0	0
1	D	243	0	136	1	0
1	E	243	0	136	0	0
1	G	243	0	136	1	0
1	H	243	0	136	2	0
1	J	243	0	136	1	0
1	K	243	0	136	0	0
2	C	819	0	717	2	0
2	F	584	0	489	7	0
2	I	837	0	736	4	0
2	L	710	0	557	4	0
3	C	4	0	0	0	0
3	F	4	0	0	0	0
3	I	4	0	0	0	0
3	L	4	0	0	0	0
All	All	4910	0	3587	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:49:GLU:HG2	2:F:72:CYS:HB2	1.78	0.65
1:G:12:DC:H3'	1:J:2:DC:H5'	1.81	0.61
2:F:111:LEU:O	2:F:141:ASN:HB3	2.09	0.53
1:H:10:DG:OP1	2:I:34:ARG:HD3	2.09	0.52
2:C:128:MET:HB3	2:C:131:ALA:HB3	1.92	0.51

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	116/123 (94%)	113 (97%)	2 (2%)	1 (1%)	17	43
2	F	77/123 (63%)	75 (97%)	2 (3%)	0	100	100
2	I	119/123 (97%)	117 (98%)	1 (1%)	1 (1%)	19	46
2	L	104/123 (85%)	102 (98%)	2 (2%)	0	100	100
All	All	416/492 (85%)	407 (98%)	7 (2%)	2 (0%)	29	57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	129	GLY
2	I	129	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	75/103 (73%)	70 (93%)	5 (7%)	16	39
2	F	53/103 (52%)	49 (92%)	4 (8%)	13	34
2	I	77/103 (75%)	74 (96%)	3 (4%)	32	63
2	L	56/103 (54%)	54 (96%)	2 (4%)	35	66
All	All	261/412 (63%)	247 (95%)	14 (5%)	22	49

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	71	SER
2	F	141	ASN
2	I	143	TRP
2	F	47	ARG
2	I	97	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	12/12 (100%)	-0.20	0 100 100	44, 60, 75, 78	0
1	B	12/12 (100%)	-0.26	0 100 100	48, 56, 67, 70	0
1	D	12/12 (100%)	-0.16	0 100 100	48, 62, 74, 75	0
1	E	12/12 (100%)	-0.30	0 100 100	57, 62, 74, 83	0
1	G	12/12 (100%)	-0.25	0 100 100	53, 75, 93, 96	0
1	H	12/12 (100%)	-0.19	0 100 100	64, 74, 87, 90	0
1	J	12/12 (100%)	-0.13	0 100 100	47, 62, 90, 97	0
1	K	12/12 (100%)	-0.13	0 100 100	61, 68, 90, 92	0
2	C	118/123 (95%)	-0.16	1 (0%) 86 85	39, 62, 88, 104	0
2	F	85/123 (69%)	0.50	9 (10%) 6 4	40, 65, 154, 185	0
2	I	121/123 (98%)	-0.01	3 (2%) 57 54	47, 72, 99, 109	0
2	L	110/123 (89%)	0.09	4 (3%) 42 37	40, 77, 136, 152	0
All	All	530/588 (90%)	0.03	17 (3%) 47 42	39, 66, 128, 185	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	125	CYS	4.8
2	F	88	CYS	3.7
2	F	115	THR	3.4
2	F	90	LEU	3.2
2	I	45	CYS	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	F	2003	1/1	0.60	0.06	188,188,188,188	0
3	ZN	F	2004	1/1	0.86	0.10	176,176,176,176	0
3	ZN	F	2001	1/1	0.94	0.08	69,69,69,69	0
3	ZN	C	2002	1/1	0.95	0.13	84,84,84,84	0
3	ZN	I	2001	1/1	0.96	0.07	100,100,100,100	0
3	ZN	L	2001	1/1	0.96	0.13	65,65,65,65	0
3	ZN	C	2003	1/1	0.97	0.05	96,96,96,96	0
3	ZN	L	2002	1/1	0.97	0.10	72,72,72,72	0
3	ZN	I	2003	1/1	0.98	0.06	69,69,69,69	0
3	ZN	L	2004	1/1	0.98	0.05	122,122,122,122	0
3	ZN	C	2004	1/1	0.98	0.04	79,79,79,79	0
3	ZN	F	2002	1/1	0.98	0.09	76,76,76,76	0
3	ZN	L	2003	1/1	0.98	0.04	162,162,162,162	0
3	ZN	C	2001	1/1	0.99	0.09	72,72,72,72	0
3	ZN	I	2004	1/1	0.99	0.06	78,78,78,78	0
3	ZN	I	2002	1/1	0.99	0.07	90,90,90,90	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.