



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 06:10 pm BST

PDB ID : 2AW5
Title : Crystal structure of a human malic enzyme
Authors : Papagrigoriou, E.; Berridge, G.; Smee, C.; Bray, J.; Arrowsmith, C.; Edwards, A.; Weigelt, J.; Sundstrom, M.; Oppermann, U.; Gileadi, O.; von Delft, F.; Structural Genomics Consortium (SGC)
Deposited on : 2005-08-31
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	575	<div> <div>%</div> <div> <div>76%</div> <div>14%</div> <div>8%</div> </div> </div>
1	B	575	<div> <div>16%</div> <div> <div>78%</div> <div>13%</div> <div>7%</div> </div> </div>
1	C	575	<div> <div>5%</div> <div> <div>74%</div> <div>17%</div> <div>7%</div> </div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 11920 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADP-dependent malic enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	531	Total	C	N	O	S	0	0	0
			4004	2567	675	744	18			
1	B	536	Total	C	N	O	S	0	0	0
			3919	2500	670	730	19			
1	C	533	Total	C	N	O	S	0	0	0
			3997	2556	675	748	18			

There are 69 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	CLONING ARTIFACT	UNP P48163
A	-9	HIS	-	CLONING ARTIFACT	UNP P48163
A	-8	HIS	-	CLONING ARTIFACT	UNP P48163
A	-7	HIS	-	CLONING ARTIFACT	UNP P48163
A	-6	HIS	-	CLONING ARTIFACT	UNP P48163
A	-5	HIS	-	CLONING ARTIFACT	UNP P48163
A	-4	HIS	-	CLONING ARTIFACT	UNP P48163
A	-3	SER	-	CLONING ARTIFACT	UNP P48163
A	-2	SER	-	CLONING ARTIFACT	UNP P48163
A	-1	GLY	-	CLONING ARTIFACT	UNP P48163
A	0	VAL	-	CLONING ARTIFACT	UNP P48163
A	1	ASP	-	CLONING ARTIFACT	UNP P48163
A	2	LEU	-	CLONING ARTIFACT	UNP P48163
A	3	GLY	-	CLONING ARTIFACT	UNP P48163
A	4	THR	-	CLONING ARTIFACT	UNP P48163
A	5	GLU	-	CLONING ARTIFACT	UNP P48163
A	6	ASN	-	CLONING ARTIFACT	UNP P48163
A	7	LEU	-	CLONING ARTIFACT	UNP P48163
A	8	TYR	-	CLONING ARTIFACT	UNP P48163
A	9	PHE	-	CLONING ARTIFACT	UNP P48163
A	10	GLN	-	CLONING ARTIFACT	UNP P48163
A	11	SER	-	CLONING ARTIFACT	UNP P48163
A	12	MET	-	CLONING ARTIFACT	UNP P48163

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	MET	-	CLONING ARTIFACT	UNP P48163
B	-9	HIS	-	CLONING ARTIFACT	UNP P48163
B	-8	HIS	-	CLONING ARTIFACT	UNP P48163
B	-7	HIS	-	CLONING ARTIFACT	UNP P48163
B	-6	HIS	-	CLONING ARTIFACT	UNP P48163
B	-5	HIS	-	CLONING ARTIFACT	UNP P48163
B	-4	HIS	-	CLONING ARTIFACT	UNP P48163
B	-3	SER	-	CLONING ARTIFACT	UNP P48163
B	-2	SER	-	CLONING ARTIFACT	UNP P48163
B	-1	GLY	-	CLONING ARTIFACT	UNP P48163
B	0	VAL	-	CLONING ARTIFACT	UNP P48163
B	1	ASP	-	CLONING ARTIFACT	UNP P48163
B	2	LEU	-	CLONING ARTIFACT	UNP P48163
B	3	GLY	-	CLONING ARTIFACT	UNP P48163
B	4	THR	-	CLONING ARTIFACT	UNP P48163
B	5	GLU	-	CLONING ARTIFACT	UNP P48163
B	6	ASN	-	CLONING ARTIFACT	UNP P48163
B	7	LEU	-	CLONING ARTIFACT	UNP P48163
B	8	TYR	-	CLONING ARTIFACT	UNP P48163
B	9	PHE	-	CLONING ARTIFACT	UNP P48163
B	10	GLN	-	CLONING ARTIFACT	UNP P48163
B	11	SER	-	CLONING ARTIFACT	UNP P48163
B	12	MET	-	CLONING ARTIFACT	UNP P48163
C	-10	MET	-	CLONING ARTIFACT	UNP P48163
C	-9	HIS	-	CLONING ARTIFACT	UNP P48163
C	-8	HIS	-	CLONING ARTIFACT	UNP P48163
C	-7	HIS	-	CLONING ARTIFACT	UNP P48163
C	-6	HIS	-	CLONING ARTIFACT	UNP P48163
C	-5	HIS	-	CLONING ARTIFACT	UNP P48163
C	-4	HIS	-	CLONING ARTIFACT	UNP P48163
C	-3	SER	-	CLONING ARTIFACT	UNP P48163
C	-2	SER	-	CLONING ARTIFACT	UNP P48163
C	-1	GLY	-	CLONING ARTIFACT	UNP P48163
C	0	VAL	-	CLONING ARTIFACT	UNP P48163
C	1	ASP	-	CLONING ARTIFACT	UNP P48163
C	2	LEU	-	CLONING ARTIFACT	UNP P48163
C	3	GLY	-	CLONING ARTIFACT	UNP P48163
C	4	THR	-	CLONING ARTIFACT	UNP P48163
C	5	GLU	-	CLONING ARTIFACT	UNP P48163
C	6	ASN	-	CLONING ARTIFACT	UNP P48163
C	7	LEU	-	CLONING ARTIFACT	UNP P48163
C	8	TYR	-	CLONING ARTIFACT	UNP P48163

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	9	PHE	-	CLONING ARTIFACT	UNP P48163
C	10	GLN	-	CLONING ARTIFACT	UNP P48163
C	11	SER	-	CLONING ARTIFACT	UNP P48163
C	12	MET	-	CLONING ARTIFACT	UNP P48163


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NADP-dependent malic enzyme




• Molecule 1: NADP-dependent malic enzyme



GLU
GLU
VAL
GLN

● Molecule 1: NADP-dependent malic enzyme



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	152.50Å 136.96Å 117.84Å 90.00° 121.75° 90.00°	Depositor
Resolution (Å)	37.30 – 2.50 47.40 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.7 (37.30-2.50) 97.7 (47.40-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.207 , 0.256 0.250 , 0.296	Depositor DCC
R_{free} test set	3520 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	52.2	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11920	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/4080	0.74	2/5546 (0.0%)
1	B	0.56	0/3993	0.70	0/5440
1	C	0.57	0/4074	0.71	1/5541 (0.0%)
All	All	0.58	0/12147	0.72	3/16527 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	118	ARG	NE-CZ-NH1	5.65	123.13	120.30
1	A	215	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	C	276	VAL	CB-CA-C	-5.11	101.70	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4004	0	3879	57	0
1	B	3919	0	3687	53	0
1	C	3997	0	3832	62	0
All	All	11920	0	11398	169	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 169 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:124:PHE:CD2	1:C:192:ILE:HD11	1.90	1.06
1:B:150:VAL:HG21	1:B:228:PHE:CZ	2.13	0.83
1:B:124:PHE:CD2	1:B:192:ILE:HD11	2.20	0.77
1:B:16:TYR:O	1:B:19:THR:HB	1.87	0.74
1:A:150:VAL:HG22	1:A:191:VAL:HB	1.69	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/575 (91%)	508 (97%)	14 (3%)	3 (1%)	25	43
1	B	532/575 (92%)	517 (97%)	14 (3%)	1 (0%)	47	68
1	C	529/575 (92%)	502 (95%)	27 (5%)	0	100	100
All	All	1586/1725 (92%)	1527 (96%)	55 (4%)	4 (0%)	41	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	304	GLU
1	A	128	HIS
1	B	304	GLU
1	A	19	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/492 (82%)	376 (94%)	25 (6%)	18	35
1	B	372/492 (76%)	350 (94%)	22 (6%)	19	37
1	C	397/492 (81%)	372 (94%)	25 (6%)	18	34
All	All	1170/1476 (79%)	1098 (94%)	72 (6%)	18	35

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	142	GLU
1	B	453	ASN
1	C	361	MET
1	B	192	ILE
1	B	260	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	49	ASN
1	B	61	ASN
1	C	49	ASN
1	A	453	ASN
1	A	488	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	531/575 (92%)	0.15	5 (0%) 84 86	25, 60, 79, 87	0
1	B	536/575 (93%)	0.82	90 (16%) 1 1	25, 63, 84, 95	0
1	C	533/575 (92%)	0.48	27 (5%) 28 29	25, 63, 79, 93	0
All	All	1600/1725 (92%)	0.48	122 (7%) 13 14	25, 62, 81, 95	0

The worst 5 of 122 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	354	PHE	9.4
1	B	358	HIS	7.8
1	B	355	ALA	7.4
1	B	380	ALA	7.3
1	B	298	LEU	6.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.