



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 09:20 pm BST

PDB ID : 3AWV
Title : Crystal structure of Streptomyces tyrosinase in a complex with caddie soaked in a Cu(II)-containing solution for 80 hr: occupancy of CuA is low
Authors : Matoba, Y.; Sugiyama, M.
Deposited on : 2011-03-26
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> % </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 82% 14% • • </div> </div>
2	B	134	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 5% </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 50% 8% 42% </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3192 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	8	0
			2240	1407	413	416	4			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	SER	PHE	SEE REMARK 999	UNP Q83WS2
A	274	LEU	-	EXPRESSION TAG	UNP Q83WS2
A	275	GLU	-	EXPRESSION TAG	UNP Q83WS2
A	276	HIS	-	EXPRESSION TAG	UNP Q83WS2
A	277	HIS	-	EXPRESSION TAG	UNP Q83WS2
A	278	HIS	-	EXPRESSION TAG	UNP Q83WS2
A	279	HIS	-	EXPRESSION TAG	UNP Q83WS2
A	280	HIS	-	EXPRESSION TAG	UNP Q83WS2
A	281	HIS	-	EXPRESSION TAG	UNP Q83WS2

- Molecule 2 is a protein called MelC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	78	Total	C	N	O	S	0	3	0
			607	383	110	112	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	127	LEU	-	EXPRESSION TAG	UNP Q83WS1
B	128	GLU	-	EXPRESSION TAG	UNP Q83WS1
B	129	HIS	-	EXPRESSION TAG	UNP Q83WS1
B	130	HIS	-	EXPRESSION TAG	UNP Q83WS1
B	131	HIS	-	EXPRESSION TAG	UNP Q83WS1
B	132	HIS	-	EXPRESSION TAG	UNP Q83WS1
B	133	HIS	-	EXPRESSION TAG	UNP Q83WS1

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	134	HIS	-	EXPRESSION TAG	UNP Q83WS1

- Molecule 3 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cu 2 2	0	1
3	A	2	Total Cu 2 2	0	2

- Molecule 4 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total N O 4 1 3	0	0
4	A	1	Total N O 4 1 3	0	0
4	A	1	Total N O 4 1 3	0	0
4	A	1	Total N O 4 1 3	0	0
4	B	1	Total N O 4 1 3	0	0
4	B	1	Total N O 4 1 3	0	1


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	235	Total 235	O 235	0	0
5	B	82	Total 82	O 82	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosinase



• Molecule 2: MelC



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	65.09Å 97.85Å 55.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.40 24.56 – 1.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.40) 99.9 (24.56-1.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 1.40Å)	Xtriage
Refinement program	CNS, SHELXL-97	Depositor
R, R_{free}	0.176 , 0.218 0.166 , 0.202	Depositor DCC
R_{free} test set	3534 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3192	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/2324	1.36	27/3171 (0.9%)
2	B	0.58	0/636	1.49	9/864 (1.0%)
All	All	0.58	0/2960	1.39	36/4035 (0.9%)

There are no bond length outliers.

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	105	ARG	CD-NE-CZ	16.11	146.15	123.60
2	B	105	ARG	NE-CZ-NH1	-15.12	112.74	120.30
2	B	105	ARG	NE-CZ-NH2	10.81	125.70	120.30
1	A	178	ARG	NE-CZ-NH2	-9.96	115.32	120.30
1	A	178	ARG	NE-CZ-NH1	9.05	124.83	120.30
1	A	136[A]	ARG	NE-CZ-NH2	-8.64	115.98	120.30
1	A	136[B]	ARG	NE-CZ-NH2	-8.64	115.98	120.30
1	A	136[A]	ARG	NE-CZ-NH1	8.62	124.61	120.30
1	A	136[B]	ARG	NE-CZ-NH1	8.62	124.61	120.30
1	A	92	ARG	NE-CZ-NH2	-8.32	116.14	120.30
1	A	16	ARG	CD-NE-CZ	7.78	134.50	123.60
1	A	114	ARG	NE-CZ-NH2	7.74	124.17	120.30
1	A	229	ARG	NE-CZ-NH2	-7.67	116.47	120.30
1	A	36	ARG	NE-CZ-NH1	-7.19	116.70	120.30
1	A	16	ARG	NE-CZ-NH1	7.16	123.88	120.30
2	B	72	TYR	CB-CG-CD2	6.84	125.10	121.00
1	A	235	TYR	CB-CG-CD2	6.70	125.02	121.00
1	A	140	ARG	NE-CZ-NH1	-6.41	117.09	120.30
1	A	258	ARG	NE-CZ-NH2	6.39	123.50	120.30
1	A	55	ARG	NE-CZ-NH1	6.30	123.45	120.30
1	A	87	ASP	CB-CG-OD2	-6.14	112.78	118.30
1	A	229	ARG	NE-CZ-NH1	6.09	123.34	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	45	PHE	CB-CG-CD2	-6.04	116.57	120.80
1	A	71	GLU	OE1-CD-OE2	5.86	130.33	123.30
1	A	218	TYR	CB-CG-CD1	-5.84	117.49	121.00
1	A	272	ASP	CB-CG-OD2	5.76	123.48	118.30
1	A	92	ARG	NH1-CZ-NH2	5.58	125.53	119.40
2	B	85	ARG	NE-CZ-NH2	-5.51	117.54	120.30
2	B	45	PHE	CB-CG-CD1	5.35	124.55	120.80
1	A	140	ARG	NH1-CZ-NH2	5.29	125.22	119.40
2	B	57	ARG	NE-CZ-NH1	-5.25	117.67	120.30
2	B	85	ARG	NE-CZ-NH1	5.25	122.92	120.30
1	A	140	ARG	NE-CZ-NH2	-5.23	117.69	120.30
1	A	103	PHE	CB-CG-CD1	5.21	124.45	120.80
1	A	103	PHE	CB-CG-CD2	-5.13	117.21	120.80
1	A	112	ASP	CB-CG-OD1	5.11	122.90	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2240	0	2112	26	0
2	B	607	0	574	6	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	16	0	0	0	0
4	B	8	0	0	0	0
5	A	235	0	0	5	0
5	B	82	0	0	0	0
All	All	3192	0	2686	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:VAL:HG22	5:A:798:HOH:O	1.94	0.66
1:A:46:THR:HG22	2:B:84[B]:MET:HE1	1.81	0.61
1:A:136[A]:ARG:HD2	1:A:138:TYR:OH	2.04	0.58
1:A:46:THR:HG22	2:B:84[B]:MET:CE	2.35	0.56
1:A:111:THR:HG21	5:A:823:HOH:O	2.05	0.55
1:A:185:ARG:NH2	2:B:88:ASP:HB3	2.22	0.55
1:A:135[B]:SER:HB2	5:A:757:HOH:O	2.10	0.50
1:A:248:ASN:HB3	5:A:705:HOH:O	2.12	0.49
1:A:242:PRO:O	1:A:243:ASP:HB2	2.13	0.48
1:A:185:ARG:HH22	2:B:88:ASP:HB3	1.79	0.47
1:A:16:ARG:HG2	1:A:16:ARG:HH11	1.81	0.46
1:A:16:ARG:NH1	5:A:795:HOH:O	2.48	0.45
1:A:153:ARG:NH2	1:A:157:GLU:OE1	2.50	0.45
1:A:153:ARG:O	1:A:157:GLU:HG2	2.17	0.45
1:A:39:ASN:OD1	1:A:204[B]:GLY:O	2.36	0.44
1:A:204[B]:GLY:HA3	2:B:94:VAL:HG23	2.00	0.44
1:A:136[B]:ARG:NH2	2:B:114:GLU:OE1	2.50	0.43
1:A:42:ILE:HD13	1:A:54[A]:HIS:CE1	2.54	0.43
1:A:42:ILE:HG12	1:A:54[B]:HIS:CE1	2.54	0.42
1:A:56:SER:HB2	1:A:57:PRO:HD2	2.02	0.41
1:A:56:SER:O	1:A:178:ARG:HD3	2.21	0.41
1:A:153:ARG:HH12	1:A:156:VAL:HG11	1.86	0.41
1:A:113:GLY:O	1:A:140:ARG:HA	2.22	0.40
1:A:228:ARG:HD3	1:A:228:ARG:HA	1.80	0.40
1:A:65[A]:ARG:HG3	1:A:254:TRP:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	279/281 (99%)	269 (96%)	10 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	77/134 (58%)	74 (96%)	3 (4%)	0	100	100
All	All	356/415 (86%)	343 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/240 (100%)	235 (98%)	4 (2%)	60	31
2	B	63/94 (67%)	61 (97%)	2 (3%)	39	9
All	All	302/334 (90%)	296 (98%)	6 (2%)	53	23

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	88	TRP
1	A	103	PHE
1	A	190	HIS
2	B	67	GLU
2	B	68	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
2	B	80	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NO3	A	507	-	1,3,3	0.54	0	0,3,3	0.00	-
4	NO3	B	505	-	1,3,3	0.74	0	0,3,3	0.00	-
4	NO3	A	504	-	1,3,3	0.24	0	0,3,3	0.00	-
4	NO3	A	506	-	1,3,3	0.56	0	0,3,3	0.00	-
4	NO3	B	508[A]	-	1,3,3	0.61	0	0,3,3	0.00	-
4	NO3	A	503	-	1,3,3	0.84	0	0,3,3	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/281 (97%)	-0.17	4 (1%) 73 72	9, 14, 26, 51	0
2	B	78/134 (58%)	0.27	7 (8%) 9 8	9, 16, 42, 57	0
All	All	351/415 (84%)	-0.07	11 (3%) 49 48	9, 15, 30, 57	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	69	GLY	14.4
2	B	70	GLY	8.7
2	B	40	ALA	8.1
2	B	123	PHE	6.7
2	B	66	HIS	5.7
2	B	68	HIS	4.0
1	A	148	ALA	3.7
1	A	95	ARG	3.4
1	A	274	LEU	2.8
2	B	67	GLU	2.3
1	A	94	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NO3	A	506	4/4	0.87	0.17	21,33,36,38	0
4	NO3	A	507	4/4	0.89	0.14	28,37,40,42	0
4	NO3	B	505	4/4	0.94	0.15	21,22,27,32	0
4	NO3	A	503	4/4	0.94	0.11	15,16,26,27	0
4	NO3	A	504	4/4	0.95	0.08	20,23,30,37	0
4	NO3	B	508[A]	4/4	0.96	0.07	18,23,24,31	4
3	CU	B	502[A]	1/1	0.99	0.08	21,21,21,21	1
3	CU	A	500[A]	1/1	0.99	0.06	17,17,17,17	1
3	CU	B	502[B]	1/1	0.99	0.08	24,24,24,24	1
3	CU	A	501[A]	1/1	1.00	0.03	13,13,13,13	1

6.5 Other polymers

There are no such residues in this entry.