



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 12:49 pm BST

PDB ID : 1AXT  
Title : IMMUNE VERSUS NATURAL SELECTION: ANTIBODY ALDOLASES  
WITH THE RATES OF NATURAL ENZYMES  
Authors : Heine, A.; Wilson, I.A.  
Deposited on : 1997-10-20  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	216	
2	H	218	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3598 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IMMUNOGLOBULIN IGG2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	216	Total	C	N	O	S	0	0	0
			1674	1048	283	337	6			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	28	TYR	ASN	CONFLICT	PIR S16112
L	32	PHE	TYR	CONFLICT	PIR S16112
L	34	ASN	TYR	CONFLICT	PIR S16112
L	40	SER	PRO	CONFLICT	PIR S16112
L	46	LEU	PRO	CONFLICT	PIR S16112
L	50	LYS	ARG	CONFLICT	PIR S16112
L	89	SER	PHE	CONFLICT	PIR S16112
L	103	LYS	ARG	CONFLICT	PIR S16112

- Molecule 2 is a protein called IMMUNOGLOBULIN IGG2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	218	Total	C	N	O	S	42	0	0
			1676	1066	274	328	8			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	101	Total	O	0	0
			101	101		
3	H	147	Total	O	0	0
			147	147		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: IMMUNOGLOBULIN IGG2A



#### • Molecule 2: IMMUNOGLOBULIN IGG2A



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.50 Å 65.30 Å 132.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.15	Depositor
% Data completeness (in resolution range)	96.1 (10.00-2.15)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
Refinement program	SHELXL-96	Depositor
R, $R_{free}$	0.214 , 0.317	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3598	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.33	0/1712	0.96	2/2320 (0.1%)
2	H	0.35	0/1719	1.00	3/2339 (0.1%)
All	All	0.34	0/3431	0.98	5/4659 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	108	ARG	NE-CZ-NH1	7.04	123.82	120.30
1	L	77	ARG	NE-CZ-NH1	5.96	123.28	120.30
2	H	228	ARG	NE-CZ-NH1	5.34	122.97	120.30
2	H	149	PRO	C-N-CA	5.02	134.25	121.70
2	H	191	VAL	CB-CA-C	-5.01	101.89	111.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1674	0	1620	31	0
2	H	1676	0	1647	25	0
3	H	147	0	0	8	0
3	L	101	0	0	5	0
All	All	3598	0	3267	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:115:VAL:O	1:L:207:LYS:HD3	1.83	0.78
2:H:127:VAL:HG22	3:H:309:HOH:O	1.83	0.76
2:H:126:PRO:HB3	3:H:306:HOH:O	1.88	0.73
1:L:2:LEU:O	1:L:97:THR:HG21	1.88	0.72
2:H:203:GLN:HG2	3:H:229:HOH:O	1.89	0.71
1:L:125:LEU:O	1:L:183:LYS:HD2	1.91	0.71
1:L:39:LYS:HB2	1:L:42:GLN:NE2	2.14	0.63
1:L:39:LYS:O	1:L:42:GLN:HB2	1.99	0.62
1:L:186:TYR:CZ	1:L:211:ARG:HD2	2.35	0.62
1:L:17:ASP:O	1:L:78:VAL:HG13	2.01	0.60
2:H:153:THR:OG1	2:H:211:ALA:HB3	2.03	0.59
1:L:142:LYS:HG2	3:L:299:HOH:O	2.03	0.59
2:H:52(B):LYS:HG3	2:H:55:TYR:CZ	2.38	0.57
1:L:190:ASN:OD1	1:L:210:ASN:HB3	2.03	0.57
1:L:27(E):SER:OG	1:L:30:ASN:HB2	2.04	0.57
2:H:150:GLU:HB2	3:H:262:HOH:O	2.08	0.53
2:H:157:TRP:CZ2	2:H:191:VAL:HG22	2.44	0.53
1:L:197:THR:HA	3:L:233:HOH:O	2.09	0.52
2:H:52(B):LYS:HG3	2:H:55:TYR:OH	2.11	0.51
1:L:149:LYS:HE3	1:L:195:GLU:OE1	2.10	0.51
2:H:157:TRP:CE2	2:H:191:VAL:HG22	2.47	0.50
1:L:75:ILE:CD1	1:L:78:VAL:HG12	2.40	0.50
2:H:5:GLU:HG2	3:H:305:HOH:O	2.10	0.50
1:L:14:SER:N	1:L:107:LYS:HZ3	2.10	0.49
2:H:2:VAL:HA	2:H:25:SER:O	2.14	0.48
1:L:14:SER:O	1:L:17:ASP:HB2	2.14	0.47
1:L:25:SER:O	1:L:69:THR:OG1	2.30	0.47
1:L:147:LYS:N	1:L:147:LYS:HD3	2.29	0.47
2:H:23:VAL:HG13	3:H:249:HOH:O	2.15	0.46
2:H:171:VAL:HG22	2:H:191:VAL:HG13	1.98	0.46
2:H:17:SER:HA	2:H:82:MET:O	2.16	0.46
1:L:75:ILE:HG12	1:L:75:ILE:O	2.12	0.46
1:L:150:ILE:HG21	3:L:225:HOH:O	2.15	0.46
2:H:53:ASP:O	2:H:54:ASN:HB2	2.15	0.46
2:H:82(A):ASN:O	2:H:82(B):SER:HB2	2.16	0.46
2:H:58:HIS:HD2	3:H:282:HOH:O	1.99	0.46
1:L:191:SER:HB2	1:L:210:ASN:OD1	2.16	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:170:ASP:O	1:L:171:SER:HB2	2.16	0.45
1:L:2:LEU:HD23	1:L:27:GLN:NE2	2.32	0.45
1:L:140:TYR:CG	1:L:141:PRO:HA	2.53	0.44
2:H:137:SER:N	2:H:195:SER:OG	2.50	0.44
1:L:54:ARG:NH1	3:L:268:HOH:O	2.50	0.43
2:H:2:VAL:CG2	2:H:27:LEU:HD13	2.48	0.43
2:H:147:TYR:CE1	2:H:152:VAL:HG13	2.53	0.43
2:H:199:TRP:CG	2:H:200:PRO:HA	2.54	0.43
1:L:2:LEU:HD22	1:L:26:SER:OG	2.18	0.43
1:L:165:ASP:HA	3:L:304:HOH:O	2.18	0.42
2:H:147:TYR:HE1	2:H:150:GLU:O	2.01	0.42
1:L:190:ASN:O	1:L:210:ASN:HA	2.19	0.42
1:L:61:ARG:HH22	1:L:82:ASP:CG	2.23	0.42
2:H:108:LEU:HD13	3:H:351:HOH:O	2.20	0.42
1:L:183:LYS:HE2	1:L:187:GLU:OE2	2.20	0.42
1:L:191:SER:HA	1:L:210:ASN:OD1	2.20	0.42
2:H:23:VAL:O	2:H:23:VAL:HG23	2.20	0.41
1:L:12:PRO:HA	1:L:105:GLU:O	2.21	0.41
2:H:178:LEU:HB2	2:H:185:TYR:CE1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	214/216 (99%)	208 (97%)	6 (3%)	0	100	100
2	H	216/218 (99%)	207 (96%)	7 (3%)	2 (1%)	17	11
All	All	430/434 (99%)	415 (96%)	13 (3%)	2 (0%)	29	22

All (2) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
2	H	133	THR
2	H	136	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	193/193 (100%)	165 (86%)	28 (14%)	3	1
2	H	191/191 (100%)	167 (87%)	24 (13%)	4	1
All	All	384/384 (100%)	332 (86%)	52 (14%)	4	1

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	2	LEU
1	L	6	GLN
1	L	9	LEU
1	L	17	ASP
1	L	18	GLN
1	L	24	ARG
1	L	27	GLN
1	L	33	LEU
1	L	56	SER
1	L	69	THR
1	L	75	ILE
1	L	76	SER
1	L	81	GLU
1	L	83	LEU
1	L	104	LEU
1	L	122	SER
1	L	143	ASP
1	L	149	LYS
1	L	155	ARG
1	L	160	LEU
1	L	184	ASP
1	L	191	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	L	202	THR
1	L	203	SER
1	L	205	ILE
1	L	207	LYS
1	L	208	SER
1	L	211	ARG
2	H	3	LYS
2	H	13	GLN
2	H	24	VAL
2	H	27	LEU
2	H	52(B)	LYS
2	H	61	GLU
2	H	66	LYS
2	H	71	ARG
2	H	78	LEU
2	H	82(A)	ASN
2	H	83	ARG
2	H	93	LYS
2	H	108	LEU
2	H	128	CYS
2	H	130	ASP
2	H	134	THR
2	H	136	SER
2	H	137	SER
2	H	150	GLU
2	H	191	VAL
2	H	195	SER
2	H	202	SER
2	H	204	SER
2	H	221	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	27	GLN
1	L	34	ASN
1	L	42	GLN
1	L	124	GLN
1	L	166	GLN
2	H	82(A)	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.