



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 11:46 pm BST

PDB ID : 4AXC  
Title : Inositol 1,3,4,5,6-pentakisphosphate 2-kinase apo form  
Authors : I Banos-Sanz, J.; Sanz-Aparicio, J.; Gonzalez, B.  
Deposited on : 2012-06-12  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	<div> <div>5%</div> <div>65%</div> <div>18%</div> <div>•</div> <div>14%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INOSITOL-PENTAKISPHOSPHATE 2-KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	1	0
			3151	2007	535	596	13			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q93YN9
A	-3	GLU	-	expression tag	UNP Q93YN9
A	-2	PHE	-	expression tag	UNP Q93YN9
A	-1	GLU	-	expression tag	UNP Q93YN9
A	0	LEU	-	expression tag	UNP Q93YN9
A	54	SER	ALA	conflict	UNP Q93YN9
A	90	GLN	LYS	conflict	UNP Q93YN9
A	129	ALA	TRP	conflict	UNP Q93YN9
A	157	THR	SER	conflict	UNP Q93YN9
A	204	ILE	ASN	conflict	UNP Q93YN9
A	224	ARG	SER	conflict	UNP Q93YN9
A	321	CYS	SER	conflict	UNP Q93YN9
A	325	ILE	LEU	conflict	UNP Q93YN9
A	337	ARG	LYS	conflict	UNP Q93YN9

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	74	Total	O	0	0
			74	74		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.80Å 68.23Å 66.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	66.00 – 2.25 47.44 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.4 (66.00-2.25) 99.5 (47.44-2.25)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.242 , 0.278 0.242 , 0.276	Depositor DCC
$R_{free}$ test set	1192 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.4	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 37.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	2/3207 (0.1%)	0.66	2/4322 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	13	TRP	CD2-CE2	5.42	1.47	1.41
1	A	381	TRP	CD2-CE2	5.39	1.47	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	LEU	CA-CB-CG	6.61	130.51	115.30
1	A	248	LEU	CA-CB-CG	5.43	127.79	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3151	0	3174	60	0
2	A	1	0	0	0	0
3	A	20	0	0	0	0
4	A	6	0	8	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	74	0	0	1	0
All	All	3252	0	3182	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:395:THR:HG21	1:A:397:GLN:HE21	1.44	0.83
1:A:254:GLY:O	1:A:257:THR:HG22	1.92	0.69
1:A:66:GLN:HG3	1:A:76:ILE:HB	1.75	0.68
1:A:18:GLU:OE2	1:A:126:ARG:HD3	1.95	0.67
1:A:67:HIS:O	1:A:70:ARG:HG2	1.95	0.65
1:A:6:GLU:HA	1:A:114:PHE:HB2	1.78	0.65
1:A:15:TYR:HA	1:A:25:LEU:HD12	1.79	0.65
1:A:121:LYS:HZ3	1:A:125:GLN:HG3	1.61	0.65
1:A:23:LEU:HD22	1:A:43:LYS:HG2	1.79	0.64
1:A:277:GLN:H	1:A:397:GLN:HE22	1.47	0.63
1:A:16:ARG:NH2	1:A:148:ASP:OD1	2.34	0.59
1:A:395:THR:HG21	1:A:397:GLN:NE2	2.17	0.58
1:A:122:VAL:O	1:A:126:ARG:HG3	2.03	0.58
1:A:277:GLN:H	1:A:397:GLN:NE2	2.04	0.56
1:A:73:ASN:O	1:A:76:ILE:HG13	2.07	0.55
1:A:110:VAL:CG2	1:A:114:PHE:CD2	2.90	0.54
1:A:107:ARG:HB3	1:A:107:ARG:CZ	2.38	0.54
1:A:121:LYS:NZ	1:A:125:GLN:HG3	2.22	0.54
1:A:120:LYS:HA	1:A:120:LYS:HE2	1.90	0.53
1:A:238:ASN:H	1:A:238:ASN:HD22	1.57	0.53
1:A:254:GLY:HA3	1:A:257:THR:HG22	1.92	0.52
1:A:97:LEU:HG	1:A:307:ILE:CD1	2.39	0.52
1:A:45:ARG:O	1:A:46:ARG:HB2	2.11	0.51
1:A:25:LEU:HD23	1:A:41:ILE:HD12	1.93	0.51
1:A:164:SER:HB3	1:A:243:PHE:HB2	1.94	0.49
1:A:422:LEU:O	1:A:426:ILE:HG12	2.12	0.49
1:A:295:VAL:HG13	1:A:301:LEU:HD22	1.94	0.48
1:A:15:TYR:HA	1:A:25:LEU:CD1	2.44	0.48
1:A:61:LEU:HD12	1:A:76:ILE:HA	1.96	0.47
1:A:68:LEU:HD22	1:A:413:LEU:HD13	1.94	0.47
1:A:119:ASP:O	1:A:123:THR:HG22	2.15	0.47
1:A:190:VAL:HB	1:A:195:MET:HE2	1.97	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:HIS:HD2	1:A:303:ARG:NH2	2.14	0.46
1:A:374:SER:O	1:A:401:TYR:HA	2.16	0.46
1:A:97:LEU:O	1:A:303:ARG:NH2	2.49	0.45
1:A:261:SER:HB2	1:A:262:PRO:HD2	1.98	0.45
1:A:14:ILE:HA	1:A:121:LYS:HE2	1.99	0.45
1:A:272:LEU:HD22	1:A:276:ILE:HD13	1.99	0.45
1:A:5:LEU:HB2	1:A:110:VAL:HG23	1.99	0.44
1:A:25:LEU:CD2	1:A:41:ILE:HD12	2.48	0.44
1:A:68:LEU:HA	1:A:354:LEU:HD22	1.99	0.44
1:A:172:GLY:HA3	1:A:218:PHE:CD2	2.52	0.44
1:A:238:ASN:N	1:A:238:ASN:HD22	2.16	0.44
1:A:119:ASP:CG	1:A:136:VAL:HG22	2.38	0.43
1:A:25:LEU:HB2	1:A:39:ILE:HG22	1.99	0.43
1:A:393:LYS:HE2	1:A:394:PRO:HG3	1.99	0.43
1:A:72[A]:ASN:N	1:A:72[A]:ASN:OD1	2.52	0.43
1:A:68:LEU:HD23	1:A:69:TRP:NE1	2.34	0.42
1:A:132:ASN:HD22	1:A:132:ASN:HA	1.70	0.42
1:A:254:GLY:C	1:A:256:SER:N	2.70	0.42
1:A:97:LEU:O	1:A:303:ARG:CZ	2.68	0.42
1:A:254:GLY:O	1:A:257:THR:CG2	2.65	0.42
1:A:15:TYR:CE2	1:A:126:ARG:HG2	2.55	0.41
1:A:300:VAL:HG22	1:A:405:PHE:CZ	2.55	0.41
1:A:34:PHE:HE1	1:A:145:ILE:HD13	1.84	0.41
1:A:14:ILE:HA	1:A:121:LYS:CE	2.51	0.41
1:A:162:CYS:HA	1:A:375:PHE:O	2.21	0.40
1:A:107:ARG:HB2	5:A:2009:HOH:O	2.20	0.40
1:A:97:LEU:HG	1:A:307:ILE:HD12	2.03	0.40
1:A:328:GLN:HG2	1:A:329:PRO:HD2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/456 (84%)	369 (96%)	14 (4%)	2 (0%)	29	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	70	ARG
1	A	254	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/402 (88%)	315 (89%)	38 (11%)	6	4

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	MET
1	A	4	ILE
1	A	6	GLU
1	A	16	ARG
1	A	23	LEU
1	A	30	SER
1	A	33	LEU
1	A	46	ARG
1	A	63	SER
1	A	70	ARG
1	A	72[A]	ASN
1	A	72[B]	ASN
1	A	96	LEU
1	A	123	THR
1	A	132	ASN
1	A	151	LEU
1	A	164	SER
1	A	174	LEU
1	A	176	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	186	LEU
1	A	188	THR
1	A	189	SER
1	A	198	LEU
1	A	207	SER
1	A	238	ASN
1	A	244	LEU
1	A	248	LEU
1	A	253	SER
1	A	273	LYS
1	A	300	VAL
1	A	301	LEU
1	A	310	LEU
1	A	350	LEU
1	A	351	ASP
1	A	379	ASN
1	A	393	LYS
1	A	397	GLN
1	A	414	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
1	A	132	ASN
1	A	147	ASN
1	A	238	ASN
1	A	397	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	1438	-	4,4,4	0.84	0	6,6,6	0.30	0
4	GOL	A	1440	-	5,5,5	0.49	0	5,5,5	0.31	0
3	SO4	A	1437	-	4,4,4	1.13	0	6,6,6	0.70	0
3	SO4	A	1439	-	4,4,4	0.43	0	6,6,6	0.19	0
3	SO4	A	1436	-	4,4,4	0.76	0	6,6,6	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1440	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1440	GOL	C1-C2-C3-O3
4	A	1440	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	394/456 (86%)	0.46	23 (5%)	23 25	32, 54, 79, 96	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	253	SER	5.1
1	A	394	PRO	4.0
1	A	381	TRP	4.0
1	A	73	ASN	3.8
1	A	124	LYS	3.2
1	A	4	ILE	3.1
1	A	3	MET	3.0
1	A	182	LYS	3.0
1	A	329	PRO	2.9
1	A	82	GLU	2.8
1	A	421	LYS	2.7
1	A	8	LYS	2.7
1	A	350	LEU	2.5
1	A	303	ARG	2.5
1	A	74	GLU	2.4
1	A	75	LEU	2.4
1	A	393	LYS	2.2
1	A	378	ARG	2.1
1	A	28	ALA	2.1
1	A	395	THR	2.1
1	A	432	ARG	2.0
1	A	83	VAL	2.0
1	A	81	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	1440	6/6	0.53	0.25	63,70,72,72	0
3	SO4	A	1439	5/5	0.86	0.20	73,76,79,85	0
3	SO4	A	1436	5/5	0.88	0.20	63,63,67,69	0
2	ZN	A	1435	1/1	0.96	0.06	56,56,56,56	0
3	SO4	A	1437	5/5	0.97	0.13	54,59,63,64	0
3	SO4	A	1438	5/5	0.98	0.10	61,65,73,74	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.